Lawrence Berkeley National Laboratory

Recent Work

Title

Neutron Deficient Isotopes of Iodine

Permalink

<https://escholarship.org/uc/item/97t218d4>

Authors

Marquez, Luis

Perlman, I.

Publication Date

1949-12-29

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNCLASSIFIED

UCRL-555
Unclassified Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

NEUTRON DEFICIENT ISOTOPES OF IODINE

Luis Marquez and I. Perlman

December 29, 1949

Berkeley, California

-2-

INSTALLATIONNo. of Copies

Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brookhaven National Laboratory	8
Bureau of Medicine and Surgery	1
Bureau of Ships	1
Carbide & Carbon Chemicals Corp. (K-25)	4
Carbide & Carbon Chemicals Corp. (Y-12)	4
Chicago Operations Office	1
Cleveland Area Office	1
Columbia University (Dunning)	1
Columbia University (Failla)	1
Dow Chemical Company	1
General Electric Company, Richland	6
Hanford Operations Office	1
Idaho Operations Office	1
Iowa State College	2
Kansas City	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos	3
Mallinckrodt Chemical Works	1
Massachusetts Institute of Technology (Gaudin)	1
Massachusetts Institute of Technology (Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	2
National Bureau of Standards	2
Naval Radiological Defense Laboratory	2
NEPA Project	2
New Brunswick Laboratory	1
New York Operations Office	5
North American Aviation, Inc.	1
Oak Ridge National Laboratory	8
Patent Advisor, Washington	1
Rand Corporation	1
Sandia Base	1
Sylvania Electric Products, Inc.	1
Technical Information Branch, ORE	15
U. S. Public Health Service	1
UCLA Medical Research Laboratory (Warren)	1
University of California Radiation Laboratory	5
University of Rochester	2
University of Washington	1
Western Reserve University (Friedell)	2
Westinghouse Electric Corp.	4
Total	119

Information Division
 Radiation Laboratory
 Univ. of California
 Berkeley, California

NEUTRON DEFICIENT ISOTOPES OF IODINE

Luis Marquez and I. Perlman
Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

ABSTRACT

The iodine isotopes I^{123} , I^{122} , I^{121} , and $I^{120}(\?)$ have been made by bombarding antimony with α -particles. Their characteristics are summarized and evidence for mass assignments presented.

NEUTRON DEFICIENT ISOTOPES OF IODINE

Luis Marquez and I. Perlman
Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

During the course of some investigations of abnormal change in nuclear charge following irradiation with high-energy particles, some of the nuclides encountered proved to be light isotopes of iodine which had not been reported previously. The main study will be reported at a later date while the present communication describes some work done to characterize and identify the iodine isotopes. The mass number region of interest is that below I^{124} as higher isotopes are well known.¹

Four new isotopes of iodine have been identified with mass assignments and decay characteristics as indicated in Table 1. The method of making mass assignments for some of these was to compare shapes of excitation curves of (α, xn) reactions on antimony in which known iodine activities were produced from the Sb^{123} and the new ones to be assigned from the Sb^{121} . The excitation functions as shown in Figure 1 are fairly crude as the only objective was to compare shapes in order to assign mass numbers. The experimental procedure was to irradiate antimony (57% Sb^{121} , 43% Sb^{123}) with helium ions of different energies, to isolate the iodine by a method in which the chemical yield could be measured, and then to resolve the iodine decay curves into the several components making use of distinctive radioactive properties.

13-hr. I^{123} .-- A 13-hr. component was assigned to I^{123} since its excitation function paralleled that for 56-day I^{125} , both curves being of the shape that would be expected for $(\alpha, 2n)$ reactions; in this case, on

¹For summary of data see: G. T. Seaborg and I. Perlman, Rev. Mod. Phys. 20, 585 (1948).

Figure 1 Excitation Functions for α, xn Reactions on Antimony

FIG. 1

Sb^{121} and Sb^{123} respectively. The yields as shown include corrections for natural abundances of Sb^{121} and Sb^{123} , and are based on the assumption that both the 13-hr. I^{123} and 56-day I^{125} have one K x-ray per disintegration, and assuming one percent counting efficiency in the argon-filled Geiger tubes. The 13-hr. I^{123} has conversion electrons of 150 ± 15 kev energy as determined with a low-resolution beta-ray spectrometer and by absorption in beryllium. Gamma-rays corresponding to this energy were also observed. This activity with similar half-life and radiation characteristics has been reported recently by Mitchell, Mei, Maienschein, and Peacock.²

4.5-day I^{124} and 13-day I^{126} .— The curves for both of these isotopes using normal antimony as a target are shown as appropriately labeled solid curves in Fig. 1. The curve for I^{126} is characteristic of an (α, n) reaction as is the low-energy part of that for I^{124} . The upswing beyond 30 Mev in the I^{124} curve is due to the $(\alpha, 3n)$ reaction on Sb^{123} . The points shown for the broken line projection of the I^{124} curve were obtained from the irradiation of separated * Sb^{121} from which I^{124} can be formed only by the (α, n) reaction.

The discrepancy in yields at 20 Mev between I^{126} and I^{124} is not explained; but it may be remarked that in other cases such as the fission of bismuth³ and the (γ, n) reaction on iodine⁴, the measured yield of I^{126} has been much lower than expected, making it appear that not all of the decay events take place through the 1-Mev beta-particle upon which the yields are based.

*The sample of separated antimony consisting of 99.3% Sb^{121} , 0.7% Sb^{122} was obtained from the Isotopes Division of the U.S. Atomic Energy Commission.

²A. C. G. Mitchell, J. Y. Mei, F. C. Maienschein, and C. L. Peacock, Phys. Rev. 76, 1450 (1949).

³R. H. Goeckermann and I. Perlman, Phys. Rev. 76, 628 (1949).

⁴M. L. Perlman, Phys. Rev. 75, 988 (1949).

The half-life of I^{124} was measured as 4.5 days and the β^+ -energy as 2.1 ± 0.1 Mev. It is estimated from the yield of K x-rays that this isotope decays only about 30% by positron-emission and 70% by electron-capture.

4-min. I^{122} .-- In experiments in which an hour was consumed in chemical separation, no activity appeared which followed the excitation function of that part of the 4.5-day I^{124} curve produced by the $(\alpha, 3n)$ reaction. It was therefore assumed that I^{122} , which would be formed by the $Sb^{121}(\alpha, 3n) I^{122}$ reaction, is short-lived. More rapid chemistry showed a 4-min. iodine activity, and a single yield determination at 45 Mev showed it to be in the expected range as indicated in Fig. 1. The positron has an energy of 2.9 ± 0.1 Mev, and it is estimated that there is some electron-capture branching.

The assignment of this activity to I^{122} is also based on the fact that it was the only new activity beyond I^{123} , I^{124} , I^{125} , and I^{126} to appear at 45 Mev. A 1.8-hr. activity assigned to I^{121} (from $\alpha, 4n$ reaction) did not appear at this energy.

1.8-hr. I^{121} .-- An activity with 1.8-hr. half-life with a 1.2-Mev positron and conversion electrons of 185 kev appeared in irradiation of antimony with 60-, 100-, and 360-Mev helium ions. Its decay is followed by the appearance of 17-day Te^{121} in approximately the proper yield for a parent-daughter relationship.

30-min. $I^{120} (?)$.-- In the higher energy irradiations (100 and 360 Mev) a 30-min. activity appeared, having a hard positron of 4.0 Mev. The decay of the iodine containing this activity was followed by the appearance of a tellurium activity which could not be resolved uniquely but which does have a half-life of several days. It is possible that this activity is 4.5-day Te^{119} . This would indicate that I^{119} was present. However, the observed positron

energy of the 30-min. period is in better agreement with the prediction for I^{120} which may mean that I^{119} was also present but unobserved.

We wish to thank Dr. J. G. Hamilton and the staff of the Crocker Laboratory cyclotron and Mr. James Vale and the group operating the 184-inch cyclotron for the irradiations used in these studies. We also wish to acknowledge the contribution to some of the early phases of this work by Dr. M. Lindner now at the State College of Washington, Pullman, Washington. This work was performed under the auspices of the Atomic Energy Commission.

TABLE 1

Mass No.	Half-Life	Radiation	Energy
124	4.5 days	K, β^+	$\beta^+ = 2.1 \pm 0.1 \text{ Mev}$
123	13 hr.	K, γ, e^-	$e^- = 150 \pm 15 \text{ kev}$
122	4 min.	$\beta^+ (K, \gamma?)$	$\beta^+ = 2.9 \pm 0.1 \text{ Mev}$
121	1.8 hr.	K, β^+, γ, e^-	$e^- = 185 \pm 10 \text{ kev}$ $\beta^+ = 1.2 \pm 0.1 \text{ Mev}$
120?	$\sim 30 \text{ min.}$	$\beta^+, \gamma (K?)$	$\beta^+ = 4.0 \pm 0.2 \text{ Mev}$