UC Irvine

UC Irvine Previously Published Works

Title

Fusion reaction spectra produced by anisotropic³He ions during ICRF

Permalink

<https://escholarship.org/uc/item/97t4704j>

Journal

Nuclear Fusion, 24(5)

ISSN

0029-5515

Author

Heidbrink, WW

Publication Date

1984

DOI

10.1088/0029-5515/24/5/011

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

FUSION REACTION SPECTRA PRODUCED BY ANISOTROPIC ^3He IONS DURING ICRF

W.W. HEIDBRINK (Plasma Physics Laboratory,
Princeton University, Princeton, New Jersey,
United States of America)

ABSTRACT. For 'beam-target' fusion reactions, collimated measurements of the energy spectrum of one of the reaction products can provide information on the degree of anisotropy of the reacting beam ions. Early measurements of the spectrum of 15 MeV protons produced by reactions between energetic ^3He ions and relatively cold deuterons during fast-wave minority heating in the PLT tokamak found an asymmetric, downward-shifted fusion spectrum. New spectral measurements with a thicker detector that collects the full proton energy indicate that the velocity distribution of fast ^3He ions is peaked perpendicular to the tokamak magnetic field.

During fast-wave (ICRF) heating in the PLT tokamak [1, 2], ^3He minority ions with energies of 200 keV produce 15 MeV protons through $\text{d}(^3\text{He}, \text{p})\alpha$ fusion reactions with the majority deuterium species [2, 3]. 15 MeV protons are unconfined in PLT and can be detected by using silicon surface barrier detectors mounted near the vacuum vessel wall [3, 4]. In previous measurements of the energy spectrum of the 15 MeV protons produced during ICRF [2, 3], the highest proton energies were cut off by the finite depth of the surface barrier detector. When the full energy of all the protons is measured, the asymmetric, downward-shifted spectrum reported previously becomes a spectrum with two peaks separated by about 2 MeV. Analysis of this spectrum in terms of a model two-component distribution function that has one component purely perpendicular and the other purely isotropic in velocity space indicates that $\geq 90\%$ of the high-energy ^3He ions created by the waves have velocities perpendicular to the tokamak toroidal field.

In the experiment, 15 MeV protons produced by the $\text{d}(^3\text{He}, \text{p})\alpha$ fusion reaction were detected with a proton spectrometer used previously on PLT [3]. The spectrometer was modified to use a surface barrier detector with a depleted region of 1800 μm , so that protons with energies up to 20 MeV deposited their full energy in the depleted region of the detector. The previous apparatus only collected the full energy of protons with less than 15.5 MeV. The spectrometer was collimated to measure perpendicular protons

(collimator FWHM = 6.5°) produced in the plasma between major radii of approximately 102 cm and 140 cm, with nearly constant detection efficiency over this range [3]. The ^3He cyclotron layer was at major radii of 125–143 cm for these discharges.

The measured 15 MeV proton spectra exhibit a broad, two-lobed structure (Fig. 1a) for discharges characterized by plasma currents of 500–550 kA, toroidal fields of 28–32 kG, line-average electron densities of $(1.7\text{--}3.0) \times 10^{13} \text{ cm}^{-3}$, ICRF powers of

FIG.1. (a) Proton spectrum during ^3He minority ICRF heating (240 kW) in a discharge with $I_\phi = 500 \text{ kA}$, $\bar{n}_e = 2.7 \times 10^{13} \text{ cm}^{-3}$, and $B_\phi = 30.5 \text{ kG}$. The curve is the spectrum produced by an anisotropic perpendicular ^3He beam with maximum energy of 400 keV and temperature of 30 keV [5]. The absolute accuracy of the energy measurement is $\pm 2.5\%$; the resolution of the spectrometer is 0.5 MeV (full line-width). (b) Proton spectrum during co-injection of a 40 keV deuterium neutral beam into a deuterium and ^3He plasma in the absence of ICRF. The curve is the spectrum produced by an isotropic 40 keV deuterium beam with a classical slowing-down distribution ($f(v) \propto v^{-3}$) [5].

FIG. 2. 15 MeV proton spectral width and $d(^3\text{He}, p)\alpha$ reaction rate versus line-average electron density (a), toroidal field (b), ICRF power (c), and estimated line-average ^3He density (d) with the other parameters held constant. $I_{\phi} \approx 500\text{ kA}$. The dashed curve through the reaction rate data is the fit used to calculate the spectral width predicted by Eq. 2 (solid curves).

100–500 kW at 30 MHz, and $d(^3\text{He}, p)\alpha$ reaction rates of $\geq 10^{11} \text{ s}^{-1}$. In contrast, co-injection of a 40 keV deuterium neutral beam into a deuterium and ^3He plasma in the absence of ICRF produced a relatively narrow 15 MeV proton spectrum with a single peak (Fig. 1b). The width of the proton spectrum during ICRF was found to depend less sensitively on electron density, ICRF power, toroidal magnetic field, and ^3He density than the $d(^3\text{He}, p)\alpha$ reaction rate (Fig. 2).

Operation of the spectrometer was restricted to a relatively narrow parameter regime by the requirement that the reaction rate be sufficiently large for statistically significant measurements but sufficiently small to avoid appreciable pulse pile-up. The ratio of counts in the low-energy peak to the number of counts in the central minimum was 1.68 ± 0.29 for discharges with full spectral width ≥ 2.7 MeV; the ratio of counts in the high-energy peak to counts in the central minimum was 1.38 ± 0.22 .

A twin-lobed spectrum cannot be produced by an isotropic beam of ^3He ions. Although these spectral measurements of protons at a single angle of escape do not provide sufficient information to determine uniquely the direction of anisotropy of the ^3He tail, they are consistent with theoretical predictions of a perpendicular anisotropic tail [3, 6]. Twin lobes are produced by a perpendicular ^3He distribution because, although the probability of undergoing a fusion reaction is constant throughout a ^3He ion's gyro-period, more of the protons detected in the poloidal plane make an angle parallel or antiparallel to the ^3He velocity vector than perpendicular to the ^3He velocity, resulting in a proton distribution that has more protons with a maximal Doppler shift than without any shift [5]. The spacing of the lobes implies that perpendicular ^3He ions with energy of about 40 keV in the case of the most closely spaced peaks and of about 300 keV in the case of the most widely spaced peaks made the dominant contribution to the observed proton spectra. Theoretically, the ^3He distribution is predicted to be perpendicular for energies $\geq 20 Z_{\text{eff}} T_e \approx 30 \text{ keV}$ [6]. The instrumental resolution was too poor in these experiments to determine the minimum energy at which anisotropy exists. The ratio of counts between lobes to the number of counts in the peaks implies that the isotropic contribution to the observed spectrum is less than 10% of the perpendicular contribution.

The observation that the proton spectral width is less sensitive to the parameters affecting ^3He tail heating than the $d(^3\text{He}, p)\alpha$ reaction rate (Fig. 2) is consistent with theoretical expectations. The proton energy distribution $F(E_p)$ can be approximated as

$$F(E_p) \propto \int dE_{\text{He}} \sigma(E_{\text{He}}) E_{\text{He}} f(E_{\text{He}}) \hat{F}(E_p, E_{\text{He}}) \quad (1)$$

where E_p is the proton energy, σ the fusion cross-section, $f(E_{\text{He}})$ is the energy distribution of the ^3He ions, and $\hat{F}(E_p, E_{\text{He}})$ the proton energy distribution produced by helium ions of energy E_{He} . Analytical expressions for $\hat{F}$ for isotropic and anisotropic ^3He distributions are given in Ref. [5]. For a ^3He distribution that decreases rapidly with energy, such as $f(E) \propto \exp(-E/T)$, where T is the tail temperature, the weighting factor $\sigma(E_{\text{He}}) E_{\text{He}} f(E_{\text{He}})$ peaks strongly for some energy $E_{\text{He}} = E_{\text{peak}}$ and the proton energy distribution function is approximately $F(E_p) \approx \hat{F}(E_p, E_{\text{peak}})$. Asymptotic analysis [7] of Eq. (1) using an analytical fit to the cross-section σ [8] indicates that the energy E_{peak} for which the weighting factor $\sigma E f$ is maximized depends on tail temperature according to $E_{\text{peak}} \approx 14 (\text{keV})^{1/3} T^{2/3}$. Since the spectral width scales like $\sqrt{E_{\text{He}}}$ [5], this implies that the spectral width scales approximately as $T^{1/3}$. In contrast to this weak algebraic dependence on tail temperature, the analysis predicts a strong exponential dependence of the $d(^3\text{He}, p)\alpha$ reaction rate $R_{d\text{He}}$ on tail temperature, $R_{d\text{He}} \propto \exp[-43(\text{keV}/T)^{1/3}]$. The reaction rate also depends on the number density of ^3He tail ions, while the proton spectrum produced by the minority tail depends on tail temperature alone. Combining expressions, the spectral width W is expected to scale with reaction rate R and ^3He density n as

$$W_2 = W_1 \left[1 + 0.023 T_1^{1/3} \ln \left(\frac{n_2 R_1}{n_1 R_2} \right) \right]^{-1} \quad (2)$$

The predictions of Eq. (2) are plotted in Fig. 2 by using the measured value of the reaction rate and a single normalization ($W_1 = 3.0 \text{ MeV}$, $R_1 = 4 \times 10^{11} \text{ s}^{-1}$, $n_1 \approx 1.0 \times 10^{12} \text{ cm}^{-3}$, $T_1 \approx 40 \text{ keV}$ is implied by $W_1 = 3.0 \text{ MeV}$). A major uncertainty in the application of Eq. (2) is determination of the ^3He density, which was estimated from the rise in electron density when neutral ^3He was puffed into the vacuum vessel about 50 ms before the ICRF pulse, but which differs from this value due to ^3He pumping and desorption [9]. The measured scaling of spectral width with reaction rate is consistent with the theoretical prediction of Eq. (2) within the experimental error (Fig. 2). When the ^3He gas puff was reduced so that the density rise was below the minimum detectable level ($\Delta n_e \lesssim 0.2 \times 10^{12} \text{ cm}^{-3}$) (Fig. 2d), the spectral width broadened slightly, implying that the tail temperature continued to rise with decreasing ^3He concentration;

FIG.3. A good fit to the asymmetric proton spectrum measured previously [3] is found by assuming that the true proton spectrum was produced by an anisotropic perpendicular ^3He beam with maximum energy 400 keV and temperature 50 keV but that this spectrum was distorted by a thin detector with the model response

$$E_{\text{det}} = \begin{cases} E_p, & \text{if } E_p \leq 15.4 \text{ MeV} \\ 3(15.4 \text{ MeV} - 2E_p), & \text{if } E_p > 15.4 \text{ MeV} \end{cases}$$

The model detector response is based on calculations summarized in Fig.2.5 of Ref. [10]. The counts below 12.5 MeV in the data are thought to be protons that lose energy in the walls of the collimator.

but the reaction rate fell, implying that the reduction in number density of ^3He ions had a stronger effect on the reaction rate than the increase in tail temperature.

The probable explanation for the asymmetric, downward-shifted proton spectra observed previously in PLT [3] is that the thinner detector used in those experiments failed to collect the full energy of the protons in the upper lobe of the distribution (Fig.3).

In conclusion, measurements of the spectrum of 15 MeV protons produced during fast-wave minority heating in the PLT tokamak indicate that $\approx 90\%$ of the ≈ 200 keV ^3He ions near the major radius of the device are anisotropic in velocity space. The width of the proton spectrum is increased by increasing the RF power, by reducing the electron or ^3He density, and by adjusting the toroidal field so that the ^3He resonance layer is in the centre of the discharge, but measurements of the $\text{d}(^3\text{He}, \text{p})\alpha$ reaction rate are much more

sensitive to changes in tail temperature than are the spectral measurements. The tail temperature implied by the spectral measurements is typically between 30 and 50 keV.

ACKNOWLEDGEMENTS

The author thanks J.D. Strachan for suggesting this problem and for critically reading the manuscript and G. Hammett, R.E. Chrien, and D. Hwang for helpful discussions. G. Estep provided technical assistance. The support of the PLT and ICRF groups under J. Hosea and the Neutral Beam group under G. Schilling are gratefully acknowledged, with special credit due J.R. Wilson and P. Colestock for forming the plasmas diagnosed in this experiment. This work was supported by US Department of Energy Contract No. DE-AC02-76-CHO-3073.

REFERENCES

- [1] HOSEA, J., BOYD, D., BRETZ, N., CHRIEN, R., COHEN, S., et al., in *Plasma Physics and Controlled Nuclear Fusion Research* (Proc. 8th Int. Conf. Brussels, 1980), Vol. 2, IAEA, Vienna (1981) 95.
- [2] HWANG, D., BITTER, M., BUDNY, R., CAVALLO, A., CHRIEN, R., et al., in *Plasma Physics and Controlled Nuclear Fusion Research* (Proc. 9th Int. Conf., Baltimore, 1982), Vol. 2, IAEA, Vienna (1983) 3.
- [3] CHRIEN, R.E., STRACHAN, J.D., *Phys. Fluids* **26** (1983) 1953.
- [4] CHRIEN, R.E., COLESTOCK, P.L., EUBANK, H.P., HOSEA, J.C., HWANG, D.Q., et al., *Phys. Rev. Lett.* **46** (1981) 535.
- [5] HEIDBRINK, W.W., *Fusion Reaction Spectra Produced by Anisotropic Fast Ions in the PLT Tokamak*, Princeton Plasma Physics Lab. Rep. PPP-L-2079 (Feb. 1984).
- [6] STIX, T.H., *Nucl. Fusion* **15** (1975) 737.
- [7] A similar analysis for the $\text{d}(\text{d}, \text{n})$ fusion reaction is found in POST, R.F., *Rev. Mod. Phys.* **28** (1956) 338.
- [8] MILEY, G.H., TOWNER, H., IVICH, N., *Fusion Cross Section and Reactivities*, Rep. COO-2218-17, University of Illinois, Urbana, Ill. (1974).
- [9] CHRIEN, R.E., EUBANK, H.P., MEADE, D.M., STRACHAN, J.D., *Nucl. Fusion* **21** (1981) 1661.
- [10] CHRIEN, R.E., *Measurements of Fusion Reactions from a Tokamak Plasma*, PhD Thesis, Princeton University (1982) 46.

(Manuscript received 28 December 1983

Final manuscript received 23 March 1984)