

# UCLA

## UCLA Previously Published Works

### Title

Smoking habits and parathyroid hormone concentrations in young adults: The CARDIA study

### Permalink

<https://escholarship.org/uc/item/984127wv>

### Authors

Fujiyoshi, Akira  
Polgreen, Lynda E  
Gross, Myron D  
[et al.](#)

### Publication Date

2016-12-01

### DOI

10.1016/j.bonr.2016.04.003

Peer reviewed



## Smoking habits and parathyroid hormone concentrations in young adults: The CARDIA study



Akira Fujiyoshi<sup>a,b,\*</sup>, Lynda E. Polgreen<sup>c</sup>, Myron D. Gross<sup>d</sup>, Jared P. Reis<sup>e</sup>, Stephen Sidney<sup>f</sup>, David R. Jacobs Jr.<sup>a</sup>

<sup>a</sup> Division of Epidemiology & Community Health, School of Public Health, University of Minnesota, MN, USA

<sup>b</sup> Department of Public Health, Shiga University of Medical Science, Shiga, Japan

<sup>c</sup> Harbor-UCLA Medical Center, Torrance, CA, USA

<sup>d</sup> Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN, USA

<sup>e</sup> Division of Cardiovascular Sciences, National Heart, Lung, and Blood Institute, Bethesda, MD, USA

<sup>f</sup> Kaiser Permanente Northern California, Oakland, CA, USA

### ARTICLE INFO

#### Article history:

Received 29 December 2015

Received in revised form 29 March 2016

Accepted 27 April 2016

Available online 28 April 2016

#### Keywords:

Parathyroid hormone

Smoking

Young adults

### ABSTRACT

Conflicting results have been reported concerning a relationship between smoking and serum PTH. Our study objective was to examine whether smoking was associated with serum PTH independent of correlates of PTH among young adults, and explore potential mechanisms.

This was a cross-sectional study of healthy individuals, 24–36 years old, examined during 1992 through 1993 in California, USA (a subset of Coronary Artery Risk Development in Young Adults study).

Linear regression was used to obtain adjusted means of PTH according to smoking habit (current, former, never). Biomarkers for calcium metabolism and bone turnover (including serum concentrations of osteocalcin, bone-specific alkaline phosphatase, and 24-hour urinary excretion of calcium) and bone mineral density were similarly compared by smoking.

376 participants were analyzed (171 women, 181 black). Over half reported never smoking. We observed lower PTH in current smokers compared to non-smokers and found no evidence of an interaction by race and sex. PTH was lowest in current smokers, intermediate in former smokers, and highest in never smokers (geometric mean PTH: 23.6, 26.7, 27.4 pg/mL, respectively; *P* for trend, 0.006) after adjusting for potential confounders including calcium intake. Among the biomarkers, serum osteocalcin concentration and 24-hour urinary excretion of calcium were lowest in current smokers. We observed no smoking-related difference in bone mineral density.

In this community-based sample of young adult men and women, smoking was associated with significantly lower PTH concentration. The mechanism and clinical implication of the finding, however, remains uncertain.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Smoking is generally associated with lower concentration of vitamin D, lower bone mineral density and increased risk of fracture in the elderly. Therefore, a compensatory increase in parathyroid hormone

(PTH) may be expected (Jorde et al., 2005). However, conflicting results have been reported in the literature on the relation between smoking and PTH. While some studies reported higher PTH among smokers (i.e. expected direction) (Rapuri et al., 2000; Szulc et al., 2002; Ortego-Centeno et al., 1997), others have reported lower PTH concentration among smokers (Mellstrom et al., 1993; Landin-Wilhelmsen et al., 1995; Brot et al., 1999; Need et al., 2002). One of the largest population-based observational studies documented this unexpected finding cross-sectionally (Jorde et al., 2005), yet the underlying mechanism and clinical implication of the finding remains to be elucidated.

In this study, we investigated a cross-sectional relationship between smoking habits and concentration of PTH using a population-based sample of men and women in the USA. The primary aim was to investigate whether PTH concentration differs among current, former and

*Abbreviations:* PTH, Parathyroid hormone; CARDIA, Coronary Artery Risk Development in Young Adults; BMI, Body mass index; 25OHD, 25-hydroxyvitamin D; 1,25(OH)<sub>2</sub>D, 1,25-dihydroxyvitamin D; BAP, Bone-specific alkaline phosphatase; U-PYDcr, 24-hour urinary excretion of pyridinoline standardized for urinary excretion of creatinine; BMD, Bone mineral density.

\* Corresponding author at: Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, West Bank Office Building, 1300 S. 2nd Street, Minneapolis, MN 55454-1015, USA.

E-mail addresses: [fujiy001@umn.edu](mailto:fujiy001@umn.edu), [afujij@belle.shiga-med.ac.jp](mailto:afujij@belle.shiga-med.ac.jp) (A. Fujiyoshi).

never smokers after accounting for potential confounders. In light of inconsistencies in the literature, we did not have a hypothesis about the direction of the PTH and smoking association. The secondary aim was to explore potential mechanisms using biological markers related to calcium metabolism and bone turnover according to smoking habit.

## 2. Methods

### 2.1. Study population

We studied a subgroup of participants of the Coronary Artery Risk Development in Young Adults (CARDIA) study. A detailed description of CARDIA has been published elsewhere (Friedman et al., 1988). In brief, CARDIA is a cohort study of the evolution of cardiovascular risk among young adults, conducted since 1985. The participants for the present study were enrolled in the CARDIA study in the Oakland Clinical Center, California, who were returning for their fourth examination from June 1992 through May 1993 to participate in an ancillary study of bone mineral homeostasis (Bikle et al., 1999). All subjects, aged 24–36 years, were in good health and had a serum creatinine level <1.6 mg/dL and a serum calcium level ranging from 8.8 to 10.3 mg/dL. Within the year before the examination, all women menstruated regularly, were not taking oral contraceptive agents, and had not been pregnant  $\geq 10$  weeks or breastfeeding. Subjects with medical conditions or taking medications known to alter calcium homeostasis were excluded from the study (Bikle et al., 1999). The data were collected with written informed consent and with approval of the local Institutional Review Board. Of 402 ancillary study participants, we included those participants with no missing variables directly relevant to the primary aim (i.e. to examine association between smoking and PTH). We excluded those who gave unrealistic estimate of energy intake (800 or 8000 kcal/day in men; 600 or 6000 kcal/day in women), leaving 376 participants for analysis.

### 2.2. Measurement

Demographic information was obtained using standardized questionnaires. Weight was measured on a balance beam scale. Height was measured using a vertically mounted metal centimeter ruler. Body mass index (BMI) was computed as weight (kg) divided by square of height (m). At examination, fasting blood including creatinine, phosphate and calcium (collected between 8:00 and 11:00 AM) and 24-hour urine samples were obtained for analysis using routine laboratory procedures and an automated chemistry analyzer (Bikle et al., 1999).

PTH and other biomarkers were measured on stored samples 3 years after study completion (Bikle et al., 1999; Ettinger et al., 1997). PTH was measured using a 2-site immunoradiometric assay that detects only intact PTH (i.e. biologically active 84-amino-acid peptide) (Nussbaum et al., 1987). Serum 25-hydroxyvitamin D (25OHD) was measured by RIA in the Calcitropic Hormone Reference Laboratory, University of California, San Francisco. Serum 1,25-dihydroxyvitamin D ( $1,25(\text{OH})_2\text{D}$ ) was measured by a competitive protein-binding assay using the vitamin D receptor present in calf thymus cytosol (Bikle et al., 1999). Osteocalcin was measured using a mid-molecule-specific RIA for human osteocalcin (Taylor et al., 1988). Bone-specific alkaline phosphatase (BAP) was measured spectrophotometrically after heat inactivation (Farley et al., 1981). Free pyridinoline cross-links in urine were measured using an ELISA developed by Metra Biosystems (Seyedin et al., 1993). The results of this method were compared with that of direct assessment after HPLC purification (Black et al., 1988) for 20 subjects in each race and gender group to verify the comparability of this method with that using HPLC ( $r > 0.9$  for all groups) before adopting this method for use in assessing all subjects (Bikle et al., 1999). 24-hour urinary excretion of pyridinoline was standardized for urinary excretion of creatinine (U-PYDcr) for analysis.

### 2.3. Assessment of lifestyle information

The usual diet was assessed by a diet history interview (CARDIA dietary history) in which food models and measuring cups and spoons were used to estimate portion size. The previous month was used as a frame of reference for estimating the usual intake. Intake of dietary calories and calcium were estimated according to the nutrient data base developed by the Nutrition Coordinating Center at the University of Minnesota. The estimate of calcium intake included supplemental sources. Validity and reliability of the CARDIA dietary history has been published elsewhere (Liu et al., 1994). Sun exposure time was assessed by query for weekday and weekend separately and then calculated as hours per week. The CARDIA physical activity history questionnaire was used to quantify physical activity level, and scored accordingly (physical activity score). The amount of time per week spent in leisure, occupational, and household physical activity over the previous 12 months was assessed. We estimated total physical activity expressed in exercise units as a product of intensity and frequency. The validity and reliability of the questionnaire was published elsewhere (Jacobs et al., 1989).

Additionally, we examined bone mineral density (BMD) at various sites according to smoking habit. Dual-energy x-ray absorptiometry measurement of BMD for the whole body, total hip, left and right total arms, and lumbar spine was obtained by using a Hologic QDR 2000 densitometer (Hologic, Inc.) in the array scanning mode, separately for each area of the body (total hip, lumbar spine, and whole body). The total arm was assessed using the whole-body scan (Fujiyoshi et al., 2013). In vivo precision for BMD, based on repeated scans of 20 volunteers done 1 to 6 weeks apart and expressed as a coefficient of variation, was 0.9% for whole-body BMD.

### 2.4. Statistical analysis

We used linear regression models to obtain adjusted PTH concentrations according to smoking habits (Current, Former, and Never). Due to skewed distributions, we log-transformed the following variables in our statistical models: PTH, 25OHD, BAP, U-PYDcr and calcium intake in density. When any of these variables was used as a dependent variable, we back-transformed the results to present the adjusted geometric means for ease of interpretation. For the main analyses, we constructed the following models: Model 1 adjusted for age (years), race (Black, White), sex, and education (up to 15 years, 16 years or greater). Model 2 adjusted additionally for calcium intake in density (mg/1000 kcal/day, supplements included, log-transformed), serum concentrations of 25OHD (ng/mL, log-transformed), creatinine (mg/dL), and season (Jan–Mar, Apr–Jun, July–Sep, and Oct–Dec) of blood draw. Model 3 further adjusted for physical activity score and sun-exposure time (hours/week). Model 4 further adjusted for body mass index ( $\text{kg}/\text{m}^2$ ).

For each sex and race/ethnicity group, we observed that PTH concentrations tended to be high in never smokers, and low in current smokers, then we tested interaction by sex or race/ethnicity on the association of smoking and PTH by inserting a product term in the models. All the interaction terms were non-significant throughout the models. Therefore, we presented sex- and race-combined results. We tested linear trend of PTH across the smoking habits by using an ordinal variable (0 = never, 1 = former, and 2 = current smoker). As sensitivity analyses, we repeated the main analyses replacing calcium density in the models with either a) calcium density excluding supplemental source (i.e. dietary source of calcium only) or, b) absolute amount of calcium intake (mg/day) or, c) serum calcium concentration (mg/dL).

For the secondary aim, we estimated the adjusted means of the following biological markers: serum phosphate [mmol/L], serum  $1,25(\text{OH})_2\text{D}$  [nmol/L], 24-hour urinary calcium [mmol/day], fractional excretion of calcium (FEca, also known as calcium/creatinine clearance

ratio), which was calculated as the clearance of calcium divided by the clearance of creatinine or:

$$(U_{Ca}/P_{Ca}) \times (P_{Cr}/U_{Cr}).$$

where  $U_{Ca}$  and  $U_{Cr}$  are the 24-hour urinary excretions of calcium or creatinine, and  $P_{Ca}$  and  $P_{Cr}$  are the plasma concentrations of calcium and creatinine (Bikle et al., 1999). For bone-turnover markers, we used serum osteocalcin [ng/mL], serum BAP [IU/L], and U-PYDcr [micromole/mmol creatinine]. Osteocalcin and BAP are markers for bone formation, whereas U-PYDcr is a marker for bone resorption. We used the same sets of adjusting covariates as in the main analysis except for FEca in that serum creatinine was excluded from the models. This was because FEca itself is standardized by serum creatinine. Lastly, we compared adjusted means of the following BMDs according to smoking habit: whole body, arms (average of left and right), lumbar spine, and total hip.

All the statistical tests were considered to be significant when  $P$  values were  $<0.05$ . Statistical analysis was conducted by SAS version 9.4 (SAS Institute).

### 3. Results

We analyzed 376 participants (171 women, 181 black, age 24–36 years). Demographics of the participants according to smoking habit for each sex and race/ethnicity were given in Table 1.

In all sex-, race/ethnicity- groups, more than half of the participants (56–72%) reported never smoking. Over a quarter of black men (25.8%) and black women (26.1%) reported current smoking, whereas 10.7% of White men and 15.7% of White women did so. Across the sex and race/ethnicity strata, current smokers tended to have greater sun-exposure hours and higher BMI (with exception for calcium intake in white women and BMI for black men), and tended to have fewer

**Table 1**  
Demographics of participants according to smoking habit.

No.	Black men (N = 93)			Black women (N = 88)			White men (N = 112)			White women (N = 83)		
	Current	Former	Never	Current	Former	Never	Current	Former	Never	Current	Former	Never
	24	9	60	23	16	49	12	19	81	13	20	50
Age, years	30.8	32.0	30.4	30.4	31.1	31.1	31.1	30.8	31.4	29.7	33.1	31.8
Education, years	14	13	15	13	14	14	13	15	16	13	15	17
Body mass index, kg/m <sup>2</sup>	25.0	24.9	27.0	29.2	26.9	28.6	26.6	26.1	25.3	26.6	26.3	23.2
Energy intake, kcal/day	4140	4371	3304	2479	2066	2107	3527	3241	3199	2604	2209	2446
Calcium intake <sup>a</sup> , mg/day	1614	1263	1303	879	751	828	1506	1471	1449	1304	1391	1320
Calcium intake <sup>a</sup> in density, mg/1000 kcal/day	410	290	410	370	380	400	430	440	440	510	600	560
Physical activity score	498	428	533	239	253	298	347	504	423	359	312	387
Sun exposure time, hours/week	32.5	29.7	27.3	27.1	15.8	23.8	18.1	17.8	17.0	16.3	14.3	14.4
Calcium (serum), mg/dL	9.5	9.5	9.6	9.3	9.4	9.3	9.7	9.5	9.6	9.4	9.3	9.4
25OHD (serum), ng/mL	32.4	35.3	32.7	20.4	19.2	23.6	37.5	38.2	38.6	39.6	36.1	36.6
Creatinine (serum), mg/dL	0.99	1.04	1.03	0.79	0.75	0.80	0.89	0.92	0.95	0.78	0.74	0.78
Biomarkers related to calcium-metabolism												
Phosphate (serum), mg/dL	3.13	2.96	3.18	3.38	3.27	3.34	2.78	3.07	2.89	2.97	3.03	3.36
1,25(OH) <sub>2</sub> D (serum), pg/mL	46.1	45.9	43.8	45.3	49.0	43.1	40.3	38.0	39.8	38.9	39.7	37.9
24-hour urinary Ca, mmol/day	2.64	4.56	4.15	2.19	2.49	2.66	4.66	5.64	5.02	3.78	3.85	3.94
Fraction excretion of calcium (× 100)	0.74	0.86	0.85	0.57	0.58	0.71	1.07	1.18	1.06	0.93	0.91	1.09
Bone turnover marker												
Osteocalcin (serum), ng/mL	11.7	12.5	14.6	9.8	10.8	9.6	13.3	14.8	14.9	10.8	12.9	13.4
BAP (serum), U/L	9.2	7.6	9.8	7.2	8.2	8.4	9.0	9.9	8.5	7.1	7.4	6.7
U-PYDcr, μmol/mmol creatinine	24.9	23.5	24.5	34.3	41.2	37.6	28.8	27.4	26.9	33.6	37.3	39.2
Bone mineral density, g/cm <sup>2</sup>												
Whole body	1.27	1.32	1.30	1.17	1.15	1.16	1.16	1.19	1.18	1.11	1.11	1.08
Lumbar spine	1.14	1.20	1.13	1.16	1.08	1.13	1.02	1.08	1.02	1.03	1.05	1.04
Total hip	1.16	1.22	1.19	1.05	0.98	1.04	1.05	1.06	1.03	0.97	0.96	0.95
Arms (average of right and left)	0.98	1.02	1.02	0.81	0.79	0.81	0.94	0.95	0.95	0.80	0.79	0.77
Parathyroid hormone, pg/mL	24.4	22.2	25.7	30.5	37.8	38.4	26.0	25.4	26.9	24.3	31.2	27.0

Abbreviations: 25OHD, 25-hydroxyvitamin D; 1,25(OH)<sub>2</sub>D, 1,25-dihydroxyvitamin D; 24-hour urinary Ca, 24-hour urinary excretion of calcium; BAP: bone-specific alkaline phosphatase; U-PYDcr: urinary excretion of pyridinoline corrected with urinary excretion of creatinine.

<sup>a</sup> Calcium intake included supplemental source.

**Table 2**  
Crude and adjusted means of PTH (pg/mL) according to smoking habit.

	Current	Former	Never	Trend $P$	$P$ for pairwise comparison <sup>a</sup>	
	72	64	240		Current vs former	Current vs never
Crude, arithmetic means	26.6	29.8	29.0	–	–	–
Crude, geometric means	24.3	27.4	26.9	0.095	0.082	0.055
Model 1	23.5	26.8	27.4	0.008	0.051	0.005
Model 2	23.6	26.5	27.4	0.006	0.068	0.004
Model 3	23.5	26.6	27.4	0.005	0.051	0.003
Model 4	23.6	26.7	27.4	0.006	0.054	0.004

Trend  $P$ s were calculated using smoking status as an ordinal variable (0 = never smoker, 1 = former smoker, 2 = current smoker).

Values were all geometric means unless otherwise specified. Model 1 adjusted for age (years), race (Black, White), sex, education (up to 15 years, 16 years or greater). Model 2 adjusted for calcium intake in energy density (mg/1000 kcal/day, log-transformed), serum level of 25OHD (ng/mL, log-transformed), serum level of creatinine (mg/dL), and season (Jan–Mar, Apr–Jun, July–Sep, and Oct–Dec) in addition to Model 1. Model 3 adjusted for physical activity score and sun-exposure time (hours/week) in addition to Model 2. Model 4 adjusted for body mass index (kg/m<sup>2</sup>) in addition to Model 3.

<sup>a</sup> There was no statistically significant difference between former smokers and never smokers in all models.

education years. We observed no clear patterns by smoking habit for serum concentrations of calcium, 25OHD and creatinine across the strata.

Crude (arithmetic or geometric) and adjusted means of PTH according to smoking habit were shown in Table 2. Current smokers tended to have lower PTH concentrations compared to never smokers. With adjustment for basic demographics in Model 1 (age, race, sex, and education), a significant trend was observed such that PTH concentrations were lowest in current smokers and highest in never smokers. This

significant trend persisted across the models. Excluding supplemental source of calcium intake did not change the overall results (data not shown). Replacing energy-adjusted calcium intake with either absolute amount of calcium intake (mg/day) or serum calcium concentration (mg/dL) in the models yielded similar results (data not shown). In subgroup analysis restricted to current smokers ( $n = 72$ ), PTH concentration was not associated with number of cigarettes smoked per day (data not shown). Since the participants were not directly queried on duration and amount of cigarette smoking in the past, we did not test a dose-response relationship using pack-years of smoking. Since smoking is often associated with lower BMI, and serum PTH is associated with higher BMI (Kontogeorgos et al., 2015), there was a possibility of residual confounding by BMI on the observed association between smoking and PTH. To address this possibility, we have repeated the analyses stratified by median BMI (25.10 kg/m<sup>2</sup>) using Model 3. Results were consistent with the main one with no statistical evidence of interaction by BMI (Table 3).

Table 4 summarized results of markers related to calcium metabolism and bone turnover according to smoking habit. We observed significantly lower 24-hour urinary calcium excretion in current smokers compared to non-current smokers, with a similar pattern for FEca. Among bone turnover makers, osteocalcin was lower in current than non-current smokers with a significant linear trend. Serum concentrations of phosphate, 1,25(OH)<sub>2</sub>D, BAP, and U-PYDcr had no clear difference among the smoking groups.

We observed no significant difference in any BMD site across smoking categories after adjustment for sex and race with or without other covariates added (data not shown).

**4. Discussion**

In this generally healthy sample of black and white, young adult men and women, we observed that PTH concentration was lowest in current smokers, followed by former smokers, and highest in never smokers. This relationship appeared to be independent of factors that affect PTH concentration, such as dietary calcium intake and serum 25OH vitamin D concentration (Steingrimsdottir et al., 2005).

Smoking is generally associated with reduced bone density, and reduced concentrations of vitamin D among elderly populations. Therefore, a compensatory increase in PTH may be expected in smokers (Jorde et al., 2005). However, studies on PTH concentration comparing smoker/non-smoker are conflicting. Some studies showed lower

(Jorde et al., 2005; Mellstrom et al., 1993; Landin-Wilhelmsen et al., 1995; Brot et al., 1999; Need et al., 2002), but others reported higher PTH concentrations in smokers (Rapuri et al., 2000; Szulc et al., 2002; Ortego-Centeno et al., 1997). One of the largest population studies on this question examined 7896 men and women in Norway and found lower PTH concentrations in smokers compared to non-smokers regardless of age, sex, BMI, and serum creatinine concentration (the fifth Tromsø study) (Jorde et al., 2005). Our finding of lower PTH in smokers is consistent with the Tromsø study, and the observed relationship in our study was robust to adjustment for other factors associated with PTH including sun-exposure time, serum 25OHD concentrations.

Studies reporting higher PTH in current smokers, the opposite finding to ours, appear to have serious limitations in generalizing their results. In a study of healthy men in Spain, Ortego-Centeno and colleagues reported higher PTH in smokers ( $n = 15$ ) than non-smokers ( $n = 17$ ). However, the difference was not statistically significant ( $P$ -value not provided) and the sample size was small. (Ortego-Centeno et al., 1997) Szulc and colleagues reported statistically significantly higher PTH concentration in French current smokers ( $n = 35$ ) compared to never smokers ( $n = 93$ ) (Szulc et al., 2002). However, this comparison was based on subgroup of individuals who weighed <75 kg. Including all their participants (719 men, aged 51–85 years), the results appeared to be consistent with our findings, although not statistically significant (Szulc et al., 2002). In a study of 489 elderly women (65–77 years) in the USA, Rapuri and colleagues showed higher PTH in heavy smokers than non-smokers. However, the difference was not statistically significant, nor was it adjusted for serum 25OHD and other possible confounders (Rapuri et al., 2000).

Potential mechanisms have been proposed to explain lower PTH in smokers such as direct toxic effect of smoking on the parathyroid cells, enhanced degradation or suppressed secretion of PTH by smoking (Jorde et al., 2005; Brot et al., 1999; Need et al., 2002). Among those, Need and colleagues hypothesized that direct impairment of osteoblast function by smoking was likely to result in a slight increase in serum ionized calcium, leading to suppression of PTH (Need et al., 2002). Our finding of lower osteocalcin (bone formation marker) in current smokers than non-current smokers is consistent with this hypothesis and consistent with another study on healthy perimenopausal women (Brot et al., 1999) although the other bone formation marker, BAP, did not show a significant pattern in our study and we did not have variables to estimate serum ionized calcium. We also observed lower 24-hour urinary calcium excretion in current smokers than non-smokers, which seems consistent with the fifth Tromsø study reporting a lower but non-significant 24-hour urinary calcium excretion in smokers ( $n = 54$ ) compared to non-smokers ( $n = 151$ ) based on the small subgroup who underwent biochemical assessments (Jorde et al., 2005). The lower calcium excretion in current smokers could be due to lower osteoclast function in smokers and a response to low PTH levels. These actions allow for the maintenance calcium balance between formation and reabsorption from bone. Understanding the mechanism and clinical implication of the finding will require further study.

In comparing BMD at various sites, we observed no significant difference according to smoking habit in the study sample. Although smoking is a known risk factor for osteoporosis and reduced BMD in the elderly, it may require longer exposure time and/or a larger sample size for adverse effect of smoking on BMD to be evident given the relatively young age (24–36 years) of our study participants, the age range at which bone mass reaches its peak (Davies et al., 2005). Alternate methods of modeling exposure to smoking such as use of pack-year might have led to different findings (Callreus et al., 2013). However, we did not model exposure to smoking using pack-year given limitations in the method for querying smoking history in CARDIA.

A few studies have explicitly reported PTH concentration among former smokers relative to those in current and never smokers. The fifth Tromsø study reported similar concentration between former and never smokers, higher than current smokers. However, they took into

**Table 3**  
Adjusted means of PTH according to smoking habit stratified by median BMI.

		Current	Former	Never	Trend <i>P</i>	<i>P</i> for interaction by median BMI
BMI < median*	No.	31	36	121		
	adjusted PTH <sup>†</sup> , pg/mL	21.4	25.5	25.8	0.042	0.765
BMI ≥ median*	No.	41	28	119		
	adjusted PTH <sup>†</sup> , pg/mL	25.9	26.9	29.1	0.071	

\* Median BMI was 25.10 kg/m<sup>2</sup>. Participants were fairly evenly distributed by median BMI across the group of smoking habits, sex, and race/ethnicity. Percentage (%) of the participants with BMI < median for each group was as follows: never (50.4), former (56.3), and current (43.1) smoker, men (48.3), women (52.1), black (40.3), white (59.0).

† Values were geometric means adjusted for the same set of covariates as Model 3 in the text as follows: age (years), race (Black, White), sex, education (up to 15 years, 16 years or greater), calcium intake in energy density (mg/1000 kcal/day, log-transformed), serum level of 25OHD (ng/mL, log-transformed), serum level of creatinine (mg/dL), season (Jan–Mar, Apr–Jun, July–Sep, and Oct–Dec), physical activity score and sun-exposure time (hours/week).

To obtain *P*-values, smoking status was treated as an ordinal variable (0 = never smoker, 1 = former smoker, 2 = current smoker). *P* for interaction by median BMI was obtained by inserting a product term (smoking habit x median BMI) in the model.

**Table 4**  
Adjusted means of factors related to calcium metabolism and bone turnover.

	Smoking habit			Trend P	P for pairwise comparison	
	Current	Former	Never		Current vs former	Current vs never
Phosphate, serum, mg/dL	3.10	3.10	3.16	0.291	0.989	0.365
1,25(OH) <sub>2</sub> D, serum, pg/mL	42.4	43.2	41.3	0.332	0.677	0.476
24-h urinary Ca, mmol/day	3.22	4.18	4.03	0.042	0.024	0.018
FEca ( $\times 100$ ) <sup>a</sup>	0.80	0.90	0.93	0.082	0.287	0.074
Osteocalcin, serum, ng/mL	11.5	13.3	13.2	0.019	0.030	0.009
BAP <sup>b</sup> , serum, IU/L	7.1	8.0	7.8	0.126	0.065	0.068
U-PYDcr <sup>b</sup> , $\mu$ mol/mmol creatinine	28.6	29.7	29.5	0.550	0.462	0.472

With exception for FEca, the mean values were adjusted for the same covariates as Model 4 in the main analyses as follows: age (years), race (Black, White), sex, education (up to 15 years, 16 years or greater), energy density of calcium intake (mg/1000 kcal/day, log-transformed), levels of 25OHD (ng/mL, log-transformed) and creatinine (mg/dL), season (Jan–Mar, Apr–Jun, July–Sep, and Oct–Dec), physical activity score, sun-exposure time (hours/week) and body mass index (kg/m<sup>2</sup>). Trend Ps were calculated using smoking status as an ordinal variable (0 = never smoker, 1 = former smoker, 2 = current smoker).

Abbreviations: 1,25(OH)<sub>2</sub>D: 1,25-dihydroxyvitamin D; 24-hour urinary Ca: 24-hour urinary excretion of calcium; FEca: fractional excretion of calcium; U-PYDcr: urinary excretion of pyridinoline corrected with urinary excretion of creatinine; BAP: bone-specific alkaline phosphatase.

<sup>a</sup> For FEca, serum creatinine concentration was excluded from the model adjustment because this measure is standardized with creatinine concentration.

<sup>b</sup> Values were given in geometric means.

account only those who quit smoking for one year or longer, not accounting for recent quitters. In a study of 405 postmenopausal women, PTH concentrations were lowest in current, intermediate in former, and highest in never smokers (Need et al., 2002), which is consistent with our finding. The association between smoking and PTH may persist for a while, when quitting, then decline over time. To better clarify the relationship in future, a large-scale longitudinal study with repeated assessment for smoking and PTH is warranted. Such study should be able to examine not only whether the association persists longitudinally, but also dose-response of smoking and length of smoking cessation (former smokers) in relation to PTH.

Limitations need to be considered in interpreting our findings. First, the study was cross-sectional in design, making it difficult to prove temporal relationships between exposure (smoking) and outcome (PTH) although we believe it is uncommon for individuals to quit or not to initiate smoking due to high PTH concentration. Second, given the observational nature of the study, we cannot refute the possibility of residual or unmeasured confounding on the results. Third, our data were collected and measured in the 1990s. Some measurements used in the study may differ from the ones more recently used. Strengths of our study include availability of important factors associated with PTH enabling us to control for potential confounders including serum 25OHD and creatinine, calcium intake, body mass index and physical activity. Likewise, various markers of calcium metabolism and bone turnover allowed us to explore potential mechanism of the findings. Another strength is enrollment of black and white men and women aged 24 to 36 years, making the results likely generalizable to healthy young adults.

## 5. Conclusions

In summary, we found significantly lower concentration of PTH in current smokers, followed by former smokers, compared to never smokers in a population-based sample of young adult men and women. The relation was not explained by determinants of PTH such as dietary calcium intake and serum vitamin D. Additionally, we observed low serum osteocalcin and low 24-hour urinary calcium excretion in current smokers but no difference in BMD according to smoking habits. The clinical implication and potential mechanism of the findings, however, remain unclear, warranting further research.

## Grant sources

The Coronary Artery Risk Development in Young Adults Study (CARDIA) is supported by contracts HHSN268201300025C, HHSN268201300026C, HHSN268201300027C, HHSN268201300028C, HHSN268201300029C, and HHSN268200900041C from the National

Heart, Lung, and Blood Institute and the Intramural Research Program of the National Institute on Aging (NIA), and an intra-agency agreement between NIA and NHLBI (AG0005). Additional support was given to one of the authors (AF) by the Fulbright Program (2014 RS).

## Disclosure statement

The authors have nothing to disclose.

## References

- Bikle, D.D., Ettinger, B., Sidney, S., Tekawa, I.S., Tolan, K., 1999. Differences in calcium metabolism between black and white men and women. *Miner. Electrolyte Metab.* 25, 178–184.
- Black, D., Duncan, A., Robins, S.P., 1988. Quantitative analysis of the pyridinium crosslinks of collagen in urine using ion-paired reversed-phase high-performance liquid chromatography. *Anal. Biochem.* 169, 197–203.
- Brot, C., Jorgensen, N.R., Sorensen, O.H., 1999. The influence of smoking on vitamin D status and calcium metabolism. *Eur. J. Clin. Nutr.* 53, 920–926.
- Callreus, M., McGuigan, F., Akesson, K., 2013. Adverse effects of smoking on peak bone mass may be attenuated by higher body mass index in young female smokers. *Calcif. Tissue Int.* 93, 517–525.
- Davies, J.H., Evans, B.A., Gregory, J.W., 2005. Bone mass acquisition in healthy children. *Arch. Dis. Child.* 90, 373–378.
- Ettinger, B., Sidney, S., Cummings, S.R., Libanati, C., Bikle, D.D., Tekawa, I.S., Tolan, K., Steiger, P., 1997. Racial differences in bone density between young adult black and white subjects persist after adjustment for anthropometric, lifestyle, and biochemical differences. *J. Clin. Endocrinol. Metab.* 82, 429–434.
- Farley, J.R., Chesnut 3rd, C.H., Baylink, D.J., 1981. Improved method for quantitative determination in serum of alkaline phosphatase of skeletal origin. *Clin. Chem.* 27, 2002–2007.
- Friedman, G.D., Cutter, G.R., Donahue, R.P., Hughes, G.H., Hulley, S.B., Jacobs Jr., D.R., Liu, K., Savage, P.J., 1988. CARDIA: study design, recruitment, and some characteristics of the examined subjects. *J. Clin. Epidemiol.* 41, 1105–1116.
- Fujiyoshi, A., Polgreen, L.E., Hurley, D.L., Gross, M.D., Sidney, S., Jacobs Jr., D.R., 2013. A cross-sectional association between bone mineral density and parathyroid hormone and other biomarkers in community-dwelling young adults: the CARDIA study. *J. Clin. Endocrinol. Metab.* 98, 4038–4046.
- Jacobs Jr., D.R., Hahn, L.P., Haskell, W.L., Pirie, P., Sidney, S., 1989. Validity and reliability of short physical activity history: cardia and the Minnesota heart health program. *J. Cardpulm. Rehabil.* 9, 448–459.
- Jorde, R., Saleh, F., Figenschau, Y., Kamycheva, E., Haug, E., Sundsfjord, J., 2005. Serum parathyroid hormone (PTH) levels in smokers and non-smokers. The fifth Tromso study. *Eur. J. Endocrinol.* 152, 39–45.
- Kontogeorgos, G., Trimpou, P., Laine, C.M., Olerod, G., Lindahl, A., Landin-Wilhelmsen, K., 2015. Normocalcaemic, vitamin D-sufficient hyperparathyroidism - high prevalence and low morbidity in the general population: A long-term follow-up study, the WHO MONICA project, Gothenburg, Sweden. *Clin. Endocrinol.* 83, 277–284.
- Landin-Wilhelmsen, K., Wilhelmsen, L., Lappas, G., Rosen, T., Lindstedt, G., Lundberg, P.A., Wilske, J., Bengtsson, B.A., 1995. Serum intact parathyroid hormone in a random population sample of men and women: relationship to anthropometry, life-style factors, blood pressure, and vitamin D. *Calcif. Tissue Int.* 56, 104–108.
- Liu, K., Slattery, M., Jacobs Jr., D., Cutter, G., McDonald, A., Van Horn, L., Hilner, J.E., Caan, B., Bragg, C., Dyer, A., et al., 1994. A study of the reliability and comparative validity of the cardia dietary history. *Ethn. Dis.* 4, 15–27.
- Mellstrom, D., Johansson, C., Johnell, O., Lindstedt, G., Lundberg, P.A., Obrant, K., Schoon, I.M., Toss, G., Ytterberg, B.O., 1993. Osteoporosis, metabolic aberrations, and increased risk for vertebral fractures after partial gastrectomy. *Calcif. Tissue Int.* 53, 370–377.

- Need, A.G., Kemp, A., Giles, N., Morris, H.A., Horowitz, M., Nordin, B.E., 2002. Relationships between intestinal calcium absorption, serum vitamin D metabolites and smoking in postmenopausal women. *Osteoporos. Int.* 13, 83–88.
- Nussbaum, S.R., Zahradnik, R.J., Lavigne, J.R., Brennan, G.L., Nozawa-Ung, K., Kim, L.Y., Keutmann, H.T., Wang, C.A., Potts Jr., J.T., Segre, G.V., 1987. Highly sensitive two-site immunoradiometric assay of parathyrin, and its clinical utility in evaluating patients with hypercalcemia. *Clin. Chem.* 33, 1364–1367.
- Ortego-Centeno, N., Munoz-Torres, M., Jodar, E., Hernandez-Quero, J., Jurado-Duce, A., de la Higuera Torres-Puchol, J., 1997. Effect of tobacco consumption on bone mineral density in healthy young males. *Calcif. Tissue Int.* 60, 496–500.
- Rapuri, P.B., Gallagher, J.C., Balhorn, K.E., Ryschon, K.L., 2000. Smoking and bone metabolism in elderly women. *Bone* 27, 429–436.
- Seyedin, S.M., Kung, V.T., Daniloff, Y.N., Hesley, R.P., Gomez, B., Nielsen, L.A., Rosen, H.N., Zuk, R.F., 1993. Immunoassay for urinary pyridinoline: The new marker of bone resorption. *J. Bone Miner. Res.* 8, 635–641.
- Steingrimsdottir, L., Gunnarsson, O., Indridason, O.S., Franzson, L., Sigurdsson, G., 2005. Relationship between serum parathyroid hormone levels, vitamin D sufficiency, and calcium intake. *JAMA* 294, 2336–2341.
- Szulc, P., Garnero, P., Claustrat, B., Marchand, F., Duboeuf, F., Delmas, P.D., 2002. Increased bone resorption in moderate smokers with low body weight: the Minos study. *J. Clin. Endocrinol. Metab.* 87, 666–674.
- Taylor, A.K., Linkhart, S.G., Mohan, S., Baylink, D.J., 1988. Development of a new radioimmunoassay for human osteocalcin: evidence for a midmolecule epitope. *Metabolism* 37, 872–877.