

UCLA

UCLA Previously Published Works

Title

Magnetic Resonance Imaging-Guided Biopsy in Active Surveillance of Prostate Cancer

Permalink

<https://escholarship.org/uc/item/98p1w1m7>

Journal

Journal of Urology, 207(4)

ISSN

0021-0005

Authors

Kinnaird, Adam

Yerram, Nitin K

O'Connor, Luke

et al.

Publication Date

2022-04-01

DOI

10.1097/ju.0000000000002343

Peer reviewed



Published in final edited form as:

J Urol. 2022 April ; 207(4): 823–831. doi:10.1097/JU.0000000000002343.

Magnetic Resonance Imaging-Guided Biopsy in Active Surveillance of Prostate Cancer

Adam Kinnaird^{1,2,3,4,*}, Nitin K. Yerram^{5,*}, Luke O'Connor⁵, Wayne Brisbane¹, Vidit Sharma¹, Ryan Chuang¹, Rajiv Jayadevan¹, Michael Ahdoot⁵, Michael Daneshvar⁵, Alan Priester⁶, Merdie Delfin¹, Elizabeth Tran¹, Danielle E. Barsa¹, Anthony Sisk⁷, Robert E. Reiter¹, Ely Felker⁸, Steve Raman⁸, Lorna Kwan¹, Peter L. Choyke⁹, Maria J. Merino¹⁰, Bradford J. Wood¹¹, Baris Turkbey⁹, Peter A. Pinto⁵, Leonard S. Marks^{1,†}

¹Department of Urology, David Geffen School of Medicine, UCLA, Los Angeles, California

²Division of Urology, Department of Surgery, University of Alberta, Edmonton, Alberta, Canada

³Alberta Centre for Urologic Research and Excellence (ACURE), Edmonton, Alberta, Canada

⁴Cancer Research Institute of Northern Alberta (CRINA), Edmonton, Alberta, Canada

⁵Urologic Oncology Branch, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

⁶Department of Bioengineering, UCLA, Los Angeles, California

⁷Department of Pathology & Laboratory Medicine, UCLA, Los Angeles, California

⁸Department of Radiological Sciences, UCLA, Los Angeles, California

⁹Molecular Imaging Program, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

¹⁰Laboratory of Pathology, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

¹¹Center for Interventional Oncology, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

Abstract

†Correspondence: Department of Urology, David Geffen School of Medicine, UCLA, Wasserman Bldg., 3rd Floor, Los Angeles, California 90095 (telephone: 310-794-3070; FAX: 310-794-0987; lmarks@mednet.ucla.edu).

*Equal study contribution.

Conflict of Interest: Dr. Marks is co-founder of Avenda Health. Dr. Priester is data scientist at Avenda Health.

Disclosures: NIH and Philips have a Cooperative Research and Development Agreement.

Licensed Patents/Royalties: Philips: NIH receives royalties for licensed patents from Philips and in turn, BW, BT, PP, PC receive resulting royalties from NIH. NIH may own intellectual property in the field (multiple patents and patents pending-available upon request).

Intellectual Property: NIH & BW.

Non-Endorsement Language: The content of this manuscript does not necessarily reflect the views, policies, or opinions of the U.S. Department of Health and Human Services. The mention of commercial products, their source, or their use in connection with material reported herein is not to be construed as an actual or implied endorsement of such products by the United States government. Opinions expressed are those of the authors, not necessarily the NIH nor the NCI.

Purpose: The underlying premise of prostate cancer active surveillance (AS) is that cancers likely to metastasize will be recognized and eliminated before cancer-related disease can ensue. Our study was designed to determine the prostate cancer upgrading rate when biopsy guided by magnetic resonance imaging (MRGBx) is used before entry and during AS.

Materials and Methods: The cohort included 519 men with low- or intermediate-risk prostate cancer who enrolled in prospective studies (NCT00949819 and NCT00102544) between February 2008 and February 2020. Subjects were preliminarily diagnosed with Gleason Grade Group (GG) 1 cancer; AS began when subsequent MRGBx confirmed GG1 or GG2. Participants underwent confirmatory MRGBx (targeted and systematic) followed by surveillance MRGBx approximately every 12 to 24 months. The primary outcome was tumor upgrading to GG3.

Results: Upgrading to GG3 was found in 92 men after a median followup of 4.8 years (IQR 3.1–6.5) after confirmatory MRGBx. Upgrade-free probability after 5 years was 0.85 (95% CI 0.81–0.88). Cancer detected in a magnetic resonance imaging lesion at confirmatory MRGBx increased risk of subsequent upgrading during AS (HR 2.8; 95% CI 1.3–6.0), as did presence of GG2 (HR 2.9; 95% CI 1.1e8.2). In men who upgraded GG3 during AS, upgrading was detected by targeted cores only in 27%, systematic cores only in 25% and both in 47%. In 63 men undergoing prostatectomy, upgrading from MRGBx was found in only 5 (8%).

Conclusions: When AS begins and follows with MRGBx (targeted and systematic), upgrading rate (GG3) is greater when tumor is initially present within a magnetic resonance imaging lesion or when pathology is GG2 than when these features are absent.

Abstract

Study Need and Importance: Active surveillance (AS) has become a preferred strategy for managing men with low-risk prostate cancer (PCa). Improved entry and followup criteria could help eliminate men up front who are destined to require active treatment and, during surveillance, identify those remaining men who progress. In most AS programs, men have entered based on an ultrasoundguided biopsy (USGBx) revealing Grade Group (GG) 1 PCa. However, in part because of the limitations of USGBx many men thus biopsied will ultimately exhibit PCa of increased risk and require active treatment. Biopsies employing guidance by magnetic resonance imaging (MRGBx) provide characterization of prostate pathology which is more accurate than USGBx. Thus, use of MRGBx is of increasing interest for men before entry and during AS.

What We Found: We studied a cohort of 519 men on AS with low- or intermediate-risk PCa who enrolled in prospective studies between February 2008 and February 2020. All men underwent confirmatory MRGBx (targeted and systematic), followed by surveillance MRGBx approximately every 24 months. We found that cancer detected in a magnetic resonance imaging (MRI)-visible lesion at confirmatory MRGBx increased risk of subsequent upgrading during AS, as did presence of GG2 (see figure). In men undergoing prostatectomy after AS, upgrading from the most recent MRGBx was found in only 8%.

Limitations: Limitations include lack of a comparator arm of men confirmed and followed with USGBx, and the results are from 2 centers with in-depth experience with MRGBx, potentially limiting generalizability.

Interpretation for Patient Care: When AS begins and follows with MRGBx (targeted and systematic), upgrading rate (\geq GG3) is greater when tumor is initially present within an MRI-visible lesion or when pathology is GG2 than when these features are absent.

Graphical Abstract




Figure. Probability of freedom from upgrading to \geq GG3 during AS of PCa after an MRI-guided confirmatory biopsy that showed no cancer (blue), GG1 (red) or GG2 (green).

Keywords

image-guided biopsy; prostatic neoplasms; observation; magnetic resonance imaging

Active surveillance (AS) has become the preferred strategy for managing men with low-risk prostate cancer (PCa).^{1,2} The proportion of men with low-risk PCa managed by AS increased from 15% in 2010 to 42% in 2015.³ When patients are selected and followed carefully, nearly all will avoid PCa-related disease in the near and intermediate term.^{4,5} In one study of 1,818 men with low-risk PCa who were followed in AS for many years, only 4 men died from PCa.⁶ However, the rate of definitive treatment in these men was 48% at 10 years.

Improved entry and followup criteria could help eliminate men up front who are destined to require active treatment and, during surveillance, identify those remaining men who progress. In most AS programs, men have entered on the basis of an ultrasound-guided biopsy (USGBx) revealing a cancer with Gleason Score of 6 (Gleason grade group [GG] 1), ie low risk, with periodic biopsies to confirm that low risk is maintained. However, because of the limitations of USGBx, many men thus biopsied will ultimately exhibit PCa of increased risk and require active treatment.^{6,7} Biopsies employing guidance by magnetic resonance imaging (MRGBx) provide characterization of prostate pathology which is more accurate than USGBx.^{8–10} Thus, use of MRGBx is of increasing interest for use in men before entry and during AS.^{11–15}

The 2 centers reporting here initiated AS programs at approximately the same period when their work with MRGBx began.^{8,16} During that interval, prospectively acquired databases structured around MRGBx have been maintained at each site. In contrast to other studies, all patients in this large, combined series began AS with a confirmatory MRGBx; all patients had MRGBx at scheduled intervals per protocol during years of AS; at each biopsy session both targeted and systematic biopsies were obtained; and followup is among the longest yet reported using serial MRGBx to detect upgrading.

METHODS

Study Design

This cohort study used prospectively collected data from 2 institutions (University of California, Los Angeles [UCLA; [NCT00949819](#)] and National Cancer Institute [NCI; [NCT00102544](#)]) between February 2008 and February 2020. Both studies were approved by the respective institutional review boards to evaluate the use of image fusion devices for targeted biopsies (IRB No. 11–002281). The electromagnetic tracking device used at NCI is now commercially available as the UroNav[®] platform (Philips Healthcare, Eindhoven, Noord-Brabant, Netherlands) and the mechanical tracking image-fusion device used at UCLA is commercially available as the Artemis platform (Eigen, Grass Valley, California). Investigator experience with the MRI-ultrasound fusion systems has been described previously.^{17,18}

Patients

All patients in this study had an initial diagnostic biopsy of GG1 PCa, obtained by various methods in community settings. Within 12 months of initial diagnostic biopsy subjects underwent confirmatory MRGBx. Patients were included in the analytical cohort if confirmatory biopsy revealed GG2 cancer and at least 1 followup surveillance biopsy was obtained. Patients were excluded if they received immediate PCa treatment, were followed up elsewhere, withdrew consent or died from a cause unrelated to PCa. A flow diagram of participants is shown in figure 1.

Imaging and Biopsy Methods

All patients underwent 3.0 Tesla multiparametric magnetic resonance imaging (MRI) as previously described.^{8,15}

All biopsies were performed using MRI-ultrasound fusion technology (Artemis at UCLA and UroNav or a pre-UroNav prototype at NCI). Each biopsy procedure combined conventional 12-core systematic sampling in addition to targeted biopsy directed to MRI visible lesions (at least 2 cores per Prostate Imaging–Reporting and Data System [PI-RADS[®]] 3–5 lesion).¹⁹ Targeted and systematic cores were obtained at each biopsy session per the protocol of each institution. MRI and pathology interpretations were performed by expert genitourinary radiologists and pathologists. Prior to the adoption of the PI-RADS scoring system, Likert scoring systems were used.^{20,21}

Followup

After confirmatory biopsy, patients were monitored with digital rectal examination and prostate specific antigen (PSA) testing approximately every 12 to 24 months. Patients also underwent multiparametric MRI and MRI-guided surveillance biopsies approximately every 12 to 24 months.

Outcomes

The main outcome of interest was pathological upgrading of PCa to GG3, a point at which AS is generally considered inappropriate. The secondary outcome was pathological upgrading to GG2. Whole mount pathology was used to determine the pathological GG in men undergoing radical prostatectomy after AS.

Statistical Analyses

We conducted survival analyses for upgrading to GG3 as well as GG2 and present upgrade-free probabilities and Kaplan-Meier curves. Log-rank test was used to test strata. We calculated upgrade-free probabilities and 95% Wald confidence intervals in SAS® (SAS, Cary, North Carolina). Continuous variables that were not normally distributed were tested between groups using the Wilcoxon rank-sum or median test as appropriate. We conducted Cox proportional hazard regression to calculate the hazard ratios of upgrading, controlling for patient, biopsy and MRI characteristics from the confirmatory biopsy. We *a priori* chose the following variables to include in the regression: age, ethnicity (nonHispanic White vs African American vs Asian vs other/unknown), family history of PCa, PSA, PSA density (≥ 0.15 vs <0.15 ng/ml/cc), maximum cancer core length, number of positive systematic cores (0 vs 1 vs ≥ 2), number of positive targeted cores (0 vs 1 vs ≥ 2) and GG.

RESULTS

Characteristics of Patients

Patient characteristics at time of MRI-guided confirmatory biopsy are shown in table 1. From February 2008 to January 2019, a total of 519 patients (UCLA, 330; National Institutes of Health [NIH], 189) met inclusion criteria, signed consent forms and were enrolled in the study.

MRI-Guided Confirmatory Biopsy

Of the 519 men enrolled, confirmatory biopsy revealed PCa $<GG2$ in 444 and GG2 in 75 (fig. 1). A median of 5 (IQR 4–6) targeted cores were obtained from subjects with MRI-visible lesions (362) and a median of 12 (IQR 12–13) systematic cores were taken from all men (table 1). Targeted and systematic cores were both required to detect highest GG (supplementary table 2, <https://www.jurology.com>).

Followup During Surveillance

Followup data were available for outcomes through February 2020. The median followup period was 4.8 years (IQR 3.1–6.5) after MRI-guided confirmatory biopsy (table 2). Patients

underwent a median of 1 (range 0–6) followup surveillance MRI study and a median of 2 (range 0–5) followup surveillance biopsy sessions within 5 years after confirmatory biopsy.

Risk of Upgrading to GG3

Upgrading to GG3 occurred in 92 men (table 2). The median followup time for men who did not upgrade GG3 was 5.1 years (IQR 3.5–6.7). Upgrade-free probability at 2, 5 and 7 years after confirmatory biopsy were 0.95 (95% CI 0.93–0.96), 0.85 (95% CI 0.81–0.88) and 0.77 (95% CI 0.72–0.82), respectively (fig. 2, upper panel). The upgrade-free probability at 5 years for men who had cancer detected in targeted cores or combined biopsy was 0.75 (95% CI 0.59–0.86) and 0.69 (95% CI 0.56–0.79), respectively (fig. 2, middle panel). For men with cancer detected on systematic cores only, or when no cancer was detected, the upgrade-free probability at 5 years was 0.89 (95% CI 0.79–0.94) and 0.93 (95% CI 0.87–0.97), respectively.

At confirmatory MRGBx cancer found in targeted but not systematic cores was associated with tumor upgrading to GG3 during AS (table 3). If tumor was identified in 1 targeted core (HR 2.75; 95% CI 1.25–6.03) or more than 1 targeted core (HR 3.38; 95% CI 1.65–6.91), men were more likely to upgrade during AS than if all targeted cores were cancer-free. Presence of cancer in systematic cores was not independently associated with tumor upgrading. Family history of PCa (HR 1.58; 95% CI 1.02–2.44) and GG2 identified at any confirmatory biopsy (HR 2.93; 95% CI 1.05–8.19) were independently associated with upgrading to GG3. At followup surveillance biopsies upgrading to GG3 was detected by only targeted cores in 27%, only systematic cores in 25% and both in 47% (table 2).

Risk of Upgrading to GG2

Upgrading to GG2 occurred in 164 men. The median followup time for men who did not upgrade GG2 was 4.7 years (IQR 3.3–6.3). Upgrade-free probability at 2, 5 and 7 years after confirmatory biopsy was 0.83 (95% CI 0.80–0.87), 0.62 (95% CI 0.56–0.67) and 0.55 (95% CI 0.49–0.61), respectively (see supplementary figure, <https://www.jurology.com>, upper panel).

At confirmatory MRGBx cancer found in targeted but not systematic cores was associated with tumor upgrading to GG2 during AS (supplementary table 1, <https://www.jurology.com>). If tumor was identified in 1 targeted core (HR 2.07; 95% CI 1.17–3.65) or more than 1 targeted core (HR 1.90; 95% CI 1.10–3.29), men were more likely to upgrade during AS than if all targeted cores were cancer-free. Presence of cancer in systematic cores was not independently associated with tumor upgrading.

Radical Prostatectomy after AS

Of the 519 men entering AS after confirmatory MRGBx, 116 subsequently underwent active treatment. Sixty-seven men underwent radical prostatectomy a median interval of 30.5 months (IQR 17.9, 51.6) after confirmatory MRGBx. Study of excised prostates via whole mount pathology was available in 63 men (94%; table 4). In 58 men (92%), final GG was the same or lower than that found at the preceding MRGBx. Upgrading at surgery (GG3) was not associated with duration of AS ($p=0.17$). The time to surgery was not related to

Grade Group (GG1 or GG2) at confirmatory MRGBx ($p=0.11$). At final pathology, 11 men (15%) had GG4–5 cancer and 2 (3%) had positive lymph nodes.

DISCUSSION

The present work supports and expands that of other AS investigators.^{1,6,11–14} First, AS proved to be generally safe and effective throughout years of followup. The 5-year upgrading-free probability (GG3) was 85%. Second, the value of confirmatory MRGBx as a starting point for AS is established, since some 10% of men who were thought to be candidates for AS on the basis of conventional USGBx were found on MRGBx to have PCa GG3. Third, tumor detected within an MRI lesion deserves special attention during AS, since the chances of tumor upgrading were increased threefold over others. Fourth, the role of both targeted and systematic biopsies during surveillance is reinforced, since 25% of GG3 cancers seen during followup were diagnosed only by systematic biopsy. Fifth, with only an 8% upgrading rate at prostatectomy, MRGBx reliably characterizes the tumor. And sixth, we confirm the need for vigilance when GG2 lesions are surveilled because of a nearly threefold increase in upgrade rate compared to men entering with GG1.

The primary outcome was upgrading to GG3. Men with GG1 or GG2 tumors are considered eligible for AS, but the presence of GG3 excludes AS from consideration.^{2,15} Men with GG3 disease have significantly greater rates of biochemical recurrence and overall mortality compared to men with GG2.^{22,23} Thus, MRGBx was used here to exclude men with GG3 at the start of AS and to discontinue AS when GG3 was found during surveillance. Since accurate characterization of prostate pathology is critical for men undergoing AS, and since MRGBx is more sensitive for cancer detection than USGBx,⁹ these data support the use of MRGBx for selection and followup of men undergoing AS.

The date of confirmatory biopsy was used as the start of AS, differentiating the present study from others which have used initial diagnostic biopsy (any method) as the starting point. When entry biopsies are performed via ultrasound guidance alone, later upgrading during AS may be as high as 30%.⁶ In the present study only 18% of men with GG1 pathology at confirmatory MRGBx experienced upgrading to GG3 lesions. The results of the present study support the importance of MRGBx to confirm findings of conventional biopsy before entry into AS, and further suggest that such a confirmatory biopsy may be the appropriate starting point for AS.

A cancer-containing targeted core from an MRI lesion at confirmatory biopsy (GG1 or GG2) nearly tripled the risk of tumor upgrading compared to finding no tumor or tumor only in a systematic core. The overall risk of tumor upgrading to GG3 after 7 years of AS was ~40% if cancer was found in a targeted core compared to only ~10% if cancer was found in a systematic core or no cancer was identified (fig. 2).

For men in AS, systematic biopsies provide important information beyond that obtained from targeted biopsy, as suggested by Klotz and colleagues and confirmed herein.²⁴ In the present report, systematic cores detected upgrading missed by targeted biopsy in 25% of men. Tumor present outside MRI lesions may be the result of misregistration on fusion

biopsy, segmentation error, operator or technical inaccuracies, MRI failing to identify cancer or any combination thereof.²⁵ Further, combination of targeted and systematic cores most accurately predicted final tumor grade found at radical prostatectomy.⁸ In the present cohort, 63 men underwent radical prostatectomy (and had available whole mount slides for analysis) after elimination from AS; the upgrading rate in this group from prior MRGBx to final pathology was only 8% (vs 56% by systematic cores only, supplementary table 3, <https://www.jurology.com>) compared to historical rates of 20%–30% seen with USGBx.²⁶

While an adverse MRI change during AS (increased PI-RADS or PRECISE scoring) has been associated with tumor upgrading,²⁷ omission of surveillance biopsy based on a stable MRI may leave high-grade tumors undetected. In previous work, AS by serial MRI alone (without biopsy) failed to detect 17%–22% of tumor upgrades.^{11,13} These data are in keeping with the present study in which 14% of men with negative MRIs upgraded during AS. These findings may be due to the failure of MRI to visualize clinically significant cancers in as many as 15%–25% of cases.^{28–30} The possibility of a falsely negative MRI adds to the importance of combining systematic and targeted biopsy.

The present study, because it included men with GG2, adds information about upgrading for men entering AS with intermediate-risk pathology. Here the finding of GG2 at confirmatory biopsy was independently associated with increased risk of upgrading to GG3. Thus is confirmed an earlier report showing that men with GG2 upgrade to GG3 more often than men with GG1 by severalfold.¹⁵ Others have reported increased PCa progression and even metastasis during AS in men with GG2 initially diagnosed by USGBx.^{1,4} These data indicate that even with the accuracy of MRGBx, increased vigilance during surveillance of men with GG2 is warranted.

Since results were similar at the 2 sites despite use of different biopsy platforms, the findings appear to be unrelated to type of fusion device employed and may apply to other MRI-targeted methods, as well.

Limitations of this study include lack of a comparator arm of men confirmed and followed with USGBx. Relatively few men with GG2 were included (75); 80% of subjects were Caucasian. No attempt was made to stratify GG2 tumors, (eg percent pattern 4). Genomics were not studied. Men were able to choose definitive treatment at any time. And because the 2 centers have longstanding, in-depth experience with MRGBx, the results may not yet be generalizable. Despite the limitations, the prospective data herein provide evidence that for men in AS MRGBx can provide a degree of accuracy about prostate pathology that has not been available with USGBx.

CONCLUSIONS

When AS begins and follows with MRGBx (targeted and systematic), upgrading rate (GG3) is greater when tumor is initially present within an MRI lesion or when pathology is GG2 than when these features are absent.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Support:

Supported in part by R01CA218547, R01CA195505, the Intramural Research Program of the National Cancer Institute, the Center for Interventional Oncology, intramural NIH Grants Z1A CL040015, 1ZIDBC011242, UL1TR000124 from University of California, Los Angeles Clinical and Translational Science Institute, the University Hospital Foundation in Edmonton, Alberta, and generous donation from Mr. Leo Broks.

Abbreviations and Acronyms

AS	active surveillance
GG	Gleason grade group
MRGBx	magnetic resonance imaging-guided biopsy
MRI	magnetic resonance imaging
NCI	National Cancer Institute
NIH	National Institutes of Health
PCa	prostate cancer
PI-RADS®	Prostate Imaging eReporting and Data System
PSA	prostate specific antigen
UCLA	University of California, Los Angeles
USGBx	ultrasound-guided biopsy

REFERENCES

1. Klotz L, Vesprini D, Sethukavalan P et al. : Long-term follow-up of a large active surveillance cohort of patients with prostate cancer. *J Clin Oncol* 2014; 33: 272. [PubMed: 25512465]
2. Chen RC, Rumble RB, Loblaw DA et al. : Active surveillance for the management of localized prostate cancer (Cancer Care Ontario Guideline): American Society of Clinical Oncology clinical practice guideline endorsement. *J Clin Oncol* 2016; 34: 2182. [PubMed: 26884580]
3. Mahal BA, Butler S, Franco I et al. : Use of active surveillance or watchful waiting for low-risk prostate cancer and management trends across risk groups in the United States, 2010–2015. *JAMA* 2019; 321: 704. [PubMed: 30743264]
4. Maggi M, Cowan JE, Fasulo V et al. : The long-term risks of metastases in men on active surveillance for early stage prostate cancer. *J Urol* 2020; 204: 1222. [PubMed: 33157570]
5. Bokhorst LP, Valdagni R, Rannikko A et al. : A decade of active surveillance in the PRIAS study: an update and evaluation of the criteria used to recommend a switch to active treatment. *Eur Urol* 2016; 70: 954. [PubMed: 27329565]
6. Tosoian JJ, Mamawala M, Epstein JI et al. : Active surveillance of grade group 1 prostate cancer: long-term outcomes from a large prospective cohort. *Eur Urol* 2020; 77: 675. [PubMed: 31918957]
7. Klotz L, Vesprini D, Sethukavalan P et al. : Long-term follow-up of a large active surveillance cohort of patients with prostate cancer. *J Clin Oncol* 2015; 33: 272. [PubMed: 25512465]

8. Ahdoot M, Wilbur AR, Reese SE et al. : MRI-targeted, systematic, and combined biopsy for prostate cancer diagnosis. *N Engl J Med* 2020; 382: 917. [PubMed: 32130814]
9. Kasisivsanathan V, Rannikko AS, Borghi M et al. : MRI-targeted or standard biopsy for prostate cancer diagnosis. *N Engl J Med* 2018; 378: 1767. [PubMed: 29552975]
10. Norris JM, Kinnaird A, Margolis DJ et al. : Developments in MRI-targeted prostate biopsy. *Curr Opin Urol* 2020; 30: 1. [PubMed: 31644434]
11. Chesnut GT, Vertosick EA, Benfante N et al. : Role of changes in magnetic resonance imaging or clinical stage in evaluation of disease progression for men with prostate cancer on active surveillance. *Eur Urol* 2020; 77: 501. [PubMed: 31874726]
12. Klotz L, Pond G, Loblaw A et al. : Randomized study of systematic biopsy versus magnetic resonance imaging and targeted and systematic biopsy in men on active surveillance (ASIST): 2-year postbiopsy follow-up. *Eur Urol* 2019; 77: 311. [PubMed: 31708295]
13. Chu CE, Loneragan PE, Washington SL et al. : Multiparametric magnetic resonance imaging alone is insufficient to detect grade reclassification in active surveillance for prostate cancer. *Eur Urol* 2020; 78: 515. [PubMed: 32631744]
14. Stavrinides V, Giganti F, Trock B et al. : Five-year outcomes of magnetic resonance imaging-based active surveillance for prostate cancer: a large cohort study. *Eur Urol* 2020; 78: 443. [PubMed: 32360049]
15. Jayadevan R, Felker ER, Kwan L et al. : Magnetic resonance imaging-guided confirmatory biopsy for initiating active surveillance of prostate cancer. *JAMA Netw Open* 2019; 2: e1911019. [PubMed: 31509206]
16. Filson CP, Natarajan S, Margolis DJA et al. : Prostate cancer detection with magnetic resonance-ultrasound fusion biopsy: the role of systematic and targeted biopsies. *Cancer* 2016; 122: 884. [PubMed: 26749141]
17. Marks L, Young S and Natarajan S: MRI—ultrasound fusion for guidance of targeted prostate biopsy. *Curr Opin Urol* 2013; 23: 43. [PubMed: 23138468]
18. Natarajan S, Marks LS, Margolis DJA, et al. : Clinical application of a 3D ultrasound-guided prostate biopsy system. *Urol Oncol* 2011; 29: 334. [PubMed: 21555104]
19. Weinreb JC, Barentsz JO, Choyke PL et al. : PI-RADS prostate imaging—reporting and data system: 2015, version 2. *Eur Urol* 2016; 69: 16.
20. Afshari Mirak S, Bajgirani AM, Hosseiny M et al. : PD64–06 Comparison of performance of PI-RADSV2 and a quantitative PI-RADSV1 based protocol IN 3T multiparametric MRI for detection, grading and staging of prostate cancer using whole mount histopathology as reference standard IN 569 patients. *J Urol, suppl.*, 2019; 201: e1183.
21. Gaur S, Harmon S, Mehralivand S et al. : Prospective comparison of PI-RADS version 2 and qualitative in-house categorization system in detection of prostate cancer. *J Magn Reson Imaging* 2018; 48: 1326. [PubMed: 29603833]
22. Erickson A, Sandeman K, Lahdensuo K et al. : New prostate cancer grade grouping system predicts survival after radical prostatectomy. *Hum Pathol* 2018; 75: 159. [PubMed: 29447924]
23. Epstein JI, Zelefsky MJ, Sjöberg DD et al. : A contemporary prostate cancer grading system: a validated alternative to the Gleason score. *Eur Urol* 2016; 69: 428. [PubMed: 26166626]
24. Klotz L, Loblaw A, Sugar L et al. : Active surveillance magnetic resonance imaging study (ASIST): results of a randomized multicenter prospective trial. *Eur Urol* 2019; 75: 300. [PubMed: 30017404]
25. Kinnaird A, Sharma V, Chuang R et al. : Risk of prostate cancer after a negative magnetic resonance imaging guided biopsy. *J Urol* 2020; 204: 1180. [PubMed: 32614257]
26. Godtman RA, Schafferer M, Pihl CG et al. : Long-term outcomes after deferred radical prostatectomy in men initially treated with active surveillance. *J Urol* 2018; 200: 779. [PubMed: 29730198]
27. O'Connor LP, Wang AZ, Yerram NK et al. : Changes in magnetic resonance imaging using the prostate cancer radiologic estimation of change in sequential evaluation criteria to detect prostate cancer progression for men on active surveillance. *Eur Urol Oncol* 2020; 4: 227. [PubMed: 33867045]

28. Ahmed HU, El-Shater Bosaily A, Brown LC et al. : Diagnostic accuracy of multi-parametric MRI and TRUS biopsy in prostate cancer (PROMIS): a paired validating confirmatory study. *Lancet* 2017; 389: 815. [PubMed: 28110982]
29. Lovegrove CE, Miah S, El-Shater Bosaily A et al. : Comparison of transrectal ultrasound biopsy to transperineal template mapping biopsies stratified by multiparametric magnetic resonance imaging score in the PROMIS trial. *J Urol* 2019; 203: 100. [PubMed: 31335254]
30. Le JD, Tan N, Shkolyar E et al. : Multifocality and prostate cancer detection by multiparametric magnetic resonance imaging: correlation with whole-mount histopathology. *Eur Urol* 2015; 67: 569. [PubMed: 25257029]




Figure 1.
Flow diagram of participants.




Figure 2. Probability of upgrading-free survival to GG3 during AS of PCa after an MRI-guided confirmatory biopsy. *Upper panel*, overall probability of upgrading-free survival. *Middle panel*, probability of upgrading-free survival stratified by cancer-containing core type at confirmatory biopsy (*black*, negative systematic and negative targeted cores; *blue*, positive systematic and negative targeted cores; *yellow*, negative systematic and positive targeted cores; *green*, positive systematic and positive targeted cores). *Lower panel*, probability of

upgrading-free survival stratified by GG at confirmatory biopsy (*blue*, no cancer; *red*, GG1; *green*, GG2). *Sys*, systematic biopsy.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.

Baseline patient characteristics

	Total	UCLA	NIH	p Value
Total No. pts	519	330	189	
Mean yrs age (SD)	62.2 (7.3)	62.8 (7.6)	61.2 (6.6)	0.02
No. ethnicity (%):				<0.01
NonHispanic White	414 (80)	255 (77)	159 (84)	
African American	41 (8)	21 (6)	20 (11)	
Asian	24 (5)	20 (6)	4 (2)	
Other/unknown	40 (8)	34 (10)	6 (3)	
No. family history of PCa (%)	145 (28)	79 (24)	66 (35)	<0.01
No. abnormal rectal exam (%)	12 (2)	12 (4)	-	-
Median ng/ml PSA (IQR)	5.2 (3.0–8.1)	5.6 (2.7–8.3)	5.0 (3.7–7.2)	0.57
Median cc MRI volume (IQR)	48 (35–68)	48 (35–71)	48 (37–66)	0.77
No. ng/ml/cc PSA density (%):				0.62
<0.15	417 (80)	263 (80)	154 (81)	
0.15	102 (20)	67 (20)	35 (19)	
No. MRI score (%):				0.15
Negative (1 or 2)	156 (30)	103 (31)	53 (28)	
3	226 (44)	137 (42)	89 (47)	
4	105 (20)	74 (22)	31 (16)	
5	31 (6)	16 (5)	15 (8)	
Median mm max Ca core length of highest tumor grade (IQR)	2.5 (1.0–4.2)	2.0 (1.0–4.0)	3.0 (1.3–5.0)	0.18
Median No. systematic cores taken (IQR)	12 (12–13)	12 (11–13)	12 (12–14)	<0.01
No. pos systematic cores (%):				0.17
0	230 (45)	146 (45)	84 (44)	
1	146 (29)	99 (31)	47 (25)	
>2	135 (26)	24 (7)	58 (31)	
Median No. targeted cores taken (IQR) *	5 (4–6)	5 (4–6)	4 (4–6)	0.77
No. pos targeted cores (%): *				<0.01
0	212 (59)	145 (64)	67 (49)	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

	Total	UCLA	NIH	p Value
1	61 (17)	39 (17)	22 (16)	
2	88 (24)	41 (18)	47 (35)	
No. GG (%):				0.24
No Ca	181 (35)	116 (35)	67 (35)	
1	263 (51)	173 (52)	89 (47)	
2	75 (14)	41 (12)	33 (17)	

Values in bold are statistically significant. Baseline Is considered at time of MRI-guided confirmatory biopsy.

* From men with MRI score 3–5 only.

Summary of AS followup

Table 2.

	Total	UCLA	NIH	p Value
Total No. pts	519	330	189	
Median yrs length followup (IQR)	4.8 (3.1–6.5)	4.6 (3.0–6.4)	5.0 (3.3–6.5)	0.22
Followup MRIs within 5 yrs:				<0.01
Median total No. (range)	1 (0–6)	1 (0–5)	3 (0–6)	
Median yrs between MRIs (IQR)	1.1 (1.0–1.7)	2.0 (1.2–2.1)	1.1 (1.0–1.3)	
Followup biopsy sessions within 5 yrs:				<0.01
Median total No. (range)	2 (0–5)	2 (1–5)	1 (0–4)	
Median yrs between MRIs (IQR)	1.2 (1.0–2.0)	1.1 (1.0–2.0)	1.7 (1.1–2.0)	
No. upgrading at followup biopsy:				< 0.01
Upgrading to GG3	92	39	53	
GG3	47	22	25	
GG4	33	13	20	
GG5	11	4	7	
No. upgrading to GG2 (among <GG2):* †	164	93	71	< 0.01
GG2	115	73	42	
GG3	26	11	15	
GG4	13	6	7	
GG5	9	3	6	
No. pts upgrading GG3 by biopsy core type (%): ‡				< 0.01
Targeted core only	25 (2)	17 (44)	8 (15)	
Systematic core only	23 (25)	19 (49)	4 (8)	
Both targeted and systematic cores	43 (47)	3 (8)	40 (75)	
Data missing	1 (1)	0 (0)	1 (2)	
No. pts upgrading GG2 by biopsy core type (%): ‡				< 0.01
Targeted core only	50 (30)	41 (44)	9 (13)	
Systematic core only	49 (30)	40 (43)	9 (13)	
Both targeted and systematic cores	63 (38)	12 (13)	51 (72)	
Data missing	2 (1)	0 (0)	2 (3)	

	Total	UCLA	NIH	p Value
No. received treatment:	116	73	43	< 0.001
Surgery	67	36	31	
Radiation therapy	30	18	12	
Focal therapy	15	15	0	
Other	4	4	0	

Values in bold are statistically significant.

* Of 444 cases.

† Of 92 cases.

‡ Of 164 cases.

Table 3.

Risk factors of upgrading to GG3 after MRI-guided confirmatory biopsy in 506 cases

	HR (95% CI)
Family history of PCa	1.58 (1.02–2.44)
No. pos systematic cores (reference, 0):	
1	0.95 (0.45–2.00)
2	1.27 (0.65–2.50)
No. pos targeted cores (reference, 0):	
Neg MRI (no targeted cores)	1.46 (0.79–2.70)
1	2.75 (1.25–6.03)
2	3.38 (1.65–6.91)
GG (reference, no cancer):	
1	1.40 (0.57–3.44)
2	2.93 (1.05–8.19)

Values in bold are statistically significant.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4.

Radical prostatectomy outcomes after MRI-guided confirmatory biopsy

	Total
Total No. pts	63
No. final pathology after prostatectomy (%):	
No Ca	1 (2)
GG1	3 (5)
GG2	41 (65)
GG3	7 (11)
GG4	8 (13)
GG5	3 (5)
No. change in final pathology from most recent MRI-guided biopsy (%):	
Upgrade	5 (8)
Downgrade	23 (37)
No change	35 (56)
Median mos time to surgery after a confirmatory biopsy that diagnosed (IQR)*:	
Any GG	30.5 (17.9, 51.6)
No Ca	42.9 (22.6, 65.0)
GG1	33.0 (19.1, 54.6)
GG2	22.6 (16.0, 30.8)
Median mos time to surgery for final pathology (IQR)*:	
<GG3	28.2 (16.6, 49.4)
GG3	35.4 (30.0, 54.8)
No. pathological pos lymph nodes at surgery (%)*	2 (3)

* Of 61 patients.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript