

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Observation of the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay.

Permalink

<https://escholarship.org/uc/item/98r9s6w8>

Journal

European Physical Journal C: Particles and Fields, 84(10)

ISSN

1434-6044

Authors

Hayrapetyan, A

Tumasyan, A

Adam, W

et al.

Publication Date

2024

DOI

10.1140/epjc/s10052-024-13114-9

Peer reviewed



Observation of the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay

CMS Collaboration*

CERN, 1211 Geneva 23, Switzerland

Received: 29 January 2024 / Accepted: 8 July 2024
© CERN for the benefit of the CMS collaboration 2024

Abstract Using proton–proton collision data corresponding to an integrated luminosity of 140 fb^{-1} collected by the CMS experiment at $\sqrt{s} = 13 \text{ TeV}$, the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay is observed for the first time, with a statistical significance exceeding 5 standard deviations. The relative branching fraction, with respect to the $\Lambda_b^0 \rightarrow \psi(2S) \Lambda$ decay, is measured to be $\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+)/\mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S) \Lambda) = [3.38 \pm 1.02 \pm 0.61 \pm 0.03]\%$, where the first uncertainty is statistical, the second is systematic, and the third is related to the uncertainties in $\mathcal{B}(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$ and $\mathcal{B}(\Xi^- \rightarrow \Lambda \pi^-)$.

1 Introduction

Multibody decays of beauty hadrons present a rich laboratory to search for intermediate resonances in the decay products. When decay products contain a charmonium state, such intermediate resonances could decay into a charmonium meson and a hadron, which could be a manifestation of their exotic nature. An important turning point in exotic spectroscopy was achieved at the LHC, when the LHCb Collaboration reported the observation of statistically significant $J/\psi p$ pentaquark-like structures in the decay of the lightest beauty baryon $\Lambda_b^0 \rightarrow J/\psi p K^-$ [1]. Various interpretations of these structures have been proposed [2, 3], including tightly bound hidden-charm $[c\bar{c}uud]$ pentaquark states [4, 5], loosely bound molecular baryon-meson states [6–8], or being due to a double triangle singularity [9]. More recently, additional exotic states have been reported by LHCb in the decays $\Lambda_b^0 \rightarrow J/\psi p K^-$ [10], $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ [11], $B_s^0 \rightarrow J/\psi p \bar{p}$ [12], and $B^- \rightarrow J/\psi \Lambda \bar{p}$ [13]. Up to now, the hidden-charm pentaquark candidates have been reported only in $J/\psi p$ and $J/\psi \Lambda$ systems. Investigation of other channels with heavier baryons in the decay products, such as Ξ^- and Ω^- , could unveil the existence of doubly or triply strange pentaquarks [14, 15].

* e-mail: cms-publication-committee-chair@cern.ch (corresponding author)

In this paper, we report on the search for the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay, where the $J/\psi \rightarrow \mu^+ \mu^-$, $\Xi^- \rightarrow \Lambda \pi^-$, and $\Lambda \rightarrow p \pi^-$ channels are used to reconstruct the intermediate decay products. Charge-conjugate states are implied throughout the text. The measurement of the ratio of branching fractions

$$\mathcal{R} \equiv \frac{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+)}{\mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S) \Lambda)} = \frac{N(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+)}{N(\Lambda_b^0 \rightarrow \psi(2S) \Lambda)} \times \frac{\epsilon_{\psi(2S) \Lambda} \mathcal{B}(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)}{\epsilon_{J/\psi \Xi^- K^+} \mathcal{B}(\Xi^- \rightarrow \Lambda \pi^-)} \quad (1)$$

is also reported, where N is the measured Λ_b^0 yield and ϵ is the total efficiency. The normalization channel is chosen to be $\Lambda_b^0 \rightarrow \psi(2S) \Lambda$, with the subsequent $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ and $J/\psi \rightarrow \mu^+ \mu^-$ decays, because of its similar decay topology and kinematics to the signal decay, leading to the reduction of many systematic uncertainties. The branching fractions of the intermediate decays $\mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)$ and $\mathcal{B}(\Lambda \rightarrow p \pi^-)$ cancel in the ratio. Invariant mass distributions of the three two-body combinations for the signal channel are also presented in order to look for intermediate resonances.

The analysis uses proton–proton (pp) collision data recorded by the CMS experiment in 2016–2018, at $\sqrt{s} = 13 \text{ TeV}$, corresponding to an integrated luminosity of 140 fb^{-1} [16–18]. Tabulated results are provided in the HEPData record for this analysis [19].

2 The CMS detector and simulated event samples

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter, and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. Forward

calorimeters extend the pseudorapidity coverage provided by the barrel and endcap detectors. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [20].

Muons are measured in the pseudorapidity range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. Matching muons to tracks measured in the silicon tracker results in a transverse momentum (p_T) resolution for muons with p_T up to 100 GeV of 1% in the barrel and 3% in the endcaps. The silicon tracker used in 2016 measured charged particles within the range $|\eta| < 2.5$. For nonisolated particles of $1 < p_T < 10$ GeV and $|\eta| < 1.4$, the track resolutions were typically 1.5% in p_T and 25–90 μm in the transverse impact parameter [21]. At the start of 2017, a new pixel detector was installed [22]; the upgraded tracker measured nonisolated particles of $1 < p_T < 10$ GeV up to $|\eta| < 3$ with typical resolutions of 1.5% in p_T and 20–75 μm in the transverse impact parameter [23].

Events of interest are selected using a two-tiered trigger system [24]. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4 μs [25]. The second level, known as the high-level trigger (HLT), consists of a farm of computing processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage. All events used in this analysis are selected by a set of triggers requiring two identified muons of opposite charge plus an additional track to form a secondary vertex, displaced from the region of the pp interactions. The trigger demanded for each muon to have $p_T > 4$ GeV and to pass within 2 cm of the beam axis. The dimuon system was required to have $p_T > 6.9$ GeV, invariant mass between 2.9 and 3.3 GeV, a vertex fit probability greater than 10%, a separation of the secondary vertex relative to the beam axis in the transverse plane larger than 3 standard deviations (s.d.), and a cosine of the angle in the transverse plane between the dimuon momentum vector and the vector joining the beam axis and the dimuon vertex greater than 0.9. The additional track was required to have $p_T > 0.8$ (1.2) GeV and an impact parameter with respect to the beam axis greater than 0 (2) s.d., for data collected in 2016 (2017–2018). Finally, the two muons and the additional track were required to originate from the same vertex with a χ^2 per degree of freedom (dof) less than 10.

Monte Carlo (MC) simulated event samples are generated with PYTHIA v8.240 [26] using the CP5 underlying event tune [27]. The EVTGEN 1.6.0 [28] program models

the beauty baryon decays $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ and $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ with a phase space decay model, followed by the $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ and $J/\psi \rightarrow \mu^+ \mu^-$ decays. Final-state radiation is included in EVTGEN using PHOTOS 3.61 [29]. The events are then passed through a detailed GEANT4-based simulation [30] of the CMS detector, including also the decays of long-lived hyperons $\Xi^- \rightarrow \Lambda \pi^-$ and $\Lambda \rightarrow p \pi^-$, followed by the trigger and reconstruction algorithms identical to those used for the collision data. The simulation includes additional interactions due to multiple pp collisions in each bunch crossing, with the same distribution as observed in the experiment.

3 Event reconstruction and selection

The reconstruction for all the decays considered in this analysis starts by finding two muons of opposite charge, which must match those that triggered the event readout and pass the soft-muon identification criteria [31]. The offline selection for both muons requires $p_T(\mu^\pm) > 3$ GeV, $|\eta(\mu^\pm)| < 2.4$, χ^2 fit probability to a common dimuon vertex $P_{\text{vtx}}(\mu^+ \mu^-) > 1\%$, and dimuon invariant mass $2.9 < m(\mu^+ \mu^-) < 3.3$ GeV.

The $\Lambda \rightarrow p \pi^-$ candidates are selected from displaced two-prong vertices as described in Ref. [32]. The track with the higher momentum is assumed to be the proton one, and together with the pion track it is fit to a common vertex with their invariant mass constrained to the known Λ hyperon mass of $m_{\text{PDG}}(\Lambda) = 1115.683$ MeV [33]. The χ^2 fit probability for the Λ vertex is required to be $P_{\text{vtx}}(p \pi^-) > 1\%$.

For the signal channel, to form the $\Xi^- \rightarrow \Lambda \pi^-$ candidates, an additional high-purity [21] track assumed to be a pion is selected with $p_T > 0.2$ GeV. This track and the selected Λ candidate are then fit to a common vertex with the $\Lambda \pi^-$ mass constrained to the known Ξ^- hyperon mass of $m_{\text{PDG}}(\Xi^-) = 1321.71$ MeV [33]. To form the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ candidate, a high-purity track is chosen with an assigned kaon mass and $p_T(K^+) > 1.2$ GeV, which aligns with the HLT p_T requirement. The final reconstruction step in the signal channel is the $\mu^+ \mu^- \Xi^- K^+$ vertex fit with a χ^2 probability above 1%, where the dimuon mass is constrained to the world-average J/ψ meson mass of 3096.9 MeV [33].

For the normalization channel, two high-purity tracks of opposite charges with $p_T > 0.4$ GeV, assumed to be pions from the $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ decay, are selected. One of them is required to have $p_T > 1.2$ GeV to match the HLT p_T requirement. The Λ_b^0 candidates are obtained by a vertex fit of the $\mu^+ \mu^- \pi^+ \pi^- \Lambda$ system with a $J/\psi \rightarrow \mu^+ \mu^-$ mass constraint, as for the signal channel. The invariant mass of the $J/\psi \pi^+ \pi^-$ candidates is required to be in the range $3.60 < m(J/\psi \pi^+ \pi^-) < 3.95$ GeV.

From all reconstructed pp collision points in each event, the primary vertex (PV) is chosen as the one with the smallest Λ_b^0 pointing angle, which is the angle between the momentum of the Λ_b^0 candidate and the vector from the PV to the reconstructed Λ_b^0 candidate vertex. If any of the tracks used in the Λ_b^0 candidate reconstruction were included in the fit of the chosen PV, they are removed, and the PV is refitted.

Selection criteria for the signal channel $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ are optimized using the Punzi figure of merit [34]. The signal efficiency is evaluated using simulated event samples. Estimation of the background yield involves combining the collision data from the Λ_b^0 mass sideband, excluding the signal region which spans twice the mass resolution around the known Λ_b^0 mass. Additionally, the wrong-sign candidates ($J/\psi \Xi^- K^-$ and $J/\psi \Xi^+ K^+$) from the full mass range are included, after ensuring that the mass distribution of the wrong-sign candidates matches that of the correct-sign ones. Combining these two background sources reduces the impact of the statistical uncertainty in the optimization procedure. The variables used in the optimization include the p_T of all decay products; the flight length significance in the transverse plane of the Λ_b^0 , Λ , and Ξ^- baryon candidates and the corresponding pointing angles; the impact parameter significance with respect to the PV in the transverse plane for the tracks; the vertex fit probabilities; and the mass windows for hyperon candidates. The order of the cuts is determined randomly, and in several rounds of optimization this order was different each time; all rounds have converged to the same final set of optimized cuts. The resulting criteria are summarized in Table 1. The background is reduced by a factor of 15 after the optimization, whereas the signal efficiency is 70% of the initial selection described above. The selection criteria in the normalization channel $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ are chosen to be the same, wherever possible, as in the signal channel, to reduce the systematic uncertainties. The $J/\psi\pi^+\pi^-$ mass is required to be within 11.1 MeV of the known $\psi(2S)$ meson mass of 3686.1 MeV [33], which corresponds to approximately 2.5 times the mass resolution.

For the measurement of \mathcal{R} defined in Eq. (1), the pion from the Ξ^- decay is required to have $p_T > 0.4$ GeV. Additionally, the HLT requirements are repeated offline by requiring $p_T(\mu) > 4$ GeV, $p_T(J/\psi) > 6.9$ GeV, $P_{\text{vtx}}(\mu^+\mu^-) > 5\%$, and track (kaon for the signal channel, the harder of the two pions in the normalization channel) impact parameter above 2 s.d. with respect to the PV. These extra criteria ensure that events from potentially inadequately modeled phase space regions are avoided, as the reliability of the efficiency evaluation from simulated samples in those regions is questionable. Nevertheless, the reconstruction algorithm works reliably in those regions, and thus the corresponding events are used to study the mass distribution, as discussed in the following section.

Table 1 Optimized selection criteria for the signal decay mode $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$. The first two requirements are applied using the momenta before the corresponding mass constraint

Variable	Selection
$ m(p\pi^-) - m_{\text{PDG}}(\Lambda) $	< 8 MeV
$ m(\Lambda\pi^-) - m_{\text{PDG}}(\Xi^-) $	< 6 MeV
$p_T(\Lambda_b^0)$	> 11.5 GeV
$p_T(J/\psi)$	> 6.5 GeV
$p_T(\Xi^-)$	> 2.6 GeV
$p_T(\Lambda)$	> 2.2 GeV
$p_T(K^+)$	> 1.2 GeV
$\mu^+\mu^-\Xi^-K^+$ vertex fit probability	$> 5\%$
$\Lambda\pi^-$ vertex fit probability	$> 5\%$
$p\pi^-$ vertex fit probability	$> 1\%$
Ξ^- vertex displacement from Λ_b^0 vertex	> 3 s.d.
Λ vertex displacement from Ξ^- vertex	> 0 s.d.
Λ_b^0 vertex displacement from PV	> 3 s.d.
Angle between Ξ^- momentum and displacement	< 0.0447 rad
Angle between Λ momentum and displacement	< 0.14 rad
Angle between Λ_b^0 momentum and displacement	< 0.0447 rad
PV impact parameter for pion from Ξ^- decay	> 0.4 s.d.
PV impact parameter for kaon	> 0.4 s.d.

In less than 5% of the events, multiple Λ_b^0 candidates in the same channel are found. The rate is consistent in both channels and all candidates are used in the analysis. Selecting a single candidate has a negligible effect on the results.

4 Invariant mass distributions

The measured mass distribution of the $\psi(2S)\Lambda$ candidates is shown in Fig. 1 (left) together with the results of an unbinned maximum likelihood fit. The signal is modeled with a Student's t -distribution [35] with all parameters (mean, σ , n) free. The combinatorial background is described by an exponential function with a free slope parameter and normalization. The fitted mass of 5619.3 ± 0.3 MeV is in agreement with the world-average Λ_b^0 mass of 5619.60 ± 0.17 MeV [33], and the mass resolution of 8.90 ± 0.40 MeV is slightly larger than, yet in agreement with, its value of 8.52 MeV found in simulation. The measured yield is $N(\Lambda_b^0 \rightarrow \psi(2S)\Lambda) = 1744 \pm 63$. The χ^2 between the binned distribution and the fit function is 76.6 for 94 degrees of freedom, demonstrating the good quality of the fit.

The measured invariant mass distribution of the selected $J/\psi \Xi^- K^+$ candidates is shown in Fig. 1 (lower). A narrow peak at the Λ_b^0 mass is seen on top of a smooth background.




Fig. 1 Measured $\psi(2S)\Lambda$ (upper) and $J/\psi \Xi^- K^+$ (lower) invariant mass distributions and overlaid fit results

The Λ_b^0 signal is modeled with a Student's t -distribution with mean and σ floating, but the n parameter fixed to the value found by fitting the simulated distribution, because of the limited signal yield of $N(\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+) = 46 \pm 11$. The background is fitted with an exponential function. The Λ_b^0 mass returned by the fit (5625.9 ± 3.2 MeV) is within 2 s.d. of the world-average value [33]. The width of the signal peak (σ) is found to be 10.4 ± 3.3 MeV, consistent within 1.2 s.d. with the value found in simulation, 6.6 ± 0.2 MeV. The fit quality is good, as demonstrated by the $\chi^2/\text{dof} = 30.1/45$ for the binned distribution.

The signal significance is evaluated using the likelihood ratio technique by applying the background-only and signal-plus-background hypotheses. In these two fits, a Gaussian constraint is applied on the background shape parameter to the one obtained from a fit to the wrong-sign data. Similarly, a Gaussian constraint is applied to the signal shape parameter n (from simulation) and the resolution $\sigma = \sigma_{\text{MC}}$ (8.90/8.52). The correction factor is extracted from the normalization channel and accounts for the difference in the widths of the peak between the measured and simulated event samples. The

mean value of the peak is also Gaussian-constrained with a central value and uncertainty equal to the known Λ_b^0 mass and its uncertainty [33], respectively. The fit with the signal-plus-background model with these constraints returns a signal yield of 36 ± 8 and is presented in Appendix A. Since the conditions of Wilks' theorem [36] are satisfied, the asymptotic formulae of Ref. [37] (Eqs. (12) and (52)) are used to determine the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ signal significance, which is found to be 5.8 standard deviations. To evaluate the effect of the choice of the model for fitting the signal significance, several alternative models of signal and background were tested, including double-Gaussian or Johnson [38] functions for the signal and a second-degree polynomial or a modified threshold function for the background. An alternative without a constraint on the background shape was also tested. The significance obtained with the alternative models varies in the range from 5.3 to 5.9 standard deviations. This allows us to claim the first observation of the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay.

The sensitivity of this analysis to potential pentaquark signals in the intermediate invariant mass distributions of the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay is limited by the low signal yield. The background-subtracted two-body invariant mass distributions, obtained with the $s\mathcal{P}$ Plot technique [39], are shown in Fig. 2. The distributions do not show any clear peaks and agree, within uncertainties, with the predictions from the phase space simulation. The distributions are also consistent with the results of extracting the yields by fitting the Λ_b^0 signal in each of the five intermediate invariant mass bins.

For the measurement of \mathcal{R} (Eq. (1)), more stringent requirements are used, as explained at the end of Sect. 3, and the measured signal yields decrease to 1179 ± 47 and 23 ± 7 for the $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ and $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ channels, respectively, using unconstrained fits as for Fig. 1. These are the baseline results referred to in Sect. 6. The corresponding mass distributions and fits are presented in Fig. 3.

5 Efficiencies

Efficiencies for the signal and normalization channels are calculated using simulated event samples. The total efficiency is calculated by factorizing into two components: detector acceptance and a combined trigger, reconstruction, and selection efficiency.

As only the ratio of the total efficiencies is needed to measure \mathcal{R} , the systematic uncertainties associated with the muon and track reconstruction are reduced. The obtained efficiency ratio is $\epsilon_{\psi(2S)\Lambda} / \epsilon_{J/\psi \Xi^- K^+} = 5.06 \pm 0.29$, where the uncertainty reflects the limited size of the simulated samples. Efficiencies for different years of data-taking are combined with weights corresponding to the integrated luminosity collected




Fig. 2 Intermediate invariant mass distributions of the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay. The filled circles and empty squares show the measured background-subtracted distributions and the results from the simulation with a phase-space model, respectively




Fig. 3 Measured $\psi(2S)\Lambda$ (upper) and $J/\psi \Xi^- K^+$ (lower) invariant mass distributions and corresponding fits used for the measurement of \mathcal{R}

in each year. The efficiency for the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ channel is significantly lower than that for the $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ channel for several reasons including the low energy release in the $\Xi^- \rightarrow \Lambda \pi^-$ decay, resulting in a low-momentum pion track.

6 Systematic uncertainties

Many systematic uncertainties, related to the muon reconstruction and identification as well as to the trigger efficiency, partially cancel in the measured ratios. Since the signal and normalization channels have the same number of tracks in the final state, most uncertainties related to track reconstruction also cancel in the measured ratio \mathcal{R} . However, the p_T spectrum of kaons from the $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ decay is observed to differ from that of the highest- p_T pion in the $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ channel used for normalization. Despite the signal and normalization channels having the same number of final-state tracks, an uncertainty of 2.3% [40] is included, which reflects the difference in tracking efficiency between the measured and simulated event samples. The MC event samples are validated using the normalization channel by comparing the measured distributions of variables used in the event selection, after background subtraction, to those found in simulation; no significant discrepancies are found in most of the distributions. A small discrepancy was observed in the $p_T(\Lambda_b^0)$ distribution, and the MC event samples for both channels were reweighted using $p_T(\Lambda_b^0)$ -dependent weights so that the $p_T(\Lambda_b^0)$ distribution in the weighted simulation sample matches the background-subtracted distribution measured in the $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ channel. The efficiency ratio evaluated using these weighted MC samples is found to be $\epsilon_{\psi(2S)\Lambda} / \epsilon_{J/\psi \Xi^- K^+} = 4.82 \pm 0.39$, which is lower by 4.7%, yet still in agreement with the value reported in the previous section. An uncertainty of 4.7% is assigned to account for potential mismodeling of the $p_T(\Lambda_b^0)$ spectrum.

The systematic uncertainty related to the choice of the signal model is evaluated by testing three different models. For the normalization channel $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ the signal shape parameters are floating, while for the signal channel $\Lambda_b^0 \rightarrow J/\psi \Xi^- K^+$ the mass resolution parameters are fixed to those found in simulation, after correcting the width of the peak for the ratio between the two resolutions in the

measured and simulated event samples evaluated in the normalization channel. The tested models simultaneously vary the signal and normalization channels and use a Student's t -distribution, a double-Gaussian, and a Johnson function [38] to model the signal. The largest deviation in the ratio of the Λ_b^0 signal yields from the baseline value is taken as the systematic uncertainty.

The systematic uncertainty related to the choice of the background model is estimated in a similar way, with three alternative models: a second-degree polynomial, a threshold function [41, 42] multiplied by an exponential, and a threshold function multiplied by a first-degree polynomial.

By requiring the $J/\psi\pi^+\pi^-$ invariant mass to be near the $\psi(2S)$ mass, we aim to select $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ decays. Nevertheless, other $\Lambda_b^0 \rightarrow J/\psi\pi^+\pi^-\Lambda$ decays, either from different intermediate resonances or from four-body nonresonant decays may contribute. To estimate this contribution, we use the $s\mathcal{P}$ lot technique to subtract the background under the Λ_b^0 peak and plot the $J/\psi\pi^+\pi^-$ mass in an expanded mass region. The $J/\psi\pi^+\pi^-$ mass is fitted with a signal component for the $\psi(2S)$ and a background component for everything else. The integral of the background over the range used to select $\psi(2S)$ events yields 30 events, which is 2.5% of the total yield (1179 events). This value is used as the systematic uncertainty related to non- $\psi(2S)$ contributions in the normalization channel.

The uncertainty in the efficiency ratio due to the limited size of the simulated samples, calculated to be 5.6% in Sect. 5, is considered as a systematic uncertainty.

In order to assess the reliability of the efficiency evaluation from the simulated samples, the selection criteria on muon and J/ψ p_T , dimuon vertex probability, track impact parameter, and p_T of the soft pion from Ξ^- decay are tightened, one at a time, until the signal efficiency decreases by 10 or 20% with respect to that obtained with the selection used for the \mathcal{R} measurement. The analysis is repeated each time, and the value of \mathcal{R} is re-calculated and compared to the baseline \mathcal{R} value. The differences (d) between the two values and their uncertainties (δd), which also account for the correlation between the two values, are evaluated. The largest value of $\sqrt{d^2 - (\delta d)^2}$ among the different variations of the selection criteria is found to be 14.3% and is used as the systematic uncertainty in the efficiency ratio.

Table 2 summarizes the previously discussed systematic uncertainties in the ratio \mathcal{R} . The total uncertainty is calculated as the sum in quadrature of the individual sources.

7 Branching fraction ratio measurement

The branching fraction of the newly observed $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$ decay, with respect to the $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ one,

Table 2 The relative systematic uncertainties in the measurement of \mathcal{R}

Source	Uncertainty (%)
Tracking efficiency	2.3
$p_T(\Lambda_b^0)$ spectrum	4.7
Signal model	3.9
Background model	6.7
Non- $\psi(2S)$ contribution	2.5
Limited size of MC samples	5.6
Selection efficiency	14.3
Total	18.2

is measured using Eq. (1) to be

$$\begin{aligned}\mathcal{R} &\equiv \frac{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+)}{\mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S)\Lambda)} \\ &= [3.38 \pm 1.02 \text{ (stat)} \pm 0.61 \text{ (syst)} \pm 0.03 \text{ (}\mathcal{B}\text{)}]\%,\end{aligned}$$

where the last uncertainty is related to the uncertainties in the branching fractions $\mathcal{B}(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 34.68 \pm 0.30\%$ and $\mathcal{B}(\Xi^- \rightarrow \Lambda\pi^-) = 99.887 \pm 0.035\%$ [33].

8 Summary

The $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$ decay is observed with a significance exceeding 5 standard deviations using $\sqrt{s} = 13$ TeV proton–proton collision data corresponding to an integrated luminosity of 140 fb^{-1} collected by the CMS experiment. The branching fraction is measured with respect to the $\Lambda_b^0 \rightarrow \psi(2S)\Lambda$ decay to be $\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+)/\mathcal{B}(\Lambda_b^0 \rightarrow \psi(2S)\Lambda) = [3.38 \pm 1.02 \text{ (stat)} \pm 0.61 \text{ (syst)} \pm 0.03 \text{ (}\mathcal{B}\text{)}]\%$. The distributions of intermediate invariant masses $m(J/\psi\Xi^-)$, $m(J/\psi K^+)$, and $m(\Xi^- K^+)$ from the $\Lambda_b^0 \rightarrow J/\psi\Xi^-K^+$ decay are also presented. This is the first discovered multi-body decay containing the $J/\psi\Xi^-$ system, which opens the possibility to search for doubly-strange hidden-charm pentaquarks when more data are collected. The new results are important for understanding the strong interaction processes in hadronic decays of beauty baryons and the possible formation of exotic multi-quark states.

Acknowledgements We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Arme-

nia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA). Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22r1-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science – EOS" – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), under Germany's Excellence Strategy – EXC 2121 "Quantum Universe" – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing, funded by the EU NexGeneration program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF "a way of making Europe", and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B37G660013 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

Data Availability Statement This manuscript has associated data in a data repository. [Author's comment: Some data used for this analysis may be made available. Almost certainly not all data will be released near term.]

Code Availability Statement Code/software cannot be made available for reasons disclosed in the code availability statement. [Author's com-

ment: Some software (the core software) is available. Not all code used is available.]

Declarations

Conflict of interest The authors declare that they have no Conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³.

A Invariant mass distribution fit with constraints

The fit of the measured $J/\psi \Xi^- K^+$ invariant mass distribution with constraints on the background shape parameter, signal shape parameter, resolution, and the mean value of the peak is presented in Fig. 4. The fit quality is good, as demonstrated by $\chi^2/\text{dof} = 35.6/44$ for the binned distribution.




Fig. 4 Measured $J/\psi \Xi^- K^+$ invariant mass distribution and overlaid constrained fit result


References

- LHCb Collaboration, Observation of $J/\psi p$ resonances consistent with pentaquark states in $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays. *Phys. Rev. Lett.* **115**, 072001 (2015). <https://doi.org/10.1103/PhysRevLett.115.072001>. arXiv:1507.03414
- N. Brambilla et al., The XYZ states: experimental and theoretical status and perspectives. *Phys. Rep.* **873**, 1 (2020). <https://doi.org/10.1016/j.physrep.2020.05.001>. arXiv:1907.07583
- H.-X. Chen et al., An updated review of the new hadron states. *Rep. Prog. Phys.* **86**, 026201 (2023). <https://doi.org/10.1088/1361-6633/aca3b6>. arXiv:2204.02649
- L. Maiani, A.D. Polosa, V. Riquer, The new pentaquarks in the diquark model. *Phys. Lett. B* **749**, 289 (2015). <https://doi.org/10.1016/j.physletb.2015.08.008>. arXiv:1507.04980
- E. Santopinto, A. Giachino, Compact pentaquark structures. *Phys. Rev. D* **96**, 014014 (2017). <https://doi.org/10.1103/PhysRevD.96.014014>. arXiv:1604.03769
- L. Roca, J. Nieves, E. Oset, LHCb pentaquark as a $\bar{D}^* \Sigma_c - \bar{D}^* \Sigma_c^*$ molecular state. *Phys. Rev. D* **92**, 094003 (2015). <https://doi.org/10.1103/PhysRevD.92.094003>. arXiv:1507.04249
- M.I. Eides, V.Y. Petrov, M.V. Polyakov, Pentaquarks with hidden charm as hadroquarkonia. *Eur. Phys. J. C* **78**, 36 (2018). <https://doi.org/10.1140/epjc/s10052-018-5530-9>. arXiv:1709.09523
- M.-L. Du et al., Interpretation of the LHCb P_c states as hadronic molecules and hints of a narrow $P_c(4380)$. *Phys. Rev. Lett.* **124**, 072001 (2020). <https://doi.org/10.1103/PhysRevLett.124.072001>. arXiv:1910.11846
- S.X. Nakamura, $P_c(4312)^+$, $P_c(4380)^+$, and $P_c(4457)^+$ as double triangle cusps. *Phys. Rev. D* **103**, 111503 (2021). <https://doi.org/10.1103/PhysRevD.103.L111503>. arXiv:2103.06817
- LHCb Collaboration, Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$. *Phys. Rev. Lett.* **122**, 222001 (2019). <https://doi.org/10.1103/PhysRevLett.122.222001>. arXiv:1904.03947
- LHCb Collaboration, Evidence of a $J/\psi \Lambda$ structure and observation of excited Ξ^- states in the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay. *Sci. Bull.* **66**, 1278 (2021). <https://doi.org/10.1016/j.scib.2021.02.030>. arXiv:2012.10380
- LHCb Collaboration, Evidence for a new structure in the $J/\psi p$ and $J/\psi \bar{p}$ systems in $B_s^0 \rightarrow J/\psi p \bar{p}$ decays. *Phys. Rev. Lett.* **128**, 062001 (2022). <https://doi.org/10.1103/PhysRevLett.128.062001>. arXiv:2108.04720
- LHCb Collaboration, Observation of a $J/\psi \Lambda$ resonance consistent with a strange pentaquark candidate in $B^- \rightarrow J/\psi \Lambda \bar{p}$ decays. *Phys. Rev. Lett.* **131**, 031901 (2023). <https://doi.org/10.1103/PhysRevLett.131.031901>. arXiv:2210.10346
- K. Azizi, Y. Sarac, H. Sundu, Investigation of hidden-charm double strange pentaquark candidate P_{cs} via its mass and strong decays. *Eur. Phys. J. C* **82**, 543 (2022). <https://doi.org/10.1140/epjc/s10052-022-10495-7>. arXiv:2112.15543
- F.-L. Wang, X.-D. Yang, R. Chen, X. Liu, Hidden-charm pentaquarks with triple strangeness due to the $\Omega_c^{(*)} \bar{D}_s^{(*)}$ interactions. *Phys. Rev. D* **103**, 054025 (2021). <https://doi.org/10.1103/PhysRevD.103.054025>. arXiv:2101.11200
- CMS Collaboration, Precision luminosity measurement in proton–proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS. *Eur. Phys. J. C* **81**, 800 (2021). <https://doi.org/10.1140/epjc/s10052-021-09538-2>. arXiv:2104.01927
- CMS Collaboration, CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV, CMS Physics Analysis Summary CMS-PAS-LUM-17-004 (2018). <https://cds.cern.ch/record/2621960>
- CMS Collaboration, CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV, CMS Physics Analysis Summary CMS-PAS-LUM-18-002 (2019). <https://cds.cern.ch/record/2676164>
- HEPData record for this analysis (2024). <https://doi.org/10.17182/hepdata.145642>
- CMS Collaboration, The CMS experiment at the CERN LHC. *JINST* **3**, S08004 (2008). <https://doi.org/10.1088/1748-0221/3/08/S08004>
- C.M.S. Collaboration, Description and performance of track and primary-vertex reconstruction with the CMS tracker. *JINST* **9**, P10009 (2014). <https://doi.org/10.1088/1748-0221/9/10/P10009>. arXiv:1405.6569
- CMS Tracker Collaboration, The CMS Phase-1 pixel detector upgrade. *JINST* **16**, P02027 (2021). <https://doi.org/10.1088/1748-0221/16/02/P02027>. arXiv:2012.14304
- CMS Collaboration, Track impact parameter resolution for the full pseudo-rapidity coverage in the 2017 dataset with the CMS Phase-1 pixel detector. CMS Detector Performance Note CMS-DP-2020-049, 2020. <https://cds.cern.ch/record/2743740>
- CMS Collaboration, The CMS trigger system. *JINST* **12**, P01020 (2017). <https://doi.org/10.1088/1748-0221/12/01/P01020>. arXiv:1609.02366
- CMS Collaboration, Performance of the CMS Level-1 trigger in proton–proton collisions at $\sqrt{s} = 13$ TeV. *JINST* **15**, P10017 (2020). <https://doi.org/10.1088/1748-0221/15/10/P10017>. arXiv:2006.10165
- T. Sjöstrand et al., An introduction to PYTHIA 8.2. *Comput. Phys. Commun.* **191**, 159 (2015). <https://doi.org/10.1016/j.cpc.2015.01.024>. arXiv:1410.3012
- CMS Collaboration, Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements. *Eur. Phys. J. C* **80**, 4 (2020). <https://doi.org/10.1140/epjc/s10052-019-7499-4>. arXiv:1903.12179
- D.J. Lange, The EvtGen particle decay simulation package. *Nucl. Instrum. Methods A* **462**, 152 (2001). [https://doi.org/10.1016/S0168-9002\(01\)00089-4](https://doi.org/10.1016/S0168-9002(01)00089-4)
- E. Barberio, Z. Was, PHOTOS—a universal Monte Carlo for QED radiative corrections: version 2.0. *Comput. Phys. Commun.* **79**, 291 (1994). [https://doi.org/10.1016/0010-4655\(94\)90074-4](https://doi.org/10.1016/0010-4655(94)90074-4)
- GEANT4 Collaboration, Geant4—a simulation toolkit. *Nucl. Instrum. Methods A* **506**, 250 (2003). [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8)
- CMS Collaboration, Performance of the CMS muon detector and muon reconstruction with proton–proton collisions at $\sqrt{s} = 13$ TeV. *JINST* **13**, P06015 (2018). <https://doi.org/10.1088/1748-0221/13/06/P06015>. arXiv:1804.04528
- C.M.S. Collaboration, CMS tracking performance results from early LHC operation. *Eur. Phys. J. C* **70**, 1165 (2010). <https://doi.org/10.1140/epjc/s10052-010-1491-3>. arXiv:1007.1988
- Particle Data Group Collaboration, Review of particle physics. *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022). <https://doi.org/10.1093/ptep/ptac097>
- G. Punzi, Sensitivity of searches for new signals and its optimization, in *Proceedings of PHYSTAT 2003, Statistical Problems in Particle Physics, Astrophysics and Cosmology*, eConf C030908 (2003), p. MODT002. arXiv:physics/0308063
- S. Jackman, *Bayesian Analysis for the Social Sciences* (Wiley, New Jersey, 2009). <https://doi.org/10.1002/9780470686621>
- S.S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses. *Ann. Math. Stat.* **9**, 60 (1938). <https://doi.org/10.1214/aoms/1177732360>
- G. Cowan, K. Cranmer, E. Gross, O. Vitells, Asymptotic formulae for likelihood-based tests of new physics. *Eur. Phys. J. C* **71**, 1554 (2011) (Erratum: 10.1140/epjc/s10052-011-1554-0). <https://doi.org/10.1140/epjc/s10052-013-2501-z>. arXiv:1007.1727




















38. N.L. Johnson, Systems of frequency curves generated by methods of translation. *Biometrika* **36**, 149 (1949). <https://doi.org/10.2307/2332539>
39. M. Pivk, F.R. Le Diberder, \mathcal{S} Plot: a statistical tool to unfold data distributions. *Nucl. Instrum. Methods A* **555**, 356 (2005). <https://doi.org/10.1016/j.nima.2005.08.106>. [arXiv:physics/0402083](https://arxiv.org/abs/physics/0402083)
40. CMS Collaboration, Tracking performances for charged pions with Run2 legacy data, CMS Detector Performance Note CMS-DP-2022-012, 2022. <https://cds.cern.ch/record/2810814>
41. CMS Collaboration, Observation of a new excited beauty strange baryon decaying to $\Xi_b^- \pi^+ \pi^-$. *Phys. Rev. Lett.* **126**, 252003 (2021). <https://doi.org/10.1103/PhysRevLett.126.252003>. [arXiv:2102.04524](https://arxiv.org/abs/2102.04524)
42. CMS Collaboration, Study of excited Λ_b^0 states decaying to $\Lambda_b^0 \pi^+ \pi^-$ in proton-proton collisions at $\sqrt{s} = 13$ TeV. *Phys. Lett. B* **803**, 135345 (2020). <https://doi.org/10.1016/j.physletb.2020.135345>. [arXiv:2001.06533](https://arxiv.org/abs/2001.06533)

CMS Collaboration




Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan ¹















Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J. W. Andrejkovic , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicovic ,
A. Escalante Del Valle , P. S. Hussain , M. Jeitler ², N. Krammer , D. Liko , I. Mikulec , J. Schieck ²,
R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger , C.-E. Wulz ²

















Universiteit Antwerpen, Antwerpen, Belgium

M. R. Darwish ³, T. Janssen , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

E. S. Bols , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , H. El Faham , S. Lowette , I. Makarenko ,
D. Müller , A. R. Sahasransu , S. Tavernier , M. Tytgat ⁴, S. Van Putte , D. Vannerom 















Université Libre de Bruxelles, Bruxelles, Belgium

B. Clerbaux , G. De Lentdecker , L. Favart , D. Hohov , J. Jaramillo , A. Khalilzadeh , K. Lee ,
M. Mahdavihorrani , A. Malara , S. Paredes , L. Pétré , N. Postiau , L. Thomas , M. Vanden Bemden ,
C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , Y. Hong , J. Knolle , L. Lambrecht , G. Mestdach , C. Rendón , A. Samalan ,
K. Skovpen , N. Van Den Bossche , L. Wezenbeek 




















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , G. Bruno , C. Caputo , C. Delaere , I. S. Donertas , A. Giammanco , K. Jaffel , Sa. Jain ,
V. Lemaitre , J. Lidrych , P. Mastrapasqua , K. Mondal , T. T. Tran , S. Wertz 









Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G. A. Alves , E. Coelho , C. Hensel , T. Menezes De Oliveira , A. Moraes , P. Rebello Teles , M. Soeiro 








Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W. L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho ,
J. Chinellato ⁵, E. M. Da Costa , G. G. Da Silveira ⁶, D. De Jesus Damiao , S. Fonseca De Souza , J. Martins ⁷,
C. Mora Herrera , K. Mota Amarilo , L. Mundim , H. Nogima , A. Santoro , A. Sznajder , M. Thiel ,
A. Vilela Pereira 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil







































C. A. Bernardes ⁶, L. Calligaris , T. R. Fernandez Perez Tomei , E. M. Gregores , P. G. Mercadante ,
S. F. Novaes , B. Orzari , Sandra S. Padula 










































































































Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 
























University of Sofia, Sofia, Bulgaria

A. Dimitrov , T. Ivanov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 


















Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, ChileS. Keshri , S. Thakur **Beihang University, Beijing, China**T. Cheng , Q. Guo, T. Javaid , M. Mittal , L. Yuan **Department of Physics, Tsinghua University, Beijing, China**G. Bauer ^{8,9}, Z. Hu , J. Liu, K. Yi ^{8,10}**Institute of High Energy Physics, Beijing, China**G. M. Chen ¹¹, H. S. Chen ¹¹, M. Chen ¹¹, F. Iemmi , C. H. Jiang, A. Kapoor ¹², H. Liao , Z.-A. Liu ¹³, F. Monti , M. A. Shahzad ¹¹, R. Sharma ¹⁴, J. N. Song ¹³, J. Tao , C. Wang ¹¹, J. Wang , Z. Wang ¹¹, H. Zhang **State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China**A. Agapitos , Y. Ban , A. Levin , C. Li , Q. Li , Y. Mao, S. J. Qian , X. Sun , D. Wang , H. Yang, L. Zhang , C. Zhou **Sun Yat-Sen University, Guangzhou, China**Z. You **University of Science and Technology of China, Hefei, China**N. Lu **Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE)-Fudan University, Shanghai, China**X. Gao ¹⁵, D. Leggat, H. Okawa , Y. Zhang **Zhejiang University, Hangzhou Zhejiang, China**Z. Lin , C. Lu , M. Xiao **Universidad de Los Andes, Bogota, Colombia**C. Avila , D. A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J. A. Reyes Vega**Universidad de Antioquia, Medellin, Colombia**J. Mejia Guisao , F. Ramirez , M. Rodriguez , J. D. Ruiz Alvarez **University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac **University of Split, Faculty of Science, Split, Croatia**M. Kovac , T. Sculac **Institute Rudjer Boskovic, Zagreb, Croatia**P. Bargassa , V. Brigljevic , B. K. Chitroda , D. Ferencek , S. Mishra , A. Starodumov ¹⁶, T. Susa **University of Cyprus, Nicosia, Cyprus**A. Attikis , K. Christoforou , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P. A. Razis , H. Rykaczewski, H. Saka , A. Stepennov **Charles University, Prague, Czech Republic**M. Finger , M. Finger Jr. , A. Kveton **Escuela Politecnica Nacional, Quito, Ecuador**E. Ayala **Universidad San Francisco de Quito, Quito, Ecuador**E. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**H. Abdalla ¹⁷, Y. Assran ^{18,19}

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, EgyptM. Abdullah Al-Mashad , M. A. Mahmoud **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**R. K. Dewanjee ²⁰, K. Ehataht , M. Kadastik , T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken **Department of Physics, University of Helsinki, Helsinki, Finland**H. Kirschenmann , K. Osterberg , M. Voutilainen **Helsinki Institute of Physics, Helsinki, Finland**S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , K. T. S. Kallonen , M. S. Kim , R. Kinnunen, T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti, L. Martikainen , M. Myllymäki , M. M. Rantanen , H. Siikonen , E. Tuominen , J. Tuominiemi **Lappeenranta-Lahti University of Technology, Lappeenranta, Finland**P. Luukka , H. Petrow , T. Tuuva[†]**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J. L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , V. Lohezic , J. Malcles , J. Rander, A. Rosowsky , M. Ö. Sahin , A. Savoy-Navarro ²¹, P. Simkina , M. Titov , M. Tornago **Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**C. Baldenegro Barrera , F. Beaudette , A. Buchot Perraguin , P. Busson , A. Cappati , C. Charlot , F. Damas , O. Davignon , A. De Wit , G. Falmagne , B. A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , J. Motta , M. Nguyen , C. Ochando , L. Portales , R. Salerno , U. Sarkar , J. B. Sauvan , Y. Sirois , A. Tarabini , E. Vernazza , A. Zabi , A. Zghiche **Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France**J.-L. Agram ²², J. Andrea , D. Appar , D. Bloch , J.-M. Brom , E. C. Chabert , C. Collard , S. Falke , U. Goerlach , C. Grimault, R. Haeberle , A.-C. Le Bihan , M. A. Sessini , P. Van Hove **Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , J. Choi , D. Contardo , P. Depasse , C. Dozen ²³, H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg, G. Grenier , B. Ille , I. B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao **Georgian Technical University, Tbilisi, Georgia**G. Adamov, I. Lomidze , Z. Tsamalaidze ¹⁶**RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**V. Botta , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , N. Röwert , M. Teroerde **RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M. Y. Lee , L. Mastrolorenzo, M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , A. Novak , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , A. Sharma , A. Stein , F. Torres Da Silva De Araujo ²⁴, L. Vigilante, S. Wiedenbeck , S. Zaleski**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**C. Dziwok , G. Flüge , W. Haj Ahmad ²⁵, T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 


















Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , S. Bhattacharya , F. Blekman ²⁶, K. Borras ²⁷, D. Brunner , A. Campbell , A. Cardini , C. Cheng, F. Colombina , S. Consuegra Rodríguez , G. Correia Silva , M. De Silva , G. Eckerlin, D. Eckstein , L. I. Estevez Banos , O. Filatov , E. Gallo ²⁶, A. Geiser , A. Giraldi , G. Greau, V. Guglielmi , M. Guthoff , A. Hinzmann , A. Jafari ²⁸, L. Jeppe , N. Z. Jomhari , B. Kaech , M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁹, W. Lohmann ³⁰, R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , J. Metwally, A. B. Meyer , G. Milella , A. Mussgiller , A. Nürnberg , Y. Otari, D. Pérez Adán , E. Ranken , A. Raspereza , B. Ribeiro Lopes , J. Rübenach, A. Saggio , M. Scham ^{31,27}, S. Schnake ²⁷, P. Schütze , C. Schwanenberger ²⁶, D. Selivanova , M. Shchedrolosiev , R. E. Sosa Ricardo , L. P. Sreelatha Pramod , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²⁷, C. Wissing , S. Wuchterl , Y. Yang , A. Zimmermann Castro Santos

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , M. Bonanomi , P. Connor , M. Eich, K. El Morabit , Y. Fischer , A. Fröhlich, C. Garbers , E. Garutti , A. Grohsjean , M. Hajheidari, J. Haller , H. R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcar , T. Kramer , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K. J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf







Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ³², R. Hofsaess , M. Horzela , U. Husemann , M. Klute , R. Koppenhöfer , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H. J. Simonis , N. Trevisani , R. Ulrich , J. van der Linden , R. F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , S. Wunsch, X. Zuo





Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, P. Assiouras , G. Daskalakis , A. Kyriakis, A. Papadopoulos ³², A. Stakia 





National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, A. Panagiotou, I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos



National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsiolitis, A. Zacharopoulou










University of Ioánnina, Ioannina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios , C. Kamtsikis, P. Katsoulis, P. Kokkas , P. G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

M. Bartók ³³, C. Hajdu , D. Horvath ^{34,35}, F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary




































































































M. Csanád , K. Farkas , M. M. A. Gadallah ³⁶, Á. Kadlecik , P. Major , K. Mandal , G. Pásztor , A. J. Rádl ³⁷, G. I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary



















P. Raics, B. Ujvari ³⁸, G. Zilizi 

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Karancsi ³³, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, HungaryT. Csorgo ³⁷, F. Nemes ³⁷, T. Novak **Panjab University, Chandigarh, India**J. Babbar , S. Bansal , S. B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra ³⁹, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , M. Meena , K. Sandeep , T. Sheokand, J. B. Singh , A. Singla **University of Delhi, Delhi, India**A. Ahmed , A. Bhardwaj , A. Chhetri , B. C. Choudhary , A. Kumar , M. Naimuddin , K. Ranjan , S. Saumya **Saha Institute of Nuclear Physics, HBNI, Kolkata, India**S. Acharya ⁴⁰, S. Baradia , S. Barman ⁴¹, S. Bhattacharya , D. Bhowmik, S. Dutta , S. Dutta, B. Gomber ⁴⁰, P. Palit , G. Saha , B. Sahu ⁴⁰, S. Sarkar**Indian Institute of Technology Madras, Madras, India**M. M. Ameen , P. K. Behera , S. C. Behera , S. Chatterjee , P. Jana , P. Kalbhor , J. R. Komaragiri ⁴², D. Kumar ⁴², L. Panwar ⁴², R. Pradhan , P. R. Pujahari , N. R. Saha , A. Sharma , A. K. Sikdar , S. Verma **Tata Institute of Fundamental Research-A, Mumbai, India**T. Aziz, I. Das , S. Dugad, M. Kumar , G. B. Mohanty , P. Suryadevara**Tata Institute of Fundamental Research-B, Mumbai, India**A. Bala , S. Banerjee , R. M. Chatterjee, M. Guchait , Sh. Jain , S. Karmakar , S. Kumar , G. Majumder , K. Mazumdar , S. Mukherjee , S. Parolia , A. Thachayath **National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**S. Bahinipati ⁴³, A. K. Das, C. Kar , D. Maity ⁴⁴, P. Mal , T. Mishra , V. K. Muraleedharan Nair Bindhu ⁴⁴, K. Naskar ⁴⁴, A. Nayak ⁴⁴, P. Sadangi, P. Saha , S. K. Swain , S. Varghese ⁴⁴, D. Vats ⁴⁴**Indian Institute of Science Education and Research (IISER), Pune, India**A. Alpana , S. Dube , B. Kansal , A. Laha , A. Rastogi , S. Sharma **Isfahan University of Technology, Isfahan, Iran**H. Bakhshiansohi ⁴⁵, E. Khazaie ⁴⁶, M. Zeinali ⁴⁷**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**S. Chenarani ⁴⁸, S. M. Etesami , M. Khakzad , M. Mohammadi Najafabadi **University College Dublin, Dublin, Ireland**M. Grunewald **INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy**M. Abbrescia ^{a,b}, R. Aly ^{a,c,49}, A. Colaleo ^{a,b}, D. Creanza ^{a,c}, B. D'Anzi ^{a,b}, N. De Filippis ^{a,c}, M. De Palma ^{a,b}, A. Di Florio ^{a,c}, W. Elmetenawee ^{a,b,49}, L. Fiore ^a, G. Iaselli ^{a,c}, G. Maggi ^{a,c}, M. Maggi ^a, I. Margjeka ^{a,b}, V. Mastrapasqua ^{a,b}, S. My ^{a,b}, S. Nuzzo ^{a,b}, A. Pellicchia ^{a,b}, A. Pompili ^{a,b}, G. Pugliese ^{a,c}, R. Radogna ^a, G. Ramirez-Sanchez ^{a,c}, D. Ramos ^a, A. Ranieri ^a, L. Silvestris ^a, F. M. Simone ^{a,b}, Ü. Sözbilir ^a, A. Stamerra ^a, R. Venditti ^a, P. Verwilligen ^a, A. Zaza ^{a,b}**INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy**G. Abbiendi ^a, C. Battilana ^{a,b}, D. Bonacorsi ^{a,b}, L. Borgonovi ^a, R. Campanini ^{a,b}, P. Capiluppi ^{a,b}, F. R. Cavallo ^a, M. Cuffiani ^{a,b}, G. M. Dallavalle ^a, T. Diotallevi ^{a,b}, F. Fabbri ^a, A. Fanfani ^{a,b}, D. Fasanella ^{a,b}, P. Giacomelli ^a, L. Giommi ^{a,b}, C. Grandi ^a, L. Guiducci ^{a,b}, S. Lo Meo ^{a,50}, L. Lunerti ^{a,b}, S. Marcellini ^a, G. Masetti ^a, F. L. Navarria ^{a,b}, A. Perrotta ^a, F. Primavera ^{a,b}, A. M. Rossi ^{a,b}, T. Rovelli ^{a,b}, G. P. Siroli ^{a,b}**INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy**S. Costa ^{a,b,51}, A. Di Mattia ^a, R. Potenza ^{a,b}, A. Tricomi ^{a,b,51}, C. Tuve ^{a,b}




INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

G. Barbagli , G. Bardelli , B. Camaiani , A. Cassese , R. Ceccarelli , V. Ciulli , C. Civinini , R. D'Alessandro , E. Focardi , T. Kello , G. Latino , P. Lenzi , M. Lizzo , M. Meschini , S. Paoletti , A. Papanastassiou , G. Sguazzoni , L. Viliani 

























INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola , D. Piccolo 












INFN Sezione di Genova^a, Università di Genova^b, Genoa, Italy

P. Chatagnon , F. Ferro , E. Robutti , S. Tosi 































INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milan, Italy

A. Benaglia , G. Boldrini , F. Brivio , F. Cetorelli , F. De Guio , M. E. Dinardo , P. Dini , S. Gennai , R. Gerosa , A. Ghezzi , P. Govoni , L. Guzzi , M. T. Lucchini , M. Malberti , S. Malvezzi , A. Massironi , D. Menasce , L. Moroni , M. Paganoni , D. Pedrini , B. S. Pinolini , S. Ragazzi , T. Tabarelli de Fatis , D. Zuolo 












INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Naples, Italy

S. Buontempo , A. Cagnotta , F. Carnevali , N. Cavallo , A. De Iorio , F. Fabozzi , A. O. M. Iorio , L. Lista , P. Paolucci , B. Rossi , C. Sciacca 

















INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino , P. Azzi , N. Bacchetta , D. Bisello , P. Bortignon , A. Bragagnolo , R. Carlin , T. Dorigo , F. Gasparinis , U. Gasparini , A. Gozzelino , G. Grosso , L. Layer , E. Lusiani , M. Margoni , A. T. Meneguzzo , M. Migliorini , J. Pazzini , P. Ronchese , R. Rossin , F. Simonetto , G. Strong , M. Tosi , A. Triossi , S. Ventura , H. Yarar , M. Zanetti , P. Zotto , A. Zucchetta , G. Zumerle 































INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

S. Abu Zeid , C. Aimè , A. Braghieri , S. Calzaferri , D. Fiorina , P. Montagna , V. Re , C. Riccardi , P. Salvini , I. Vai , P. Vitulo 




















INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal , P. Asenov , G. M. Bilei , D. Ciangottini , L. Fanò , M. Magherini , G. Mantovani , V. Mariani , M. Menichelli , F. Moscatelli , A. Piccinelli , M. Presilla , A. Rossi , A. Santocchia , D. Spiga , T. Tedeschi 







































INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

P. Azzurri , G. Bagliesi , R. Bhattacharya , L. Bianchini , T. Boccali , E. Bossini , D. Bruschini , R. Castaldi , M. A. Ciocci , M. Cipriani , V. D'Amantes , R. Dell'Orso , S. Donato , A. Giassi , F. Ligabue , D. Matos Figueiredo , A. Messineo , M. Musich , F. Palla , A. Rizzi , G. Rolandi , S. Roy Chowdhury , T. Sarkar , A. Scribano , P. Spagnolo , R. Tenchini , G. Tonelli , N. Turini , A. Venturi , P. G. Verdini 







INFN Sezione di Roma^a, Sapienza Università di Roma^b, Rome, Italy

P. Barria , M. Campana , F. Cavallari , L. Cunqueiro Mendez , D. Del Re , E. Di Marco , M. Diemoz , F. Errico , E. Longo , P. Meridiani , J. Mijuskovic , G. Organtini , F. Pandolfi , R. Paramatti , C. Quaranta , S. Rahatlou , C. Rovelli , F. Santanastasio , L. Soffi 

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane ^{a,b}, R. Arcidiaconos ^{a,c}, S. Argiro ^{a,b}, M. Arneodo ^{a,c}, N. Bartosik ^a, R. Bellan ^{a,b},
 A. Bellora ^{a,b}, C. Biino ^a, N. Cartiglia ^a, M. Costa ^{a,b}, R. Covarelli ^{a,b}, N. Demaria ^a, L. Finco ^a,
 M. Grippo ^{a,b}, B. Kiani ^{a,b}, F. Legger ^a, F. Luongo ^{a,b}, C. Mariotti ^a, S. Maselli ^a, A. Mecca ^{a,b},
 E. Migliore ^{a,b}, M. Monteno ^a, R. Mulargia ^a, M. M. Obertino ^{a,b}, G. Ortona ^a, L. Pacher ^{a,b}, N. Pastrone ^a,
 M. Pelliccioni ^a, M. Ruspà ^{a,c}, F. Siviero ^{a,b}, V. Sola ^{a,b}, A. Solano ^{a,b}, D. Soldi ^{a,b}, A. Staiano ^a,
 C. Tarricone ^{a,b}, D. Trocino ^a, G. Umoret ^{a,b}, E. Vlasov ^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte ^a, V. Candelise ^{a,b}, M. Casarsa ^a, F. Cossutti ^a, K. De Leo ^{a,b}, G. Della Ricca ^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra ^a, J. Hong ^a, C. Huh ^a, B. Kim ^a, D. H. Kim ^a, J. Kim ^a, H. Lee ^a, S. W. Lee ^a, C. S. Moon ^a, Y. D. Oh ^a,
 M. S. Ryu ^a, S. Sekmen ^a, Y. C. Yang ^a

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak ^a, P. Gwak ^a, H. Kim ^a, D. H. Moon ^a

Hanyang University, Seoul, Korea

E. Asilar ^a, D. Kim ^a, T. J. Kim ^a, J. A. Merlin ^a, J. Park ^a

Korea University, Seoul, Korea

S. Choi ^a, S. Han ^a, B. Hong ^a, K. Lee ^a, K. S. Lee ^a, S. Lee ^a, J. Park ^a, S. K. Park ^a, J. Yoo ^a

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh ^a











Sejong University, Seoul, Korea

H. S. Kim ^a, Y. Kim ^a, S. Lee ^a



Seoul National University, Seoul, Korea

J. Almond ^a, J. H. Bhyun ^a, J. Choi ^a, W. Jun ^a, J. Kim ^a, J. S. Kim ^a, S. Ko ^a, H. Kwon ^a, H. Lee ^a, J. Lee ^a, J. Lee ^a,
 B. H. Oh ^a, S. B. Oh ^a, H. Seo ^a, U. K. Yang ^a, I. Yoon ^a

University of Seoul, Seoul, Korea

W. Jang ^a, D. Y. Kang ^a, Y. Kang ^a, S. Kim ^a, B. Ko ^a, J. S. H. Lee ^a, Y. Lee ^a, I. C. Park ^a, Y. Roh ^a, I. J. Watson ^a,
 S. Yang ^a



Department of Physics, Yonsei University, Seoul, Korea

S. Ha ^a, H. D. Yoo ^a

Sungkyunkwan University, Suwon, Korea

M. Choi ^a, M. R. Kim ^a, H. Lee ^a, Y. Lee ^a, I. Yu ^a


College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy ^a, Y. Maghrbi ^a

Riga Technical University, Riga, Latvia

K. Dreimanis ^a, A. Gaile ^a, G. Pikurs ^a, A. Potrebko ^a, M. Seidel ^a, V. Veckalns ⁵⁸

University of Latvia (LU), Riga, Latvia

N. R. Strautnieks ^a




Vilnius University, Vilnius, Lithuania

M. Ambrozas ^a, A. Juodagalvis ^a, A. Rinkevicius ^a, G. Tamulaitis ^a

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

N. Bin Norjoharuddeen ^a, I. Yusuff ⁵⁹, Z. Zolkapli ^a

Universidad de Sonora (UNISON), Hermosillo, Mexico

J. F. Benitez , A. Castaneda Hernandez , H. A. Encinas Acosta, L. G. Gallegos Mariñez, M. León Coello ,
J. A. Murillo Quijada , A. Sehwat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , E. De La Cruz-Burelo , I. Heredia-De La Cruz ⁶⁰, R. Lopez-Fernandez ,
C. A. Mondragon Herrera, A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , I. Pedraza , H. A. Salazar Ibarguen , C. Uribe Estrada 




University of Montenegro, Podgorica, Montenegro

I. Bubanja, N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P. H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M. I. Asghar, A. Awais , M. I. M. Awan, H. R. Hoorani , W. A. Khan 

AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland

V. Avati, L. Grzanka , M. Malawski 



National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szleper , P. Zalewski 















Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolkowski , A. Muhammad 



Warsaw University of Technology, Warsaw, Poland

K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisbon, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , P. Faccioli , M. Gallinaro ,
J. Hollar , N. Leonardo , T. Niknejad , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J. W. Wulff




























Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

M. Dordevic , J. Milosevic , V. Rekovic














Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Aguilar-Benitez, J. Alcaraz Maestre , Cristina F. Bedoya , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , D. Fernández Del Val , J. P. Fernández Ramos , J. Flix , M. C. Fouz ,
O. Gonzalez Lopez , S. Goy Lopez , J. M. Hernandez , M. I. Josa , J. León Holgado , D. Moran ,
C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo ,
I. Redondo , D. D. Redondo Ferrero , L. Romero, S. Sánchez Navas , L. Urda Gómez , J. Vazquez Escobar ,
C. Willmott




















Universidad Autónoma de Madrid, Madrid, Spain

J. F. de Trocóniz 

Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Universidad de Oviedo, Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero ,
J. R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez ,
A. Trapote , C. Vico Villalba , P. Vischia 





Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , J. A. Brochero Cifuentes , I. J. Cabrillo , A. Calderon ,
 J. Duarte Campderros , M. Fernandez , C. Fernandez Madrazo , G. Gomez , C. Lasasa García ,
 C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos ,
 J. Piedra Gomez , L. Scodellaro , I. Vila , J. M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

M. K. Jayananda , B. Kailasapathy ⁶¹, D. U. J. Sonnadara , D. D. C. Wickramarathna 









University of Ruhuna, Department of Physics, Matara, Sri Lanka

W. G. D. Dharmaratna ⁶², K. Liyanage , N. Perera , N. Wickramage 


















CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez ,
 M. Bianco , B. Bilin , A. A. Bin Anuar , A. Bocci , E. Brondolin , C. Caillol , T. Camporesi , G. Cerminara ,
 N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M. M. Defranchis , M. Deile ,
 M. Dobson , F. Fallavollita ⁶³, L. Forthomme , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege ,
 L. Gouskos , M. Haranko , J. Hegeman , B. Huber, V. Innocente , T. James , P. Janot , J. Kieseler ,
 S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri , M. Mannelli , A. C. Marini ,
 M. Matthewman, F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Moortgat , M. Mulders , S. Orfanelli,
 F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , H. Qu , D. Rabady , G. Reales Gutiérrez,
 M. Rovere , H. Sakulin , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva ,
 P. Sphicas ⁶⁴, A. G. Stahl Leitner , A. Steen , S. Summers , D. Treille , P. Tropea , A. Tsirou, D. Walter ,
 J. Wanczyk ⁶⁵, K. A. Wozniak ⁶⁶, P. Zehetner , P. Zejdl , W. D. Zeuner










Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua ⁶⁷, L. Caminada ⁶⁷, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H. C. Kaestli ,
 D. Kotlinski , C. Lange , M. Missiroli ⁶⁷, L. Noehte ⁶⁷, T. Rohe 

ETH Zurich-Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T. K. Aarrestad , K. Androsov ⁶⁵, M. Backhaus , A. Calandri , C. Cazzaniga , K. Datta , A. De Cosa ,
 G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , D. Hits ,
 W. Lustermann , A.-M. Lyon , R. A. Manzoni , M. Marchegiani , L. Marchese , C. Martin Perez ,
 A. Mascellani ⁶⁵, F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , M. G. Ratti , M. Reichmann ,
 C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, D. A. Sanz Becerra , R. Seidita , J. Steggemann ⁶⁵,
 D. Valsecchi , R. Wallny









Universität Zürich, Zurich, Switzerland

C. Amsler ⁶⁸, P. Bäertschi , C. Botta , D. Brzhechko, M. F. Canelli , K. Cormier , R. Del Burgo, J. K. Heikkilä ,
 M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S. P. Liechti , A. Macchiolo , P. Meiring ,
 V. M. Mikuni , U. Molinatti , I. Neutelings , A. Reimers , P. Robmann, S. Sanchez Cruz , K. Schweiger ,
 M. Senger , Y. Takahashi , R. Tramontano

National Central University, Chung-Li, Taiwan

C. Adloff ⁶⁹, C. M. Kuo, W. Lin, P. K. Rout , P. C. Tiwari ⁴², S. S. Yu 










National Taiwan University (NTU), Taipei, Taiwan








































































































L. Ceard, Y. Chao , K. F. Chen , P.s. Chen, Z.g. Chen, W.-S. Hou , T.h. Hsu, Y.w. Kao, R. Khurana, G. Kole ,
 Y. Y. Li , R.-S. Lu , E. Paganis , A. Psallidas, X.f. Su, J. Thomas-Wilsker , L.s. Tsai, H.y. Wu, E. Yazgan

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand















C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , Z. S. Demiroglu , F. Dolek , I. Dumanoglu ⁷⁰, E. Eskut , Y. Guler ⁷¹,
 E. Gurpinar Guler ⁷¹, C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir ⁷²,
 A. Polatoz , B. Tali ⁷³, U. G. Tok , S. Turkcapar , E. Usulan , I. S. Zorbakir

Middle East Technical University, Physics Department, Ankara, TurkeyM. Yalvac ⁷⁴**Bogazici University, Istanbul, Turkey**B. Akgun , I. O. Atakisi , E. Gülmez , M. Kaya ⁷⁵, O. Kaya ⁷⁶, S. Tekten ⁷⁷**Istanbul Technical University, Istanbul, Turkey**A. Cakir , K. Cankocak ^{70,78}, Y. Komurcu , S. Sen ⁷⁹**Istanbul University, Istanbul, Turkey**O. Aydilek , S. Cerci ⁷³, V. Epshteyn , B. Hacisahinoglu , I. Hos ⁸⁰, B. Isildak ⁸¹, B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , D. Sunar Cerci ⁷³, C. Zorbilmez **Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine**A. Boyaryntsev , B. Grynyov **National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine**L. Levchuk **University of Bristol, Bristol, UK**D. Anthony , J. J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki , J. Goldstein , H. F. Heath , L. Kreczko , B. Krikler , S. Paramesvaran , S. Seif El Nasr-Storey , V. J. Smith , N. Stylianou ⁸², K. Walkingshaw Pass, R. White **Rutherford Appleton Laboratory, Didcot, UK**A. H. Ball , K. W. Bell , A. Belyaev ⁸³, C. Brew , R. M. Brown , D. J. A. Cockerill , C. Cooke , K. V. Ellis , K. Harder , S. Harper , M.-L. Holmberg ⁸⁴, J. Linacre , K. Manolopoulos, D. M. Newbold , E. Olaiya, D. Petyt , T. Reis , G. Salvi , T. Schuh, C. H. Shepherd-Themistocleous , I. R. Tomalin , T. Williams **Imperial College, London, UK**R. Bainbridge , P. Bloch , C. E. Brown , O. Buchmuller, V. Cacchio, C. A. Carrillo Montoya , G. S. Chahal ⁸⁵, D. Colling , J. S. Dancu, P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M. H. Hassanshahi , A. Howard, G. Iles , M. Knight , J. Langford , L. Lyons , A.-M. Magnan , S. Malik, A. Martelli , M. Mieskolainen , J. Nash ⁸⁶, M. Pesaresi, B. C. Radburn-Smith , A. Richards, A. Rose , C. Seez , R. Shukla , A. Tapper , K. Uchida , G. P. Uttley , L. H. Vage, T. Virdee ³², M. Vojinovic , N. Wardle , D. Winterbottom **Brunel University, Uxbridge, UK**K. Coldham, J. E. Cole , A. Khan, P. Kyberd , I. D. Reid **Baylor University, Waco, TX, USA**S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , J. Hiltbrand , A. R. Kanuganti , B. McMaster , M. Saunders , S. Sawant , C. Sutantawibul , M. Toms ⁸⁷, J. Wilson **Catholic University of America, Washington, DC, USA**R. Bartek , A. Dominguez , C. Huerta Escamilla, A. E. Simsek , R. Uniyal , A. M. Vargas Hernandez **The University of Alabama, Tuscaloosa, AL, USA**R. Chudasama , S. I. Cooper , S. V. Gleyzer , C. U. Perez , P. Rumerio ⁸⁸, E. Usai , C. West , R. Yi **Boston University, Boston, MA, USA**A. Akpinar , A. Albert , D. Arcaro , C. Cosby , Z. Demiragli , C. Erice , E. Fontanesi , D. Gastler , S. Jeon , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , S. Yuan **Brown University, Providence, RI, USA**G. Benelli , X. Coubez ²⁷, D. Cutts , M. Hadley , U. Heintz , J. M. Hogan ⁸⁹, T. Kwon , G. Landsberg , K. T. Lau , D. Li , J. Luo , S. Mondal , M. Narain [†], N. Pervan , S. Sagir ⁹⁰, F. Simpson , M. Stamenkovic , W. Y. Wong, X. Yan , W. Zhang

University of California, Davis, Davis, CA, USA

S. Abbott , J. Bonilla , C. Brainerd , R. Breedon , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P. T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , D. Pellett , W. Wei , Y. Yao , F. Zhang 























University of California, Los Angeles, CA, USA

M. Bachtis , R. Cousins , A. Datta , J. Hauser , M. Ignatenko , M. A. Iqbal , T. Lam , E. Manca , D. Saltzberg , V. Valuev 




















University of California, Riverside, Riverside, CA, USA

R. Clare , M. Gordon , G. Hanson , W. Si , S. Wimpenny  [†]










University of California, San Diego, La Jolla, CA, USA

J. G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , M. Pieri , M. Quinnan , B. V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 


Department of Physics, University of California, Santa Barbara, Santa Barbara, CA, USA

A. Barzdukas , L. Brennan , C. Campagnari , G. Collura , A. Dorsett , J. Incandela , M. Kilpatrick , J. Kim , A. J. Li , P. Masterson , H. Mei , M. Oshiro , J. Richman , U. Sarica , R. Schmitz , F. Setti , J. Shephlock , D. Stuart , S. Wang 














California Institute of Technology, Pasadena, CA, USA

A. Bornheim , O. Cerri , A. Latorre , J. M. Lawhorn , J. Mao , H. B. Newman , T. Q. Nguyen , M. Spiropulu , J. R. Vlimant , C. Wang , S. Xie , R. Y. Zhu 

Carnegie Mellon University, Pittsburgh, PA, USA

J. Alison , S. An , M. B. Andrews , P. Bryant , V. Dutta , T. Ferguson , A. Harilal , C. Liu , T. Mudholkar , S. Murthy , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 











































































University of Colorado Boulder, Boulder, CO, USA

J. P. Cumalat , W. T. Ford , A. Hassani , G. Karathanasis , E. MacDonald , N. Manganelli , F. Marini , A. Perloff , C. Savard , N. Schonbeck , K. Stenson , K. A. Ulmer , S. R. Wagner , N. Zipper 























Cornell University, Ithaca, NY, USA

J. Alexander , S. Bright-Thonney , X. Chen , D. J. Cranshaw , J. Fan , X. Fan , D. Gadkari , S. Hogan , J. Monroy , J. R. Patterson , J. Reichert , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, IL, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L. A. T. Bauerdick , D. Berry , J. Berryhill , P. C. Bhat , K. Burkett , J. N. Butler , A. Canepa , G. B. Cerati , H. W. K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V. D. Elvira , Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green , A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R. M. Harris , R. Heller , T. C. Herwig , J. Hirschauer , L. Horyn , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K. H. M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , V. Papadimitriou , N. Pastika , K. Pedro , C. Pena  ⁹¹, F. Ravera , A. Reinsvold Hall  ⁹², L. Ristori , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N. V. Tran , L. Uplegger , E. W. Vaandering , I. Zoi 


University of Florida, Gainesville, FL, USA

C. Aruta , P. Avery , D. Bourilkov , L. Cadamuro , P. Chang , V. Cherepanov , R. D. Field , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. H. Lo , K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , D. Rosenzweig , S. Rosenzweig , K. Shi , J. Wang 























Florida State University, Tallahassee, FL, USA

T. Adams , A. Al Kadhimi , A. Askew , N. Bower , R. Habibullah , V. Hagopian , R. Hashmi , R. S. Kim , S. Kim , T. Kolberg , G. Martinez , H. Prosper , P. R. Prova , O. Viazlo , M. Wulansatiti , R. Yohay , J. Zhang 












Florida Institute of Technology, Melbourne, FL, USA

B. Alsufyani, M. M. Baarmand , S. Butalla , T. Elkafrawy ⁵⁶, M. Hohlmann , R. Kumar Verma , M. Rahmani

University of Illinois Chicago, Chicago, USA

M. R. Adams , C. Bennett, R. Cavanaugh , S. Dittmer , R. Escobar Franco , O. Evdokimov , C. E. Gerber , D. J. Hofman , J. H. Lee , D. S. Lemos , A. H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , T. Roy , S. Rudrabhatla , M. B. Tonjes , N. Varelas , X. Wang , Z. Ye , J. Yoo 

























The University of Iowa, Iowa City, IA, USA

M. Alhusseini , D. Blend, K. Dilsiz ⁹³, L. Emediato , G. Karaman , O. K. Köseyan , J.-P. Merlo, A. Mestvirishvili ⁹⁴, J. Nachtman , O. Neogi, H. Ogul ⁹⁵, Y. Onel , A. Penzo , C. Snyder, E. Tiras ⁹⁶

Johns Hopkins University, Baltimore, MD, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A. V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz , T. Á. Vámi 

The University of Kansas, Lawrence, KS, USA

A. Abreu , L. F. Alcerro Alcerro , J. Anguiano , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , C. Lindsey, J. Marquez , N. Minafra , M. Murray , M. Nickel , M. Pitt , S. Popescu ⁹⁷, C. Rogan , C. Royon , R. Salvatico , S. Sanders , C. Smith , Q. Wang , G. Wilson 



Kansas State University, Manhattan, KS, USA

B. Allmond , A. Ivanov , K. Kaadze , A. Kalogeropoulos , D. Kim, Y. Maravin , K. Nam, J. Natoli , D. Roy , G. Sorrentino 

























Lawrence Livermore National Laboratory, Livermore, CA, USA

F. Rebassoo , D. Wright 

University of Maryland, College Park, MD, USA

A. Baden , A. Belloni , A. Bethani , Y. M. Chen , S. C. Eno , N. J. Hadley , S. Jabeen , R. G. Kellogg , T. Koeth , Y. Lai , S. Lascio , A. C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M. M. Paranjpe, L. Wang 


Massachusetts Institute of Technology, Cambridge, MA, USA

J. Bendavid , W. Busza , I. A. Cali , Y. Chen , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, P. Harris, D. Hoang, D. Kovalskiy , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. Mironov , C. Paus , D. Rankin , C. Roland , G. Roland , S. Rothman , Z. Shi , G. S. F. Stephans , J. Wang, Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, MN, USA

B. Crossman , B. M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , S. Pandey , M. Revering , R. Rusack , R. Saradhy , N. Schroeder , N. Strobbe , M. A. Wadud 










University of Mississippi, Oxford, MS, USA

L. M. Cremaldi 

University of Nebraska-Lincoln, Lincoln, NE, USA

K. Bloom , M. Bryson, D. R. Claes , C. Fangmeier , F. Golf , G. Haza , J. Hossain , C. Joo , I. Kravchenko , I. Reed , J. E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu , A. G. Zecchinelli 










State University of New York at Buffalo, Buffalo, NY, USA

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams 






























Northeastern University, Boston, MA, USA

E. Barberis , Y. Haddad , Y. Han , A. Krishna , J. Li , M. Lu , G. Madigan , R. Mccarthy , D. M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , A. Tishelman-Charny , B. Wang , D. Wood 











Northwestern University, Evanston, IL, USA

S. Bhattacharya , J. Bueghly , Z. Chen , K. A. Hahn , Y. Liu , Y. Miao , D. G. Monk , M. H. Schmitt , A. Taliercio , M. Velasco




University of Notre Dame, Notre Dame, IN, USA

R. Band , R. Bucci , S. Castells , M. Cremonesi , A. Das , R. Goldouzian , M. Hildreth , K. W. Ho , K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli , I. Mcalister , T. McCauley , C. Mcgrady , C. Moore , Y. Musienko ¹⁶, H. Nelson , M. Osherson , R. Ruchti , A. Townsend , M. Wayne , H. Yockey , M. Zarucki , L. Zygala 

The Ohio State University, Columbus, OH, USA

A. Basnet , B. Bylsma , M. Carrigan , L. S. Durkin , C. Hill , M. Joyce , A. Lesauvage , M. Nunez Ornelas , K. Wei , B. L. Winer , B. R. Yates 
























Princeton University, Princeton, NJ, USA

F. M. Addesa , H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , S. Higginbotham , G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , A. Shevelev , D. Stickland , C. Tully 




University of Puerto Rico, Mayaguez, PR, USA

S. Malik 

















Purdue University, West Lafayette, IN, USA

A. S. Bakshi , V. E. Barnes , S. Chandra , R. Chawla , S. Das , A. Gu , L. Gutay , M. Jones , A. W. Jung , D. Kondratyev , A. M. Koshy , M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer , J. F. Schulte , M. Stojanovic , J. Thieman , A. K. Viridi , F. Wang , W. Xie 













Purdue University Northwest, Hammond, IN, USA

J. Dolen , N. Parashar , A. Pathak 


Rice University, Houston, TX, USA

D. Acosta , A. Baty , T. Carnahan , K. M. Ecklund , P. J. Fernández Manteca , S. Freed , P. Gardner , F. J. M. Geurts , A. Kumar , W. Li , O. Miguel Colin , B. P. Padley , R. Redjimi , J. Rotter , E. Yigitbasi , Y. Zhang 






















University of Rochester, Rochester, NY, USA

A. Bodek , P. de Barbaro , R. Demina , J. L. Dulemba , C. Fallon , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , P. Parygin ⁸⁷, E. Popova ⁸⁷, R. Taus , G. P. Van Onsem 

The Rockefeller University, New York, NY, USA

K. Goulianos 

Rutgers, The State University of New Jersey, Piscataway, NJ, USA

B. Chiarito , J. P. Chou , Y. Gershtein , E. Halkiadakis , A. Hart , M. Heindl , D. Jaroslowski , O. Karacheban ³⁰, I. Laflotte , A. Lath , R. Montalvo , K. Nash , H. Routray , S. Salur , S. Schnetzer , S. Somalwar , R. Stone , S. A. Thayil , S. Thomas , J. Vora , H. Wang 

University of Tennessee, Knoxville, TN, USA

H. Acharya , D. Ally , A. G. Delannoy , S. Fiorendi , T. Holmes , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 











Texas A&M University, College Station, TX, USA

D. Aebi , M. Ahmad , O. Bouhali ⁹⁸, M. Dalchenko , R. Eusebi , J. Gilmore , T. Huang , T. Kamon ⁹⁹, H. Kim , S. Luo , S. Malhotra , R. Mueller , D. Overton , D. Rathjens , A. Safonov 









Texas Tech University, Lubbock, TX, USA

N. Akchurin , J. Damgov , V. Hegde , A. Hussain , Y. Kazhykarim , K. Lamichhane , S. W. Lee , A. Mankel , T. Mengke , S. Muthumuni , T. Peltola , I. Volobouev , A. Whitbeck 

Vanderbilt University, Nashville, TN, USA

E. Appelt , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 






















University of Virginia, Charlottesville, VA, USA

B. Cardwell , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , A. Li , C. Neu , C. E. Perez Lara 

Wayne State University, Detroit, MI, USA

P. E. Karchin 

University of Wisconsin-Madison, Madison, WI, USA

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C. K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , D. Pinna, A. Savin, V. Shang , V. Sharma , W. H. Smith , D. Teague, H. F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN, Geneva, Switzerland

S. Afanasiev , V. Andreev , Yu. Andreev , T. Aushev , M. Azarkin , A. Babaev , A. Belyaev ⁸³, V. Blinov ¹⁰⁰, E. Boos , V. Borshch , D. Budkouski , V. Chekhovsky, R. Chistov ¹⁰⁰, M. Danilov ¹⁰⁰, A. Dermenev , T. Dimova ¹⁰⁰, D. Druzhkin ¹⁰¹, M. Dubinin ⁹¹, L. Dudko , A. Ershov , G. Gavrillov , V. Gavrillov , S. Gninenko , V. Golovtsov , N. Golubev , I. Golutvin , I. Gorbunov , A. Gribushin , Y. Ivanov , V. Kachanov , L. Kardapoltsev ¹⁰⁰, V. Karjavine , A. Karneyev , V. Kim ¹⁰⁰, M. Kirakosyan, D. Kirpichnikov , M. Kirsanov , V. Klyukhin , O. Kodolova ¹⁰², D. Konstantinov , V. Korenkov , A. Kozyrev ¹⁰⁰, N. Krasnikov , A. Lanev , P. Levchenko ¹⁰³, N. Lychkovskaya , V. Makarenko , A. Malakhov , V. Matveev ¹⁰⁰, V. Murzin , A. Nikitenko ^{104, 102}, S. Obraztsov , V. Oreshkin , V. Palichik , V. Perelygin , S. Petrushanko , S. Polikarpov ¹⁰⁰, V. Popov , O. Radchenko ¹⁰⁰, M. Savina , V. Savrin , M. Sergeev , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen ¹⁰⁰, S. Slabospitskii , V. Smirnov , A. Snigirev , D. Sosnov , V. Sulimov , E. Tcherniaev , A. Terkulov , O. Teryaev , I. Tlisova , A. Toropin , L. Uvarov , A. Uzunian , A. Vorobyev[†], N. Voytishin , B. S. Yuldashev ¹⁰⁵, A. Zarubin , I. Zhizhin , A. Zhokin 

† Deceased

- 1: Also at Yerevan State University, Yerevan, Armenia
- 2: Also at TU Wien, Vienna, Austria
- 3: Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt
- 4: Also at Ghent University, Ghent, Belgium
- 5: Also at Universidade Estadual de Campinas, Campinas, Brazil
- 6: Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
- 7: Also at UFMS, Nova Andradina, Brazil
- 8: Also at Nanjing Normal University, Nanjing, China
- 9: Now at Henan Normal University, Xinxiang, China
- 10: Now at The University of Iowa, Iowa City, IA, USA
- 11: Also at University of Chinese Academy of Sciences, Beijing, China
- 12: Also at China Center of Advanced Science and Technology, Beijing, China
- 13: Also at University of Chinese Academy of Sciences, Beijing, China
- 14: Also at China Spallation Neutron Source, Guangdong, China
- 15: Also at Université Libre de Bruxelles, Bruxelles, Belgium
- 16: Also at an institute or an international laboratory covered by a cooperation agreement with CERN, Geneva, Switzerland
- 17: Also at Cairo University, Cairo, Egypt
- 18: Also at Suez University, Suez, Egypt
- 19: Now at British University in Egypt, Cairo, Egypt
- 20: Also at Birla Institute of Technology, Mesra, Mesra, India
- 21: Also at Purdue University, West Lafayette, IN, USA
- 22: Also at Université de Haute Alsace, Mulhouse, France

- 23: Also at Department of Physics, Tsinghua University, Beijing, China
- 24: Also at The University of the State of Amazonas, Manaus, Brazil
- 25: Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- 26: Also at University of Hamburg, Hamburg, Germany
- 27: Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- 28: Also at Isfahan University of Technology, Isfahan, Iran
- 29: Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- 30: Also at Brandenburg University of Technology, Cottbus, Germany
- 31: Also at Forschungszentrum Jülich, Juelich, Germany
- 32: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- 33: Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
- 34: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- 35: Now at Universitatea Babeş-Bolyai-Facultatea de Fizica, Cluj-Napoca, Romania
- 36: Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- 37: Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- 38: Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary
- 39: Also at Punjab Agricultural University, Ludhiana, India
- 40: Also at University of Hyderabad, Hyderabad, India
- 41: Also at University of Visva-Bharati, Santiniketan, India
- 42: Also at Indian Institute of Science (IISc), Bangalore, India
- 43: Also at IIT Bhubaneswar, Bhubaneswar, India
- 44: Also at Institute of Physics, Bhubaneswar, India
- 45: Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- 46: Also at Department of Physics, Isfahan University of Technology, Isfahan, Iran
- 47: Also at Sharif University of Technology, Tehran, Iran
- 48: Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- 49: Also at Helwan University, Cairo, Egypt
- 50: Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- 51: Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- 52: Also at Università degli Studi Guglielmo Marconi, Rome, Italy
- 53: Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Naples, Italy
- 54: Also at Fermi National Accelerator Laboratory, Batavia, IL, USA
- 55: Also at Università di Napoli 'Federico II', Naples, Italy
- 56: Also at Ain Shams University, Cairo, Egypt
- 57: Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- 58: Also at Riga Technical University, Riga, Latvia
- 59: Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- 60: Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- 61: Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- 62: Also at Saegis Campus, Nugegoda, Sri Lanka
- 63: Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy
- 64: Also at National and Kapodistrian University of Athens, Athens, Greece
- 65: Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- 66: Also at University of Vienna Faculty of Computer Science, Vienna, Austria
- 67: Also at Universität Zürich, Zurich, Switzerland
- 68: Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- 69: Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- 70: Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- 71: Also at Konya Technical University, Konya, Turkey
- 72: Also at Izmir Bakircay University, Izmir, Turkey
- 73: Also at Adiyaman University, Adiyaman, Turkey
- 74: Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey

- 75: Also at Marmara University, Istanbul, Turkey
76: Also at Milli Savunma University, Istanbul, Turkey
77: Also at Kafkas University, Kars, Turkey
78: Now at Istanbul Okan University, Istanbul, Turkey
79: Also at Hacettepe University, Ankara, Turkey
80: Also at Faculty of Engineering, Istanbul University-Cerrahpasa, Istanbul, Turkey
81: Also at Yildiz Technical University, Istanbul, Turkey
82: Also at Vrije Universiteit Brussel, Brussel, Belgium
83: Also at School of Physics and Astronomy, University of Southampton, Southampton, UK
84: Also at University of Bristol, Bristol, UK
85: Also at IPPP Durham University, Durham, UK
86: Also at Monash University, Faculty of Science, Clayton, Australia
87: Now at an institute or an international laboratory covered by a cooperation agreement with CERN, Geneva, Switzerland
88: Also at Università di Torino, Turin, Italy
89: Also at Bethel University, St. Paul, MN, USA
90: Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
91: Also at California Institute of Technology, Pasadena, CA, USA
92: Also at United States Naval Academy, Annapolis, MD, USA
93: Also at Bingol University, Bingol, Turkey
94: Also at Georgian Technical University, Tbilisi, Georgia
95: Also at Sinop University, Sinop, Turkey
96: Also at Erciyes University, Kayseri, Turkey
97: Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
98: Also at Texas A&M University at Qatar, Doha, Qatar
99: Also at Kyungpook National University, Daegu, Korea
100: Also at another institute or international laboratory covered by a cooperation agreement with CERN, Geneva, Switzerland
101: Also at Universiteit Antwerpen, Antwerpen, Belgium
102: Also at Yerevan Physics Institute, Yerevan, Armenia
103: Also at Northeastern University, Boston, MA, USA
104: Also at Imperial College, London, UK
105: Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan