

UC San Diego

UC San Diego Previously Published Works

Title

A cell cycle checkpoint for the endoplasmic reticulum.

Permalink

<https://escholarship.org/uc/item/98x3j4rk>

Journal

Biochimica et biophysica acta. Molecular cell research, 1867(12)

ISSN

1879-2596

Author

Niwa, Maho

Publication Date

2020-08-20

Peer reviewed

A Cell Cycle Checkpoint for the Endoplasmic Reticulum

Maho Niwa

Division of Biological Sciences, Section of Molecular Biology
University of California San Diego
NSB#1, Rm 5328, 9500 Gilman Drive, La Jolla, CA 92093-0377

Corresponding author: Maho Niwa (mniwarosen@ucsd.edu)

Abstract (150 words)

The generation of new cells is one of the most fundamental aspects of cell biology. Proper regulation of the cell cycle is critical for human health, as underscored by many diseases associated with errors in cell cycle regulation, including both cancer and hereditary diseases. A large body of work has identified regulatory mechanisms and checkpoints that ensure accurate and timely replication and segregation of chromosomal DNA. However, few studies have evaluated the extent to which similar checkpoints exist for the division of cytoplasmic components, including organelles. Such checkpoint mechanisms might be crucial for compartments that cannot be generated *de novo*, such as the endoplasmic reticulum (ER). In this review, we highlight recent work in the model organism *Saccharomyces cerevisiae* that led to the discovery of such a checkpoint that ensures that cells inherit functional ER into the daughter cell.


Evidence for cell cycle checkpoints for dividing the cytoplasm

When a eukaryotic cell undergoes division, it faces the challenge of generating genetically identical daughter cells (Ciccia and Elledge, 2010; Matellan and Monje-Casas, 2020; Musacchio and Desai, 2017). The cell must copy the entire genome—which consists of over three billion base pairs in human—without making major mistakes. This process, called DNA replication, must occur within a specific length of time. Handling three billion base pairs alone also imposes spatial and temporal constraints to the cell. In order to ensure that the genome is separated into two dividing cells, these events are assisted by several “checkpoints” at the heart of the cell cycle operation. If mistakes are found, the cell temporally halts cell division, providing an opportunity for fixing mistakes. Upon recovery, the cell resumes the cell cycle at the point at which it stopped to finish generating a new daughter cell. The importance of cell cycle checkpoints has been underscored by many human diseases, such as cancer, that occur due to the failure of the cell to recognize or fix mistakes (Holland and Cleveland, 2012). Each year, incredible resources are poured into efforts to better understand cell cycle mechanisms and

1 checkpoints with the hope of generating more effective treatments for cancer and
2 other diseases.

3
4 A unique feature of eukaryotic cells is the compartmentalization of specific
5 cellular functions into organelles that are surrounded by unique sets of membranes
6 (**Figure 1**). This framework allows cellular functions to fine-tune themselves

7 discretely, but also requires these organelles to
8 be divided during cell division. In contrast to
9 genome replication and separation, however,
10 we know relatively little about the rules that
11 govern the division of cytoplasmic components
12 or organelles (Mascanzoni et al., 2019; Vevea et
13 al., 2014; Weisman, 2003). Historically, almost



14 all cell division studies have focused on issues associated with the genome and the
15 cell cycle checkpoints ensuring that all dividing cells end up with completely and
16 accurately replicated DNA. Some experimental results suggest that the division of
17 non-genomic components, such as proteins and organelles, might be regulated
18 during the cell cycle. For example, treating mammalian cells with DNA-damaging
19 agents results in a massive reorganization of the Golgi, revealing an intimate link
20 between cell cycle DNA replication and division of Golgi (Farber-Katz et al., 2014).
21 Furthermore, these results hint at the presence of cell cycle checkpoints for the
22 proper division of other organelles, such as the endoplasmic reticulum (ER),
23 mitochondria, and endosomes, although relatively little is known about the
24 molecular mechanisms underlying their regulation. In contrast, a recent study
25 reported that the inheritance of a vacuole from the mother cell can fail without
26 halting the cell cycle, as long as a new functional vacuole can be generated in the
27 daughter cell (Jin and Weisman, 2015), revealing that there may be different
28 strategies for ensuring the presence of functional organelles in newly generated
29 cells.

30
31 In recent years, studies have begun to unravel the molecular mechanisms
32 that ensure the inheritance of functional ER to a daughter cell. Here, we provide an

overview of an exciting discovery of a cell cycle regulatory checkpoint—termed the ER stress surveillance (ERSU) cell cycle checkpoint—in the budding yeast *Saccharomyces cerevisiae* (*S. cerevisiae*) (Babour et al., 2010; Chao et al., 2019; Pina et al., 2018; Pina et al., 2016; Pina and Niwa, 2015). Since the initial discovery of the ERSU pathway, subsequent work has provided intriguing mechanistic insights on the ERSU cell cycle checkpoint. Specifically, a recent finding surprisingly revealed that increased levels of phytosphingosine (PHS), an early biosynthetic intermediate of sphingolipids, triggers the ERSU cell cycle checkpoint. As the ERSU pathway is one of the few cell cycle regulatory checkpoints for non-genomic components with significant mechanistic understanding, the principles that operate the ERSU checkpoint may provide fundamental insights into cell cycle checkpoints for other components.

The endoplasmic reticulum is a gateway of the secretory pathway

The ER is one of the biggest organelles in eukaryotic cells, with an extensive reticular structure that can spread throughout large portions of the cytoplasm. Indeed, approximately half of the total membrane of a cell consists of the ER (Alberts et al., 2002). The ER membrane is contiguous with the outer nuclear membrane, which is connected to the inner nuclear membrane via the nuclear pore complex. In addition, recent studies have revealed that the ER membrane establishes contact sites with other organelles, including mitochondria, endosomes, and peroxisomes, with certain functional significance (Cohen et al., 2018; Farre et al., 2019; Helle et al., 2013; Kvam and Goldfarb, 2006; Tamura et al., 2019; Wu et al., 2018). Whether the contact sites remain during cell division and how they affect cell cycle division are still unclear.

Beyond interacting with other essential organelles, the ER plays critical roles in various cellular functions. For example, the ER generates mature, fully functional proteins like growth factors and hormones, which are destined to be secreted outside of the cell. Similarly, the ER produces properly folded, mature transmembrane cell-surface receptors and receptor kinases. As they emerge from

1 the polyribosomes, these proteins, collectively called secretory pathway proteins,
2 are targeted and translocated into the ER lumen as linear unmodified polypeptides
3 for folding, modification, and other maturation steps (Aviram and Schuldiner, 2017;
4 Brown et al., 1995; Walter et al., 1984). Secretory pathway proteins consist of one-
5 third of the total proteome of eukaryotic cells, making this one of the primary tasks
6 of the ER. Upon association with ER-resident chaperones and other protein-folding
7 components, nascent polypeptides undergo maturation steps including folding,
8 modification, and complex formation in order to generate fully folded functional
9 proteins (Ma and Hendershot, 2001; Matlack et al., 1998; McMaster, 2001;
10 Meldolesi and Pozzan, 1998; Voeltz et al., 2002). The folding processes are
11 regulated in such a way that only properly folded proteins can exit from the ER,
12 heading towards the Golgi and their final destinations. Somehow, proteins that
13 have not completed the folding steps are recognized and remain in the ER lumen
14 to complete the process before exiting to the Golgi. Any permanently misfolded
15 proteins are also recognized by the ER and are ultimately degraded by a
16 mechanism specific to permanently misfolded proteins; this process is called ER-
17 associated degradation (**ERAD**) (Berner et al., 2018; Goder et al., 2019; Hampton
18 and Sommer, 2012; Johnson and DeBose-Boyd, 2018; McCaffrey and Braakman,
19 2016; Mehrtash and Hochstrasser, 2019; Pobre et al., 2019; Sun and Brodsky,
20 2019; Wu and Rapoport, 2018). During its lifetime, a cell encounters a series of
21 conditions necessary to produce high levels of secreted proteins. Collectively, the
22 increased need for producing secretory pathway proteins or the accumulation of
23 unfolded proteins is termed "*ER stress*". When a cell recognizes ER stress, it
24 triggers an intracellular signal-transduction pathway called the unfolded protein
25 response (**UPR**), which helps the cell keep up with the increased demands of ER
26 functions (Mori et al., 2000; Ron and Walter, 2007; Rutkowski and Kaufman, 2004).
27 In addition to producing secretory pathway proteins, the UPR regulates ERAD.
28 Thus, via the UPR, the ER regulates and meets the overall composition of the
29 cellular proteome (Hampton and Sommer, 2012; Pobre et al., 2019; Sun and
30 Brodsky, 2019; Wu and Rapoport, 2018). The ER also plays vital roles in lipid
31 biosynthesis and metabolism. Nearly all of the initial steps of cellular lipid synthesis
32 occur on the ER membrane. The ER also plays other important functions, such as

1 regulating cellular Ca²⁺ levels or detoxifying unwanted chemicals, and thus it is a
2 multi-faced functional compartment (Karagas and Venkatachalam, 2019; Mittal et
3 al., 2015; Pierro et al., 2019; Preuss et al., 1991). Importantly, organelles such as
4 the ER cannot be generated *de novo*; therefore, the ER must be inherited from the
5 mother cell.

6 7 **Division of yeast ER during the cell cycle**

8
9 Cell cycle division of the ER is better described in yeast, partly because the yeast
10 ER is less complex than the ER in human cells: the tubular ER surrounds the cortex
11 of the cell and is termed cortical ER (**cER**). The ER also surrounds the nucleus,
12 comprising the perinuclear ER (pnER), which is contiguous with the outer nuclear
13 membrane (Voeltz et al., 2002). The nuclear pore complexes, large
14 macromolecular complexes, in essence, physically separate the inner nuclear
15 membrane from the outer nuclear membrane and the yeast pnER (Forbes et al.,
16 2015; Schwartz et al., 2015). The tubular ER structures (or ER tubules) are
17 approximately 50-100 nm in diameter and form a network of membranous tubules
18 that connect the cER to the pnER (Koning et al., 2002; Preuss et al., 1991). Under
19 normal growth conditions, a tubular ER emanating from the mother pnER starts to
20 grow and enters the daughter cell by passing through the septin ring at the bud
21 neck during the G1/S phase of the cell cycle (Du et al., 2001; Estrada de Martin et
22 al., 2005; Estrada et al., 2003). Here, we refer to this tubular ER as the “*initial ER*
23 *tubule*” to distinguish from other tubular ER linking both the pnER and the cER
24 (**Figure 2**). The growth of the *initial ER tubule* is directed (**Figure 2, step I & II**)
25 and eventually anchored to the bud tip by its interactions with proteins or protein
26 complexes localized at the bud tip (**step III**), including the exocyst complex
27 (Heider and Munson, 2012; Munson and Novick, 2006). Upon anchoring to the bud
28 tip, the *initial ER tubule* changes the direction of its growth, and spreads the
29 tubular ER along the cortex of the daughter cell (**step IV**). Studies have shown that
30 the phosphatase Ptc1, which is localized at the bud tip, play important roles in re-
31 directing tubular ER growth to the lateral direction (Du et al., 2006), leading to the
32 generation of the cER in the daughter cell. Once the cER is inherited in the

daughter cell, the pnER and nucleus enter the daughter cells (**Figure 2, step V**), followed by nuclear division (**step VI**). The movement of the cER into the daughter cell has been reported to occur on the actin cable; Myo4 and She3 connect the cER to the actin cable. By contrast, the movement of the pnER is mediated by microtubules (Estrada et al., 2003). Currently, we lack a detailed understanding of the specific movements of the tubular ER linking the cER and the pnER. In addition, it remains unknown whether the movement of the Initial ER tubule (**IET**) depends on cables of actin or microtubules. Furthermore, the factors determining the number or location of such tubular ER remain elusive. Regardless of the tubular ER, it is unclear how the movement of the cER depends on actin cables while that of the pnER occurs in a microtubule-dependent manner, when the lumen of both types of ER is connected. Additionally, the movement of the pnER must be coordinated with that of the nucleus and the nuclear components. Further work is required to determine how such maneuvers are achieved in a coordinated manner with the division of the genome. Ultimately, beyond the ER and the nuclear transport, transit of other organelles through the bud neck will also have to take place before committing to undergo cytokinesis (**Figure 2**).

The functional importance of the ER and the fact that the ER cannot be synthesized *de novo* suggest that the inheritance of the functional ER must be tightly regulated. This begs interesting questions: for example, what happens if the size or functional capacity of the ER is not adequate enough to divide into two cells? If the ER does not meet the functional demands of a cell or if the ER is too small to divide into two daughter cells, these conditions might impact the progression of the cell cycle. Moreover, it is unclear whether such conditions cause the cell cycle to stop at a specific point. Given that inheriting an accurate and complete set of chromosomal DNA is fundamental to the life of a cell, most studies on cell cycles, thus far, have focused on how cells maintain accurately replicated DNA and how sister chromatids are separated in a faithful and timely manner.

1 **A cell cycle checkpoint for ER division**

2 From the late 1980s through the 1990s, the use of various DNA damage
3 agents and the isolation of conditional cell cycle mutants in yeast opened the door
4 for studying cell cycle mechanisms. These methods brought about a flurry of
5 fascinating discoveries on cell cycle checkpoints for ensuring the inheritance and
6 division of genomic information (Elledge and Harper, 1994; Liu et al., 2000;
7 Morgan, 1995; Zhou and Elledge, 2000). Given the major mechanistic differences
8 between yeast and mammalian cells in terms of cell cycle modes, researchers
9 initially predicted that the regulatory checkpoints for the major cell cycle events
10 would be different for yeast and mammalian cells. However, it turned out that
11 several key regulatory events and the timing for ensuring accurate chromosomal
12 DNA replication and sister chromatid separation are remarkably conserved across
13 yeast and mammalian cells.

14 Motivated by the discovery of cell cycle checkpoints for the genome, new
15 approaches of using well-characterized chemical agents to diminish ER function in
16 yeast cells have created opportunities for (1) learning how cells handle functionally
17 stressed ER during the cell cycle and (2) identifying cell-cycle checkpoints for
18 inheriting functional ER. The chemical agents tunicamycin (Tm) and 1,4-
19 Dithiothreitol (DTT) stress ER function by inhibiting glycosylation and disrupting
20 disulfide bonds, respectively. Indeed, previous studies have shown that triggering
21 ER stress by treatment of yeast cells with either of these agents blocks cells in the
22 G1/S phase (Arnold and Tanner, 1982; Vai et al., 1987) or leads to a *chromosome*
23 *maintenance* defect (Henry et al., 2010). Similar experiments in mammalian cells
24 also revealed that ER stress leads to a G1/S phase block (Cullinan and Diehl, 2006).
25 Subsequent studies, however, found that this G1/S phase arrest stemmed from the
26 lack of components required for budding and/or bud growth. For example, the
27 emergence and growth of the daughter cell requires properly localized polarisome
28 components at the incipient bud site (McMillan et al., 1998). The subcellular
29 localization of the polarisome subunits (e.g., EPO1, SPA2, BEM3, PEA1, BNI1, BUD6,
30 and MSB3/MSB4) to the incipient bud site depends on the early secretory pathway
31 and the functional capacity of the ER, which is the gatekeeper of the secretory

1 pathway (Bidlemaier and Snyder, 2004; Li et al., 2013). Tm treatment of
2 unbudded cells with unfolding agents at the beginning of the cell cycle causes
3 polarisome subunits to unfold, thus preventing them from exiting the ER and
4 heading towards the incipient bud site. Ultimately, the diminished localization of
5 the polarisome subunit contributes to blocking bud emergence and bud growth.
6 Thus, the primary cause of cell cycle block is the lack of the functional polarisome
7 components required for the budding step. Although these studies uncovered that
8 the lack of bud emergence ultimately induces a “morphogenesis block”, they did
9 not reveal a direct impact of the ER or ER function itself on the cell cycle (Bonilla
10 and Cunningham, 2003). The disrupted localization of the polarisome was
11 overcome by an ER stress cell cycle assay that introduced a delay in the timing of
12 ER stress induction until polarisome components could reach the incipient bud site
13 and no longer induce the morphogenesis checkpoint (Bicknell et al., 2007). This
14 delay, introduced to alpha factor-synchronized cells successfully induces ER stress
15 without bud emergence block, making it possible to investigate if and how the
16 steps in the cell cycle are impacted by ER stress (Bicknell et al., 2007). Use of this
17 ER stress cell cycle assay led to the finding that a functionally stressed ER causes
18 cytokinesis block.

19
20 Subsequent work investigated how ER stress induces cytokinesis block.
21 Successful cytokinesis requires that cortical actin patches become polarized to
22 either side of the bud neck late in the cell cycle (Doyle and Botstein, 1996;
23 Kilmartin and Adams, 1984; Mulholland et al., 1994; Novick and Botstein, 1985;
24 Waddle et al., 1996). Actin visualized upon staining with Fluor546-phalloidin
25 revealed that actin patches are localized to the cortex of the cells throughout the
26 cell cycle and become redistributed just prior to cytokinesis, regardless of ER stress
27 (Bicknell et al., 2007). Thus, the ER stress-induced cytokinesis defect is not caused
28 by a delay or an alteration in the actin patch redistribution. Instead, further
29 evidence suggests that ER stress-induced cytokinesis block is caused, at least in
30 part, by the mislocalization of the septin ring. The next section summarizes the
31 studies that have investigated the consequences of having functionally stressed ER
32 on the budding yeast cell cycle, ultimately defining the ERSU cell cycle checkpoint.

The ERSU cell cycle checkpoint hallmark event 1: cER inheritance block

The finding that ER stress causes cell cycle block underscores the importance of stopping cell division if ER function is compromised. Based on the yeast ER inheritance mechanism described, there might be at least two ways to prevent the daughter cell generation if the ER is functionally stressed: First, (1) ER inheritance into the daughter cell is not sensitive to ER stress, but ER stress-induced cytokinesis block somehow ensures that the generation of a daughter cell with non-functional ER is prevented. Alternatively, (2) the inheritance of the stressed ER is blocked and the lack of a functional ER in the daughter cell prevents cytokinesis to separate mother and daughter cells. To distinguish between these possibilities, the impact of ER stress on ER inheritance was initially visualized by an ER reporter, a fusion protein between GFP and the N-terminal transmembrane domain (amino acids 1-702) of HMG-CoA reductase isozyme 1 (Babour et al., 2010; Du et al., 2001; Hampton et al., 1996). In either case, demonstrating that stressed ER directly impacts cell cycle progression would support the idea that ER functional homeostasis is ensured during the cell cycle.

Monitoring the impact of ER stress on ER behaviors revealed that ER stress induction leads to daughter cells without the cER (Babour et al., 2010). This impact was most prominent in class of cells with a small bud index (ratio between mother and daughter cell size) (**Figure 3**). In contrast, cells with larger bud sizes, with or without a divided pnER (class II or III cells, respectively), showed minimal impact of ER stress on ER inheritance. Based on the mechanism of ER entry into the daughter cell (**Figure 2**), cells with medium/larger buds most likely have had (1) the cER inherited prior to ER stress exposure or (2) the initial ER tubule already committed to remain in the daughter. In those cases, the committed cER in the daughter cell would be unlikely to be released and returned back to the mother cell. Indeed, researchers demonstrated this experimentally using synchronized cells (Bicknell et al., 2007): ER stress at the early stage of the cell cycle and prior to commitment of the cER's presence in the daughter cell (**Figure 3**) blocked the cER from entering

the daughter cell (Pina and Niwa, 2015). This finding indicates that cells can (1) recognize the functional capacity of the ER at an early point of the cell cycle and (2) block the inheritance of a stressed ER into the daughter cell. Interestingly, if cells encountered ER stress after cER establishment in the daughter cell (**Figure 2, Steps IV-VI**), cells underwent cytokinesis to separate into mother and daughter cells. In the subsequent round of the cell cycle, however, cER inheritance was blocked, ultimately resulting in cytokinesis block. These results revealed that cytokinesis block occurs in ER-stressed cells to prevent the generation of cells without the presence of an ER.

ERSU checkpoint hallmark event 2: Septin ring transfer from the bud neck

Septin has been reported to establish the diffusion barrier between mother and daughter cells. Studies using N-terminally green fluorescent protein (GFP)-tagged individual septin subunits (CDC3, SHS1, CDC10, CDC11, and CDC12) in unstressed or ER-stressed yeast cells found that the multi-subunit septin ring moves away from the bud neck and re-localizes elsewhere in response to ER stress (**Figure 3**) (Babour et al., 2010; Chao et al., 2019). At the beginning of the cell cycle – even prior to bud emergence – five septin subunits start to assemble at the incipient bud site to generate septin rings in yeast. Once assembled, the septin ring functions as a diffusion barrier of cellular components/organelles from the mother to the daughter cell (Caudron and Barral, 2009; Chao et al., 2014; Clay et al., 2014). Towards the end of the cell cycle, the septin ring splits vertically into two rings, leading to the division of the cytoplasm. Finally, after cell division the septin ring disassembles into each subunit, prior to reassembly at the new bud site (Field and Kellogg, 1999; Versele and Thorner, 2005).

ER stress does not seem to affect the initial targeting of septin subunits: the formation of septin rings at the bud neck appears as a ring-like structure with similar kinetics during ER stress, rather than being dispersed throughout the cytoplasm in unstressed cells. Upon ER stress induction, the septin ring is '*misplaced*' or moves away from the bud neck to elsewhere on the cell surface.

1 Interestingly, the ring-like appearance is retained at the new location, even though
2 it is no longer localized at the bud neck. As the septin ring is a major player in
3 cytokinesis, its absence at the bud neck under ER stress should contribute
4 significantly to cytokinesis block. Furthermore, mislocalized septin ring persists
5 once it moves away from the bud neck to the new location, until ER function is re-
6 established and the cell prepares to re-enter the cell cycle. A more recent study
7 reported the surprising finding that the septin ring is misplaced to the bud scars in
8 response to ER stress. Furthermore, among five subunits, mislocalization of the
9 Shs1 subunit to the bud scar is a key determinant of outcome, with failure to move
10 Shs1 to the bud scar diminishing the ability of the cells to re-enter the cell cycle
11 upon re-establishment of ER homeostasis (Chao et al., 2019). These findings
12 revealed an unprecedented feature of the ERSU checkpoint. The mechanistic
13 details and functional implications of ER stress-induced septin ring mislocalization
14 will be discussed elsewhere (a review in preparation).

17 **The ERSU pathway is independent of the UPR**

18 In *S. cerevisiae*, ER stress induces: (1) ER inheritance block, (2) septin ring
19 mislocalization, resulting in (3) cytokinesis block. However, a major question
20 remained as to whether these events represent a part of the cell cycle checkpoint
21 mechanism for ensuring that a daughter cell receives a functional ER. Another
22 possibility is that these events happen to occur upon ER stress but are unrelated
23 events and do not necessarily represent a cell cycle checkpoint. A signaling
24 pathway known to be activated by ER stress is the UPR (Ron and Walter, 2007;
25 Schroder and Kaufman, 2005). The UPR in yeast cells is initiated by a single-span
26 ER transmembrane protein called Ire1, which contains an *ER luminal domain* that
27 functions as an ER stress sensor (Cox et al., 1993; Mori et al., 1993).
28 Mechanistically, ER stress is sensed by dissociation of the ER chaperone Kar2/BiP
29 from the ER luminal domain of Ire1 (Bertolotti et al., 2000; Kimata et al., 2007;
30 Oikawa et al., 2007), and the Ire1 ER luminal domain recognizes certain features of
31 unfolded proteins (Credle et al., 2005; Zhou et al., 2006). These events lead to Ire1
32 dimerization/oligomerization and autophosphorylation through its Ire1 kinase

domain (Shamu and Walter, 1996) and activation of the sequence-specific endoribonuclease (RNase) domain. Activated Ire1 RNase cleaves the UPR intron of *HAC1* mRNA, encoding for an UPR-specific transcription factor (Cox et al., 1996; Mori et al., 2000), followed by ligation of two *HAC1* mRNA exons by tRNA ligase (Sidrauski et al., 1996). The splicing of *HAC1* mRNA is a key regulatory step for the UPR, as it causes a frameshift, removing a stop codon within the intron and allowing the second exon coding sequences in frame with the coding sequence of the first exon. The second exon contains a transcriptional activation domain and, thus, Ire1-dependent splicing of *HAC1* mRNA is required for the generation of a UPR-specific transcription factor. Spliced Hac1 protein activates the transcription of UPR target genes that ultimately help to re-establish ER functional homeostasis. The UPR, particularly IRE1, is an ancient pathway conserved in all eukaryotic cells (Walter and Ron, 2011).

Given the functional roles of the UPR (Back and Kaufman, 2012; Mori, 2009; Ron and Walter, 2007), it was reasonable to consider that the UPR plays a role in inducing ER inheritance block, septin ring transfer, and cytokinesis block in response to ER stress. It was, therefore, a great surprise when these cell cycle events turned out to not require IRE1 function. In *ire1Δ* cells, ER stress caused ER inheritance block, septin ring transfer, and cytokinesis block at levels similar to wild-type (WT) cells. The lack of UPR involvement suggested the presence of a new pathway that induces these events in response to ER stress.

Identifying the first ERSU component

The lack of IRE1 involvement in the ERSU checkpoint suggested the presence of an independent pathway parallel to the UPR that coordinates the functional status of the ER with its inheritance during the cell cycle, leading to a quest for the ERSU components. The lack of an ERSU component should prevent ER inheritance block, septin ring mislocalization, and temporary cytokinesis block, having deleterious consequences on cell growth. Thus, yeast cells carrying the

deletion of a gene coding for a protein either physically or functionally linked to the ER were screened for their ability to (1) grow on media containing ER stress-inducing agents such as Tm or DTT, (2) block cER inheritance, and (3) mislocalize the septin ring under ER stress. This screen identified SLT2, which encodes a MAP kinase (6). ER-stressed *slt2* knockout (*slt2Δ*) cells were unable to block cER inheritance, mislocalize septin ring, or grow on Tm or DTT plates. Importantly, *slt2Δ* cells were able to activate the UPR pathway, since *HAC1* mRNA was spliced normally with similar kinetics as in WT cells, revealing that the functional homeostasis of the ER was disrupted in *slt2Δ* cells. The inability of *slt2Δ* cells to mount all the ERSU phenotypes demonstrated that the ERSU cell cycle phenotypes are not individual phenotypes that happen to occur under ER stress, but rather are linked under a specific cell cycle regulatory event. Despite the normal appearance of septin rings and cER inheritance, *slt2Δ* cells were incapable of cell growth under ER stress, illustrating the importance of SLT2 in cell cycle progression. Additionally, these findings revealed that cER inheritance block and septin ring transfer are critical for survival during ER stress. Taken together, a SLT2-dependent, but UPR-independent, ERSU cell cycle regulatory pathway is critical for cells to respond to ER stress (**Figure 4**).

The ERSU cell cycle checkpoint is independent of the cell wall integrity pathway

In addition to SLT2, a MAP kinase, its upstream kinases PKC1, BCK1, MKK1, and MKK2 were also found to be involved in the ERSU pathway (Babour et al., 2010). Furthermore, the cell surface component Wsc1 – but not other isoforms of Wsc proteins such as Wsc2, 3, and 4 – plays a role in the ERSU pathway. Some of these components are also involved in the cell wall integrity (CWI) pathway, which is induced by excess turgor pressure against the cell wall. For example, the CWI pathway in *S. cerevisiae* involves PKC1-SLT2 activation via Wsc1 and Wsc2. However, several experimental results clearly distinguish the ERSU checkpoint from the CWI pathway: (1) Wsc1 but not Wsc2 is involved in ERSU pathway events, such as ER inheritance block, septin ring transfer, and cytokinesis block. This is in

1 contrast to the CWI pathway, which depends on both Wsc1 and Wsc2. (2) Treating
2 cells with a known CWI pathway agonist, such as calcofular white, does not
3 activate the ERSU pathway. (3) A Wsc1-AAA mutation that disrupts the
4 internalization of Wsc1 from the cell surface has no impact on ERSU checkpoint
5 activation, despite blocking the induction of the CWI pathway. Taken together, the
6 ER stress-induced ERSU checkpoint is distinct from the previously described CWI
7 pathway.

8
9 Loss of the ability of *ire1Δ* cells to sustain their cell growth under ER stress
10 provided the functional significance of the UPR signaling pathway in response to ER
11 stress. Without the UPR pathway, cells are incapable of handling the accumulation
12 of unfolded proteins in the ER lumen. The lack of *slt2Δ* cell growth on Tm-
13 containing media also reveals the functional importance of the ERSU pathway.
14 Importantly, the cell growth of ER-stressed *slt2Δ* cells can be effectively rescued by
15 preventing the inheritance of the stressed ER into the daughter cell. This has been
16 achieved, for example, by the addition of a small quantity of nocodazol, an agent
17 that prevents microtubule polymerization, by the use of *actin1-1*, a temperature-
18 sensitive Actin mutation, or by the use of *myo4Δ* cells. Both Actin and Myo4 are
19 required for the entry of the ER into the daughter cell (Estrada et al., 2003). Results
20 of these experiments revealed that the presence of stressed ER in the daughter
21 cell leads to cell death.

24 **Initiation of the ERSU cell cycle checkpoint**

25
26 The discovery that the ERSU is an independent signaling pathway induced
27 by ER stress beyond the well-studied UPR pathway, and the identification of SLT2
28 as a key ERSU player, was rather unexpected and generated excitement. However,
29 the initiators of the ERSU cell cycle checkpoint remained a mystery. SLT2 plays
30 roles in a wide range of signaling events, including the CWI response and gene
31 silencing events (Jimenez-Gutierrez et al., 2020; Levin and Errede, 1995).
32 Consistent with being a multi-functional protein, SLT2 is localized throughout the

1 cytoplasm under both normal growth and ER stress conditions. However, SLT2 has
2 not been reported to associate physically with the ER membrane. Since the ERSU
3 hallmark events such as cER inheritance block, septin ring mislocalization, and
4 cytokinesis block, are initiated in response to ER stress, the initiation step is
5 expected to occur on the ER membrane.

6
7 Another unanticipated finding led to the identification of ERSU-initiating
8 components on the ER membrane (Pina et al., 2016). This began with the puzzling
9 observation of a differential impact of ER stress on the inheritance of the pnER and
10 cER: ER stress induction blocked cER inheritance, whereas the pnER, contiguous
11 with the cER and the inner nuclear membrane, was inherited normally into the
12 daughter cell even under ER stress (Babour et al., 2010). Since the lumen of the
13 cER and the pnER are contiguous via the presence of the tubular ER, the different
14 responses between the cER and pnER were rather difficult to reconcile. A potential
15 explanation for the observed differences between the cER and the pnER might
16 stem from the spatial differences in the extent of ER stress. Surprisingly, this effort
17 to provide a molecular explanation for these differences led to the identification of
18 ERSU components.

19
20 During ER stress, Kar2/BiP binding to the elevated levels of unfolded client
21 proteins increases, resulting in the reduction of Kar2/BiP mobility within the ER
22 lumen (Lai et al., 2010; Lajoie et al., 2012). This finding was demonstrated by a
23 fluorescent recovery after photobleach (FRAP) experiment using a GFP-tagged form
24 of Kar2/BiP (Kar2/BiP-GFP). As anticipated from the contiguous nature of the pnER
25 and cER, fluorescence recovery of Kar2/BiP-GFP was similar throughout the ER. In
26 response to ER stress, the rate of the fluorescence recovery of Kar2/BiP GFP in the
27 pnER was significantly decreased as anticipated. However, the decrease in the
28 fluorescence recovery was asymmetric, differing significantly between the pnER
29 and cER. This result thus revealed differences in Kar2/BiP mobility in the pnER
30 versus cER, which might reflect differences in the level of ER stress (Pina et al.,
31 2016).

Further investigation provided a basis for the use of asymmetric FRAP for identifying ERSU components. In UPR-deficient *ire1Δ* cells, similar magnitudes of FRAP differences between the cER and pnER were also observed. Surprisingly, however, in ERSU-deficient *slt2Δ* cells, the extent of the reduction in the mobility of KAR2/BiP-GFP of the pnER became similar in the cER (Pina et al., 2018). In ER-stressed *slt2Δ* cells, both cER and pnER entered the daughter cell; by contrast, in ER-stressed WT or *ire1Δ* cells, only pnER entered the daughter cell. Further investigation confirmed that the loss of the asymmetric difference between the cER and pnER provides a means to identify ERSU-deficient cells like *slt2Δ*, although the molecular bases of these differences between the cER and pnER were unclear.

Indeed, based on the FRAP behavior, a search for ER components that establish the differential ER chaperone behaviors under ER stress identified *bona fide* ERSU components (Pina et al., 2016). Specifically, *rtn1Δrtn2Δyop1Δ* cells lacking all three reticulon family proteins, RTN1, RTN2, and YOP1, displayed a phenotype with diminished asymmetric behaviors of the KAR2/BiP-GFP in response to ER stress (**Figure 5**). Reticulons are known to play important roles in the generation of ER membrane curvature, and therefore affect the overall architecture of the ER including the cER (Hu et al., 2011; Westrate et al., 2015). Thus, the cER of *rtn1Δrtn2Δyop1Δ* cells was significantly altered compared to that of WT cells and the FRAP behaviors of *rtn1Δrtn2Δyop1Δ* were similar to those of ERSU-deficient *slt2Δ* cells (**Figure 5**). Indeed, ER stress induction of *rtn1Δrtn2Δyop1Δ* cells displayed all of the hallmark phenotypes of ERSU-deficient cells (Pina et al., 2016).

The discovery that the asymmetric differences in the luminal conditions of the cER and pnER and that reticulon family proteins such as Rtn1 and Yop1 are involved in the ERSU pathway suggested that the specific shape of the ER contributes to the ERSU-initiating signals. Although the detailed architectural and functional relationships of the ER are not fully understood, the local distributions or functions of the ER-resident or transmembrane proteins are known to dictate specific ER shapes. For example, an ER sheet-like structure is enriched in the pnER region, whereas an ER tubule structure is concentrated in the cER region (Schwarz


and Blower, 2016; Wang and Rapoport, 2019; Westrate et al., 2015). Furthermore, ribosomes are bound to the surface of the ER sheets, marking the area of the ER that generates secretory pathway proteins and constituting the rough ER. The smooth ER that lacks surface-associated ribosomes consists of the tubular ER and is involved in lipid biosynthesis. The distributions of ER sheets and tubules are regulated by a balance of the activities of two functionally antagonistic ER structural proteins, Lunapark1 (Lnp1) and Sey1/Atlastin (Chen et al., 2013; Goyal and Blackstone, 2013; Wang and Rapoport, 2019). These proteins are associated with three-way ER junctions, impact ER structure upon association with Rtn1, Rtn2, and Yop1 proteins, and act antagonistically (Anwar et al., 2012; Chen et al., 2015; Hu et al., 2008). Loss of the *LNP1* gene causes the Sey1 protein to localize to the cER and pnER and generates a more densely reticulated ER structure in yeast and a more sheet-like ER in mammalian cells (Hu et al., 2009; Shemesh et al., 2014). The loss of *SEY1* and *YOP1* reduces branched tubular ER, resulting in an accumulation of Lnp1 at the cER and pnER. Given such relationships, altering the proportions of the reticular structure vs. ER sheets upon disruption of the *LNP1*, but not *SEY1*, gene in *rtn1Δrtn2Δyop1Δ* cells indeed restored the asymmetry of KAR2/BiP-GFP FRAP behavior between the cER and pnER. Furthermore, the lack of the *LNP1*, but not *SEY1*, gene restored the ERSU-deficient phenotype in *rtn1Δrtn2Δyop1Δ* cells. Finally, recent studies identified mutant forms of either Rtn1 or Yop1, both carrying a single amino acid change, that diminish the PHS-dependent activation of the ERSU cell cycle checkpoint (unpublished results from the Niwa lab), revealing that RTN1, RTN2, and YOP1 respond to ER stress-induced PHS to activate the ERSU events. Yop1 is independent of the role of these proteins in proper ER structure. Importantly, cells carrying a mutant Rtn1 with several amino acid changes that alter the asymmetric FRAP behaviors of KAR2/BiP-GFP (Shibata et al., 2008) was found to be ERSU defective (unpublished results from the Niwa lab), providing further confirmation that RTN1, RTN2, and YOP1 play vital roles in ER inheritance block during ER stress.

Identifying the ERSU activating signal: phytosphingosine

The next important knowledge gap is the identity of the activating signal(s) of the ERSU pathway. Since the UPR sensor Ire1 is not involved in ERSU pathway activation, it is unlikely that signals activating IRE1, for example the accumulation of unfolded proteins, are involved in the ERSU. However, the activating signal(s) still must be induced by ER stress, which can be triggered by well-characterized ER stress-inducing agents, such as Tm or DTT, or by using a temperature-sensitive allele of *ero1-1* that causes unfolded proteins to accumulate.

A number of key findings suggested that sphingolipid levels increase in response to treatment with an ER stress-inducing agent such as Tm in WT yeast cells. The increase is relatively small and transient, but similar changes in sphingolipid levels have been reported during heat shock (Jenkins, 2003; Meier et al., 2006). The early steps of sphingolipid biosynthesis are initiated on the ER membrane; thus, the ER stress-induced increase in their production might act as an ERSU initiating signal. Indeed, exogenous addition of an ER-localized sphingolipid, PHS, to unstressed cells activates all of the ERSU cell cycle checkpoint hallmark events including (1) cER inheritance, (2) septin ring mislocalization, (3) Slt2 phosphorylation, and (4) cytokinesis arrest. The involvement of PHS is specific, as other sphingolipids/ceramides, such as dihydroceramide (DHC) and ceramide, do not robustly activate the ERSU pathway (**Figure 6A**).

Subsequent experiments using pharmacological drugs such as myriocin or aureobasidin A (AbA) also provided confirmation of the role of PHS as an activating signal (Pina et al., 2018). For example, treatment of cells with AbA, which blocks the conversion of PHS to further downstream products, an increase in the levels of PHS and activated the ERSU pathway in the absence of ER stress-inducing agents. Treating cells with an ER stress-inducing agent, such as Tm or DTT, further increased the extent of ERSU pathway activation. Additionally, treating cells with myriocin, which blocks serine palmitoyltransferase (SPT), diminished the level of ERSU pathway activation. SPT generates 3-




ketohydrosphingosine from serine and palm
 biosynthetic pathway, treating cells with m
 sphingolipids including PHS (**Figure 6A**).

The involvement of PHS in ERSU pathw
 genetic experiments in yeast (Sauldiner et al., 2000; Saito and Weissman, 2013). For example, an epistatic miniarray profile study identified a series of genes in the sphingolipid biosynthetic pathway as genetic interactors of the ERSU component Slt2. For example, ORM1 and ORM2 encode subunits of an inhibitory complex of SPT; cells lacking ORM1 and ORM2 (orm1Δorm2Δ double knockout cells) have elevated levels of cellular sphingolipids (Preslow et al., 2010; Han et al., 2010). Upon treatment with Tm, orm1Δorm2Δ biosynthetic knockout cells further increased the frequency of ERSU events. Importantly, adding PHS exogenously did not activate IRE1 or the UPR pathway.

Interestingly, recent studies also reported the involvement of sphingolipids in the establishment of the nuclear envelope and the ER diffusion barrier (Clay et al., 2014; Megyeri et al., 2019). While multiple sphingolipids and ceramides have been reported to function for the ER diffusion barrier, the involvement of PHS in the ERSU checkpoint is specific, as DHC does not support the maintenance of the ERSU. Furthermore, recent preliminary work has revealed that PHS functions in the ERSU checkpoint independent of changes in the ER diffusion barrier (unpublished results in Niwa lab). Taken together, these observations are consistent with the idea that elevated PHS in response to ER stress acts as an activating signal for the ERSU pathway, leading to ER inheritance block, septin ring mislocalization, Slt2 phosphorylation, and cytokinesis block.

Sphingolipids and ER homeostasis in mammalian cells

The sphingolipid biosynthetic pathway is conserved in eukaryotic cells (Gault et al., 2010; Riezman, 2006), although some details differ depending upon the



st step of the
 llular levels of

o supported by

Figure 6: Sphingolipid biosynthetic pathways.

A. An abbreviated yeast sphingolipid biosynthetic pathway.
 Serine + Palmitoyl-CoA → SPT (Lcb2, Lcb1, Tsc3) → 3-Ketodihydrosphingosine → TSC10 → Dihydrosphingosine (DHS) → SUR2 → Phytosphingosine (PHS) → Phytoceramide.
B. A comparison of the human and yeast sphingolipid biosynthetic pathways.
 Serine + Palmitoyl-CoA → SPT (Lcb2, Lcb1, Tsc3) → 3-Ketodihydrosphingosine → TSC10 → Dihydrosphingosine (DHS) → Dihydroceramide synthase → Dihydroceramide (DHC) → Ceramide → Sphingomyelin.

(PHS) is highlighted. Treatment of cells with myriocin reduces PHS levels, while orm1Δorm2Δ cells or Aureobasidin A (ABA) treatment of cells increases the PHS levels.

B. A comparison of the human and yeast sphingolipid biosynthetic pathways.
 Serine + Palmitoyl-CoA → SPT (Lcb2, Lcb1, Tsc3) → 3-Ketodihydrosphingosine → TSC10 → Dihydrosphingosine (DHS) → Dihydroceramide synthase → Dihydroceramide (DHC) → Ceramide → Sphingomyelin.

specific species. Interestingly, a study found that ER stress induction also increases dihydrosphingosine (DHS) and DHC levels in mammalian cells (Tam et al., 2018). Like PHS in yeast, both DHS and DHC are early intermediates of the sphingolipid biosynthetic pathway in mammalian cells (**Figure 6B**) (Riezman, 2006). The increased levels of these sphingolipids are transient. The molecular mechanisms of the UPR pathway are more complex in mammalian cells than those in yeast cells: in yeast, IRE1 is the only ER transmembrane UPR signaling component, whereas the mammalian UPR pathway is activated by two additional ER transmembrane components, PERK and ATF6 (Schroder and Kaufman, 2005). While the mammalian ERSU checkpoint has yet to be defined, the additional mammalian UPR components (i.e., ATF6 and PERK) might provide means to respond to ER stress-induced DHS and DHC. Exogenously added DHS or DHC indeed activated ATF6 α , but did not affect PERK or IRE1 (Tam et al., 2018). Further studies revealed a unique DHS/DHC binding motif within the transmembrane domain of ATF6 α . A single amino acid substitution within this motif resulted in inactivation of ATF6 α by DHS and DHC. However, these ATF6 α mutants were still activated by increased levels of unfolded proteins in the ER. Conversely, mutations within the ER luminal domain of ATF6 α failed to respond to the increased levels of unfolded proteins but remained active to respond to exogenously added DHS and DHC. These results indicate that ATF6 α has two distinct stress-sensing domains, one through the ER luminal domain and the other via the transmembrane motif. Currently, it remains to be determined if the mammalian ERSU checkpoint exists and how such responses are wired, but it is possible that ATF6 α is a dual component acting on the UPR pathway and the ERSU cell cycle checkpoint for the ER in mammalian cells by two distinct domains. Furthermore, the identification of the ATF6 α DHS/DHC binding motif provides hints about the activation mechanisms of ERSU components, such as Rtn1 or Yop1, via PHS.

Many more exciting questions about the yeast ERSU cell cycle checkpoint

The discovery of the ERSU pathway has led to many interesting questions. For example, why is PHS the activating signal, instead of the more abundant sphingolipids or ceramides? Production of sphingolipids and ceramide begins at the ER and synthesis of PHS occurs at the ER (Riezman, 2006). Once converted, ceramide is transported to the Golgi where further downstream lipid biosynthetic steps continue (Funato and Riezman, 2001; Liu et al., 2017). Finally, some of these lipids reach the plasma membrane. For example, glycosphingolipids and cholesterol (ergosterol in yeast) generate a subdomain within the plasma membrane, the lipid rafts (Hurst and Fratti, 2020; Lingwood and Simons, 2010). The use of more abundant lipids as signals might require more major cellular changes before the ERSU pathway is turned on. For the ERSU pathway to work as a temporary halt of the cell cycle during ER stress, such dramatic changes might not be suitable.

How are PHS levels increased during ER stress? How are increased levels of PHS recognized? Do Rtn1, Rtn2 and Yop1 proteins somehow measure the PHS levels? Or, do alternative components exist, ultimately leading to ER inheritance block, septin ring mislocalization and Slt2 phosphorylation via Rtn1, Rtn2, and Yop1? Do PHS levels contribute to the establishment of the asymmetry between the cER and the pnER? Interestingly, a mutagenesis study of Rtn1 reported that certain mutations within the Rtn1 protein caused the FRAP profile of BiP/Kar2-GFP in the cER to differ from that in the pnER, even in the absence of ER stress induction (Shibata et al., 2008), revealing that Rtn1 is sensitive to ER functional homeostasis.

ER stress induces both the ERSU and UPR pathways. While UPR activation re-establishes ER homeostasis, the ERSU pathway halts both the cell cycle and ER inheritance. In order for cells to re-enter the cell cycle in a timely manner, certain components such as those that re-evaluate ER functional status during cell cycle block and those that establish the release from cytokinesis block, must be involved. Furthermore, after release from cytokinesis block, how do cells re-enter the cell cycle? If ER functional homeostasis does not recover within a certain period

of time, what happens to the cell? Are there time limits for cell cycle recovery? Answering these questions will be critical to fully understanding the yeast ERSU cell cycle checkpoint, a mechanism to ensure all dividing cells will have sufficient levels of the ER.

Potential for cell cycle regulation of the ER in mammalian cells

Given the complexity of the mammalian ER network, examining the effect of ER stress on the ER distribution in dividing cells is challenging. During prophase, the nuclear membranes detach from the lamins and chromatin to disassemble. In fact, the nuclear membrane components are retracted into and become dispersed throughout the ER (Anderson et al., 2009). During the metaphase-to-anaphase transition, however, the nuclear envelope (NE) re-forms from ER tubules and sheets to enclose each set of daughter chromosomes (Anderson and Hetzer, 2008). Thus, the ER is intimately involved in NE disassembly and reassembly. Furthermore, during NE disassembly, nuclear membrane proteins such as POM121, an essential transmembrane protein of the nuclear pore, have to become dispersed throughout the ER and relocate to the newly reassembled NE and nuclear pores as the NE reassembles. Similarly, studies have reported that RTN3 first is associated with chromatin-bound membranes, but becomes increasingly dispersed throughout the surrounding ER by anaphase (Anderson and Hetzer, 2008). Thus, such processes might be impacted by ER stress. Indeed, induction of ER stress resulted in the mislocalization of septin ring subunits (unpublished results from the Niwa lab). Understanding the mechanisms by which the ER is divided in normal cells and how this is perturbed under various stress conditions will contribute to our understanding of human disease. Indeed, dysregulated ER function is a prominent feature of many disorders, including diabetes, Alzheimer's disease, and Parkinson's disease, which are increasing public health concerns. Thus, studies on ER function may ultimately lead to the development of new treatments for such diseases.

Summary

The discovery of the ERSU cell cycle checkpoint pathway unveiled the exciting possibility that the division of cytoplasmic components such as organelles or protein complexes is regulated in order to generate fully functional eukaryotic cells. Further studies in this area may unveil new strategies for designing novel drugs for human diseases that are caused by improper cell cycle regulation.

Acknowledgements

The author apologizes for absence of citations within the space limit of this review. The author thanks members of the laboratory for critical reading of the manuscript. The author thanks members of the Niwa lab for careful reading of the review. The work is supported by NIH (RO1 GM087415) and Cancer Research Coordinating Committee fund (CRR-20-632388).

References

- Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., and Walter, P. (2002). Molecular Biology of the Cell, 4th edition. Molecular Biology of the Cell, 4th edition.
- Anderson, D.J., and Hetzer, M.W. (2008). Reshaping of the endoplasmic reticulum limits the rate for nuclear envelope formation. *J Cell Biol* 182, 911-924.
- Anderson, D.J., Vargas, J.D., Hsiang, P., and Hetzer, M.W. (2009). Recruitment of functionally distinct membrane proteins to chromatin mediates nuclear envelope formation in vivo. *J Cell Biol* 186, 183-191.
- Arnold, E., and Tanner, W. (1982). An obligatory role of protein glycosylation in the life cycle of yeast cells. *FEBS Lett* 148, 49-53.
- Aviram, N., and Schuldiner, M. (2007). Targeting and translocation of proteins to the endoplasmic reticulum at a glance. *J Cell Sci* 130, 4079-4085.
- Babour, A., Bicknell, A.A., Tourtelotte, J., and Niwa, M. (2010). A surveillance pathway monitors the fitness of the endoplasmic reticulum to control its inheritance. *Cell* 142, 256-269.
- Back, S.H., and Kaufman, R.J. (2010). Endoplasmic reticulum stress and type 2 diabetes. *Annu Rev Biochem* 81, 767-793.
- Berner, N., Reutter, K.R., and Wolf, D.H. (2018). Protein Quality Control of the Endoplasmic Reticulum and Ubiquitin-Proteasome-Triggered Degradation of Aberrant Proteins: Yeast Pioneers the Path. *Annu Rev Biochem* 87, 751-782.
- Bertolotti, A., Zhang, Y.H., Henderson, L.M., Harding, H.P., and Ron, D. (2000). Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol* 2, 326-332.
- Bicknell, A.A., Babour, A., Federovich, C.M., and Niwa, M. (2007). A novel role in cytokinesis reveals a housekeeping function for the unfolded protein response. *J Cell Biol* 177, 1017-1027.
- Bidlingmaier, S., and Snyder, M. (2004). Regulation of polarized growth initiation and termination cycles by the polarisome and Cdc42 regulators. *J Cell Biol* 164, 207-218.
- Bonilla, M., and Cunningham, K.W. (2003). Mitogen-activated protein kinase stimulation of Ca(2+) signaling is required for survival of endoplasmic reticulum stress in yeast. *Mol Biol Cell* 14, 4296-4305.
- Breslow, D.K., Collins, S.R., Bodenmiller, B., Aebersold, R., Simons, K., Shevchenko, A., Ejning, C.S., and Weissman, J.S. (2010). Orm family proteins mediate sphingolipid homeostasis. *Nature* 463, 1048-1053.

Brown, J.D., Ng, D.T., Ogg, S.C., and Walter, P. (1995). Targeting pathways to the
 endoplasmic reticulum membrane. Cold Spring Harb Symp Quant Biol 60, 23-30.

Caudron, F., and Barral, Y. (2009). Septins and the lateral compartmentalization of
 eukaryotic membranes. Dev Cell 16, 493-506.

Chao, J.T., Pina, F., Onishi, M., Cohen, Y., Lai, Y.S., Schuldiner, M., and Niwa, M.
 (2019). Transfer of the Septin Ring to Cytokinetic Remnants in ER Stress Directs
 Age-Sensitive Cell-Cycle Re-entry. Dev Cell 51, 173-191 e175.

Chao, J.T., Wong, A.K., Tavassoli, S., Young, B.P., Chruscicki, A., Fang, N.N., Howe,
 L.J.,
 Mayor, T., Foster, L.J., and Loewen, C.J. (2014). Polarization of the endoplasmic
 reticulum by ER-septin tethering. Cell 158, 620-632.

Chen, S., Novick, P., and Ferro-Novick, S. (2013). ER structure and function. Curr
 Opin Cell Biol 25, 428-433.

Ciccia, A., and Elledge, S.J. (2010). The DNA damage response: making it safe to
 play with knives. Mol Cell 40, 179-204.

Clay, L., Caudron, F., Denoth-Lippman, A., Boettcher, B., Buvelot Frei, S., Snapp,
 E.L., and Barral, Y. (2014). A sphingolipid-dependent diffusion barrier confines ER
 stress to the yeast mother cell. ELife 3, e01883.

Cohen, S., Valm, A.M., and Lippman-Schwartz, J. (2018). Interacting organelles.
 Curr Opin Cell Biol 53, 84-91.

Cox, J.S., Shamu, C.E., and Walter, P. (1993). Transcriptional induction of genes
 encoding endoplasmic reticulum resident proteins requires a transmembrane
 protein kinase. Cell 73, 1197-1206.

Cox, J.S., Sidrauski, C., Shamu, C., and Walter, P. (1996). Regulation of the unfolded
 protein response pathway. Molecular Biology of the Cell 7, 2944-2944.

Credle, J.J., Finer-Moore, J.S., Papas, F.R., Stroud, R.M., and Walter, P. (2005). On the
 mechanism of sensing unfolded protein in the endoplasmic reticulum. Proc Natl
 Acad Sci U S A 102, 18773-18784.

Cullinan, S.B., and Diehl, J.A. (2006). Coordination of ER and oxidative stress
 signaling: the PERK/Nrf2 signaling pathway. Int J Biochem Cell Biol 38, 317-332.

Doyle, T., and Botstein, D. (1996). Movement of yeast cortical actin cytoskeleton
 visualized in vivo. Proc Natl Acad Sci U S A 93, 3886-3891.

Du, Y., Pypaert, M., Novick, P., and Ferro-Novick, S. (2001). Aux1p/Swa2p is
 required for cortical endoplasmic reticulum inheritance in Saccharomyces
 cerevisiae. Molecular Biology of the Cell 12, 2614-2628.

Du, Y., Walker, L., Novick, P., and Ferro-Novick, S. (2006). Ptc1p regulates cortical ER inheritance via Slt2p. *EMBO J* 25, 4413-4422.

Elledge, S.J., and Harper, J.W. (1994). Cdk inhibitors: on the threshold of checkpoints and development. *Curr Opin Cell Biol* 6, 847-852.

Estrada de Martin, P., Novick, P., and Ferro-Novick, S. (2005). The organization, structure, and inheritance of the ER in higher and lower eukaryotes. *Biochem Cell Biol* 83, 752-761.

Estrada, P., Kim, J., Coleman, J., Walker, L., Dunn, B., Takizawa, P., Novick, P., and Ferro-Novick, S. (2003). Myo4p and She3p are required for cortical ER inheritance in *Saccharomyces cerevisiae*. *J Cell Biol* 163, 1255-1266.

Farber-Katz, S.E., Dippold, H.C., Buschman, M.D., Peterman, M.C., Xing, M., Noakes, C.J., Tat, J., Ng, M.M., Rahajeng, J., Cowan, D.M., et al. (2014). DNA damage triggers Golgi dispersal via DNA-PK and GOLPH3. *Cell* 156, 413-427.

Farre, J.C., Mahalingam, S.S., Proietto, M., and Subramani, S. (2019). Peroxisome biogenesis, membrane contact sites, and quality control. *EMBO Rep* 20.

Field, C.M., and Kellogg, D. (1999). Septins: cytoskeletal polymers or signalling GTPases? *Trends Cell Biol* 9, 387-394.

Forbes, D.J., Travesa, A., Nord, M.S., and Bernis, C. (2015). Nuclear transport factors: global regulation of mitosis. *Curr Opin Cell Biol* 35, 78-90.

Funato, K., and Riezman, H. (2001). Vesicular and nonvesicular transport of ceramide from ER to the Golgi apparatus in yeast. *J Cell Biol* 155, 949-959.

Gault, C.R., Obeid, L.M., and Hannun, Y.A. (2010). An overview of sphingolipid metabolism: from synthesis to breakdown. *Adv Exp Med Biol* 688, 1-23.

Goder, V., Alanis-Dominguez, E., and Bustamante-Sequeiros, M. (2019). Lipids and their (un)known effects on ER-associated protein degradation (ERAD). *Biochim Biophys Acta Mol Cell Biol Lipids*.

Goyal, U., and Blackstone, C. (2013). Untangling the web: mechanisms underlying ER network formation. *Biochim Biophys Acta* 1833, 2492-2498.

Hampton, R.Y., Gardner, R.G., and Rine, J. (1996). Role of 26S proteasome and HRD genes in the degradation of 3-hydroxy-3-methylglutaryl-CoA reductase, an integral endoplasmic reticulum membrane protein. *Molecular Biology of the Cell* 7, 2029-2044.

- 1 Hampton, R.Y., and Sommer, T. (2012). Finding the will and the way of ERAD
2 substrate retrotranslocation. *Curr Opin Cell Biol* 24, 460-466.
- 3
- 4 Han, S., Lone, M.A., Schneider, R., and Chang, A. (2010). Orm1 and Orm2 are
5 conserved endoplasmic reticulum membrane proteins regulating lipid homeostasis
6 and protein quality control. *Proc Natl Acad Sci U S A* 107, 5851-5856.
- 7
- 8 Heider, M.R., and Munson, M. (2012). Exorcising the exocyst complex. *Traffic* 13,
9 898-907.
- 10
- 11 Helle, S.C., Kanfer, G., Kolar, K., Lang, A., Michel, A.H., and Kornmann, B. (2013).
12 Organization and function of membrane contact sites. *Biochim Biophys Acta* 1833,
13 2526-2541.
- 14
- 15 Henry, K.A., Blank, H.M., Hoose, S.A., and Polymenis, M. (2010). The unfolded
16 protein response is not necessary for the G1/S transition, but it is required for
17 chromosome maintenance in *Saccharomyces cerevisiae*. *PLoS One* 5, e12732.
- 18
- 19 Holland, A.J., and Cleveland, D.W. (2012). Losing balance: the origin and impact of
20 aneuploidy in cancer. *EMBO Rep* 13, 501-514.
- 21
- 22 Hu, J., Prinz, W.A., and Rapoport, T.A. (2011). Weaving the web of ER tubules. *Cell*
23 147, 1226-1231.
- 24
- 25 Hu, J., Shibata, Y., Zhu, P.P., Voss, C., Rismanchi, N., Prinz, W.A., Rapoport, T.A.,
26 and Blackstone, C. (2009). A class of dynamin-like GTPases involved in the
27 generation of the tubular ER network. *Cell* 138, 549-561.
- 28
- 29 Hurst, L.R., and Fratti, R.A. (2020). Lipid Rafts, Sphingolipids, and Ergosterol in
30 Yeast Vacuole Fusion and Maturation. *Front Cell Dev Biol* 8, 539.
- 31
- 32 Jenkins, G.M. (2003). The emerging role for sphingolipids in the eukaryotic heat
33 shock response. *Cell Mol Life Sci* 60, 701-710.
- 34
- 35 Jimenez-Gutierrez, E., Alegria-Carrasco, E., Sellers-Moya, A., Molina, M., and Martin,
36 H. (2020). Not just the wall: the other ways to turn the yeast CWI pathway on. *Int*
37 *Microbiol* 23, 107-119.
- 38
- 39 Jin, Y., and Weisman, L.S. (2015). The vacuole/lysosome is required for cell-cycle
40 progression. *Elife* 4.
- 41
- 42 Johnson, B.M., and DeBose-Boyd, R.A. (2018). Underlying mechanisms for sterol-
43 induced ubiquitination and ER-associated degradation of HMG CoA reductase.
44 *Semin Cell Dev Biol* 81, 121-128.
- 45
- 46 Karagas, N.E., and Venkatachalam, K. (2019). Roles for the Endoplasmic Reticulum
47 in Regulation of Neuronal Calcium Homeostasis. *Cells* 8.
- 48

1 Kilmartin, J.V., and Adams, A.E. (1984). Structural rearrangements of tubulin and
2 actin during the cell cycle of the yeast *Saccharomyces*. *J Cell Biol* 98, 922-933.
3
4 Kimata, Y., Ishiwata-Kimata, Y., Ito, T., Hirata, A., Suzuki, T., Oikawa, D., Takeuchi,
5 M., and
6 Kohno, K. (2007). Two regulatory steps of ER-stress sensor Ire1 involving its cluster
7 formation and interaction with unfolded proteins. *J Cell Biol* 179, 75-86.
8
9 Koning, A.J., Larson, L.L., Cadera, E.J., Parrish, M.L., and Wright, R.L. (2002).
10 Mutations that affect vacuole biogenesis inhibit proliferation of the endoplasmic
11 reticulum in *Saccharomyces cerevisiae*. *Genetics* 160, 1335-1352.
12
13 Kvam, E., and Goldfarb, D.S. (2006). Nucleus-vacuole junctions in yeast: anatomy
14 of a membrane contact site. *Biochem Soc Trans* 34, 340-342.
15
16 Lai, C.W., Aronson, D.E., and Snapp, E.L. (2010). BiP availability distinguishes
17 states of homeostasis and stress in the endoplasmic reticulum of living cells. *Mol*
18 *Biol Cell* 21, 1909-1921.
19
20 Lajoie, P., Moir, R.D., Willis, I.M., and Snapp, E.L. (2012). Kar2p availability defines
21 distinct forms of endoplasmic reticulum stress in living cells. *Mol Biol Cell* 23, 955-
22 964.
23
24 Levin, D.E., and Errede, B. (1995). The proliferation of MAP kinase signaling
25 pathways in yeast. *Curr Opin Cell Biol* 7, 197-202.
26
27 Li, X., Ferro-Novick, S., and Novick, P. (2013). Different polarisome components
28 play distinct roles in Slt2p-regulated cortical ER inheritance in *Saccharomyces*
29 *cerevisiae*. *Mol Biol Cell* 24, 3145-3154.
30
31 Lingwood, D., and Simons, K. (2010). Lipid rafts as a membrane-organizing
32 principle. *Science* 327, 46-50.
33
34 Liu, L.K., Choudhary, V., Toulmay, A., and Prinz, W.A. (2017). An inducible ER-Golgi
35 tether facilitates ceramide transport to alleviate lipotoxicity. *J Cell Biol* 216, 131-
36 147.
37
38 Liu, Q., Guntuku, S., Cui, X.S., Matsuoka, S., Cortez, D., Tamai, K., Luo, G., Carattini-
39 Rivera, S., DeMayo, F., Bradley, A., *et al.* (2000). Chk1 is an essential kinase that is
40 regulated by Atr and required for the G(2)/M DNA damage checkpoint. *Genes Dev*
41 14, 1448-1459.
42
43 Ma, Y., and Hendershot, L.M. (2001). The unfolding tale of the unfolded protein
44 response. *Cell* 107, 827-830.
45
46 Mascanzoni, F., Ayala, I., and Colanzi, A. (2019). Organelle Inheritance Control of
47 Mitotic Entry and Progression: Implications for Tissue Homeostasis and Disease.
48 *Front Cell Dev Biol* 7, 133.
49

- 1 Matellan, L., and Monje-Casas, F. (2020). Regulation of Mitotic Exit by Cell Cycle
2 Checkpoints: Lessons From *Saccharomyces cerevisiae*. *Genes (Basel)* *11*.
- 3
- 4 Matlack, K.E., Mothes, W., and Rapoport, T.A. (1998). Protein translocation: tunnel
5 vision. *Cell* *92*, 381-390.
- 6
- 7 McCaffrey, K., and Braakman, I. (2016). Protein quality control at the endoplasmic
8 reticulum. *Essays Biochem* *60*, 227-235.
- 9
- 10 McMaster, C.R. (2001). Lipid metabolism and vesicle trafficking: more than just
11 greasing the transport machinery. *Biochem Cell Biol* *79*, 681-692.
- 12
- 13 McMillan, J.N., Sia, R.A., and Lew, D.J. (1998). A morphogenesis checkpoint
14 monitors the actin cytoskeleton in yeast. *J Cell Biol* *142*, 1487-1499.
- 15
- 16
- 17 Megyeri, M., Prasad, R., Volpert, G., Sliwa-Gonzalez, A., Haribowo, A.G., Aguilera-
18 Romero, A.,
19 Riezman, H., Barral, Y., Futerman, A.H., and Schuldiner, M. (2019). Yeast ceramide
20 synthases, Lag1 and Lac1, have distinct substrate specificity. *J Cell Sci* *132*.
- 21
- 22 Mehrtash, A.B., and Hochstrasser, M. (2019). Ubiquitin-dependent protein
23 degradation at the endoplasmic reticulum and nuclear envelope. *Semin Cell Dev*
24 *Biol* *93*, 111-124.
- 25
- 26 Meier, K.D., Deloche, O., Kajiwar, K., Funato, K., and Riezman, H. (2006).
27 Sphingoid base is required for translation initiation during heat stress in
28 *Saccharomyces cerevisiae*. *Mol Biol Cell* *17*, 1164-1175.
- 29
- 30 Meldolesi, J., and Pozzan, T. (1998). The endoplasmic reticulum Ca²⁺ store: a view
31 from the lumen. *Trends Biochem Sci* *23*, 10-14.
- 32
- 33 Mittal, B., Tulsyan, S., Kumar, S., Mittal, R.D., and Agarwal, G. (2015). Cytochrome
34 P450 in Cancer Susceptibility and Treatment. *Adv Clin Chem* *71*, 77-139.
- 35
- 36 Morgan, D.O. (1995). Principles of CDK regulation. *Nature* *374*, 131-134.
- 37
- 38 Mori, K. (2009). Signalling pathways in the unfolded protein response: development
39 from yeast to mammals. *J Biochem* *146*, 743-750.
- 40
- 41 Mori, K., Ma, W., Gething, M.J., and Sambrook, J. (1993). A transmembrane protein
42 with a cdc2+/CDC28-related kinase activity is required for signaling from the ER to
43 the nucleus. *Cell* *74*, 743-756.
- 44
- 45 Mori, K., Ogawa, N., Kawahara, T., Yanagi, H., and Yura, T. (2000). mRNA splicing-
46 mediated C-terminal replacement of transcription factor Hac1p is required for
47 efficient activation of the unfolded protein response. *Proc Natl Acad Sci U S A* *97*,
48 4660-4665.
- 49

1 Mulholland, J., Preuss, D., Moon, A., Wong, A., Drubin, D., and Botstein, D. (1994).
 2 Ultrastructure of the yeast actin cytoskeleton and its association with the plasma
 3 membrane. *J Cell Biol* 125, 381-391.
 4
 5 Munson, M., and Novick, P. (2006). The exocyst defrocked, a framework of rods
 6 revealed. *Nat Struct Mol Biol* 13, 577-581.
 7
 8 Musacchio, A., and Desai, A. (2017). A Molecular View of Kinetochore Assembly and
 9 Function. *Biology (Basel)* 6.
 10
 11 Novick, P., and Botstein, D. (1985). Phenotypic analysis of temperature-sensitive
 12 yeast actin mutants. *Cell* 40, 405-416.
 13
 14 Oikawa, D., Kimata, Y., and Kohno, K. (2007). Self-association and BiP dissociation
 15 are not sufficient for activation of the ER stress sensor Ire1. *J Cell Sci* 120, 1681-
 16 1688.
 17
 18 Pierro, C., Sneyers, F., Bultynck, G., and Roderick, H.L. (2019). ER Ca(2+) release
 19 and store-operated Ca(2+) entry - partners in crime or independent actors in
 20 oncogenic transformation? *Cell Calcium* 82, 102061.
 21
 22 Pina, F., Yagisawa, F., Obara, K., Gregerson, J.D., Kihara, A., and Niwa, M. (2018).
 23 Sphingolipids activate the endoplasmic reticulum stress surveillance pathway. *J*
 24 *Cell Biol* 217, 495-505.
 25
 26 Pina, F.J., Fleming, T., Pogliano, K., and Niwa, M. (2016). Reticulons Regulate the ER
 27 Inheritance Block during ER Stress. *Dev Cell* 37, 279-288.
 28
 29 Pina, F.J., and Niwa, M. (2015). The ER Stress Surveillance (ERSU) pathway
 30 regulates daughter cell ER protein aggregate inheritance. *Elife* 4.
 31
 32 Pobre, K.F.R., Poet, G.J., and Hendershot, L.M. (2019). The endoplasmic reticulum
 33 (ER) chaperone BiP is a master regulator of ER functions: Getting by with a little
 34 help from ERdj friends. *J Biol Chem* 294, 2098-2108.
 35
 36 Preuss, D., Mulholland, J., Kaiser, C.A., Orlean, P., Albright, C., Rose, M.D., Robbins,
 37 P.W., and Botstein, D. (1991). Structure of the yeast endoplasmic reticulum:
 38 localization of ER proteins using immunofluorescence and immunoelectron
 39 microscopy. *Yeast* 7, 891-911.
 40
 41 Riezman, H. (2006). Organization and functions of sphingolipid biosynthesis in
 42 yeast. *Biochem Soc Trans* 34, 367-369.
 43
 44 Ron, D., and Walter, P. (2007). Signal integration in the endoplasmic reticulum
 45 unfolded protein response. *Nat Rev Mol Cell Biol* 8, 519-529.
 46
 47 Rutkowski, D.T., and Kaufman, R.J. (2004). A trip to the ER: coping with stress.
 48 *Trends Cell Biol* 14, 20-28.
 49

Schroder, M., and Kaufman, R.J. (2005). The mammalian unfolded protein response. *Annu Rev Biochem* 74, 739-789.

Schuldiner, M., Collins, S.R., Thompson, N.J., Denic, V., Bhamidipati, A., Punna, T., Ihmels, J., Andrews, B., Boone, C., Greenblatt, J.F., *et al.* (2005). Exploration of the function and organization of the yeast early secretory pathway through an epistatic miniarray profile. *Cell* 123, 507-519.

Schuldiner, M., and Weissman, J.S. (2013). The contribution of systematic approaches to characterizing the proteins and functions of the endoplasmic reticulum. *Cold Spring Harb Perspect Biol* 5, a013284.

Schwartz, M., Travesa, A., Martell, S.W., and Forbes, D.J. (2015). Analysis of the initiation of nuclear pore assembly by ectopically targeting nucleoporins to chromatin. *Nucleus* 6, 40-54.

Schwarz, D.S., and Blower, M.D. (2016). The endoplasmic reticulum: structure, function and response to cellular signaling. *Cell Mol Life Sci* 73, 79-94.

Shamu, C.E., and Walter, P. (1996). Oligomerization and phosphorylation of the Ire1p kinase during intracellular signaling from the endoplasmic reticulum to the nucleus. *EMBO J* 15, 3028-3039.

Shemesh, T., Klemm, R.W., Romano, F.B., Wang, S., Vaughan, J., Zhuang, X., Tukachinsky, H., Kozlov, M.M., and Rapoport, T.A. (2014). A model for the generation and interconversion of ER morphologies. *Proc Natl Acad Sci U S A* 111, E5243-5251.

Shibata, Y., Voss, C., Rist, J.M., Hu, J., Rapoport, T.A., Prinz, W.A., and Voeltz, G.K. (2008). The reticulon and DP1/Yop1p proteins form immobile oligomers in the tubular endoplasmic reticulum. *J Biol Chem* 283, 18892-18904.

Sidrauski, C., Cox, J.S., and Walter, P. (1996). tRNA ligase is required for regulated mRNA splicing in the unfolded protein response. *Cell* 87, 405-413.

Sun, Z., and Brodsky, J.L. (2019). Protein quality control in the secretory pathway. *J Cell Biol* 218, 3171-3187.

Tam, A.B., Roberts, L.S., Chandra, V., Rivera, I.G., Nomura, D.K., Forbes, D.J., and Niwa, M. (2018). The UPR Activator ATF6 Responds to Proteotoxic and Lipotoxic Stress by Distinct Mechanisms. *Dev Cell* 46, 327-343 e327.

Tamura, Y., Kawano, S., and Endo, T. (2019). Organelle contact zones as sites for lipid transfer. *J Biochem* 165, 115-123.

Vai, M., Popolo, L., and Alberghina, L. (1987). Effect of tunicamycin on cell cycle progression in budding yeast. *Exp Cell Res* 171, 448-459.

1 Versele, M., and Thorner, J. (2005). Some assembly required: yeast septins provide
2 the instruction manual. *Trends Cell Biol* 15, 414-424.

3

4 Vevea, J.D., Swayne, T.C., Boldogh, I.R., and Pon, L.A. (2014). Inheritance of the
5 fittest mitochondria in yeast. *Trends Cell Biol* 24, 53-60.

6

7 Voeltz, G.K., Rolls, M.M., and Rapoport, T.A. (2002). Structural organization of the
8 endoplasmic reticulum. *EMBO Rep* 3, 944-950.

9

10 Waddle, J.A., Karpova, T.S., Waterston, R.H., and Cooper, J.A. (1996). Movement of
11 cortical actin patches in yeast. *J Cell Biol* 132, 861-870.

12

13 Walter, P., Gilmore, R., and Blobel, G. (1984). Protein translocation across the
14 endoplasmic reticulum. *Cell* 38, 5-8.

15

16 Walter, P., and Ron, D. (2011). The unfolded protein response: from stress pathway
17 to homeostatic regulation. *Science* 334, 1081-1086.

18

19 Wang, N., and Rapoport, T.A. (2019). Reconstituting the reticular ER network -
20 mechanistic implications and open questions. *J Cell Sci* 132.

21

22 Weisman, L.S. (2003). Yeast vacuole inheritance and dynamics. *Annu Rev Genet*
23 37, 435-460.

24

25 Westrate, L.M., Lee, J.E., Prinz, W.A., and Voeltz, G.K. (2015). Form follows function:
26 the importance of endoplasmic reticulum shape. *Annu Rev Biochem* 84, 791-811.

27

28 Wu, H., Carvalho, P., and Voeltz, G.K. (2018). Here, there, and everywhere: The
29 importance of ER membrane contact sites. *Science* 361.

30

31 Wu, X., and Rapoport, T.A. (2018). Mechanistic insights into ER-associated protein
32 degradation. *Curr Opin Cell Biol* 53, 22-28.

33

34 Zhou, B.B., and Elledge, S.J. (2000). The DNA damage response: putting
35 checkpoints in perspective. *Nature* 408, 433-439.

36

37 Zhou, J., Liu, C.Y., Back, S.H., Clark, R.L., Peisach, D., Xu, Z., and Kaufman, R.J.
38 (2006). The crystal structure of human IRE1 luminal domain reveals a conserved
39 dimerization interface required for activation of the unfolded protein response. *Proc*
40 *Natl Acad Sci U S A* 103, 14343-14348.