

# UC San Diego

## UC San Diego Previously Published Works

### Title

Characterizing the ionization potential depression in dense carbon plasmas with high-precision spectrally resolved x-ray scattering

### Permalink

<https://escholarship.org/uc/item/99h7t9fz>

### Journal

Plasma Physics and Controlled Fusion, 61(1)

### ISSN

0741-3335 1361-6587

### Authors

Kraus, D  
Bachmann, B  
Barbrel, B  
[et al.](#)

### Publication Date

2019

### DOI

10.1088/1361-6587/aadd6c

Peer reviewed

PAPER

## Characterizing the ionization potential depression in dense carbon plasmas with high-precision spectrally resolved x-ray scattering

To cite this article: D Kraus *et al* 2019 *Plasma Phys. Control. Fusion* **61** 014015

View the [article online](#) for updates and enhancements.




**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# Characterizing the ionization potential depression in dense carbon plasmas with high-precision spectrally resolved x-ray scattering

D Kraus<sup>1,2</sup> , B Bachmann<sup>3</sup>, B Barbrel<sup>4</sup>, R W Falcone<sup>4</sup>, L B Fletcher<sup>5</sup>, S Frydrych<sup>6</sup>, E J Gamboa<sup>5</sup>, M Gauthier<sup>5</sup>, D O Gericke<sup>7</sup>, S H Glenzer<sup>5</sup>, S Göde<sup>5</sup>, E Granados<sup>5</sup>, N J Hartley<sup>1,8</sup>, J Helfrich<sup>6</sup>, H J Lee<sup>5</sup>, B Nagler<sup>5</sup>, A Ravasio<sup>9</sup>, W Schumaker<sup>5</sup>, J Vorberger<sup>1</sup> and T Döppner<sup>3</sup>

<sup>1</sup> Helmholtz-Zentrum Dresden-Rossendorf, Bautzner Landstr. 400, D-01328 Dresden, Germany

<sup>2</sup> Institute of Solid State and Materials Physics, Technische Universität Dresden, D-01069 Dresden, Germany

<sup>3</sup> Lawrence Livermore National Laboratory, Livermore, CA 94550, United States of America

<sup>4</sup> Department of Physics, University of California, Berkeley, CA 94720, United States of America

<sup>5</sup> SLAC National Accelerator Laboratory, Menlo Park, CA 94309, United States of America

<sup>6</sup> Institut für Kernphysik, Technische Universität Darmstadt, Schlossgartenstraße 9, D-64289 Darmstadt, Germany

<sup>7</sup> Centre for Fusion, Space and Astrophysics, Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

<sup>8</sup> Open and Transdisciplinary Research Institute, Osaka University, Suita, Osaka 565-0871, Japan

<sup>9</sup> LULI, UMR7605, CNRS-CEA, Université Paris VI-Ecole Polytechnique, F-91128 Palaiseau Cedex, France

E-mail: [d.kraus@hzdr.de](mailto:d.kraus@hzdr.de)

Received 9 July 2018, revised 22 August 2018

Accepted for publication 29 August 2018

Published 13 November 2018



CrossMark

## Abstract

We discuss the possibility of obtaining highly precise measurements of the ionization potential depression in dense plasmas with spectrally resolved x-ray scattering, while simultaneously determining the electron temperature and the free electron density. A proof-of-principle experiment at the Linac Coherent Light Source, probing isochorically heated carbon samples, demonstrates the capabilities of this method and motivates future experiments at x-ray free electron laser facilities.

Keywords: warm dense matter, dense plasma, ionization potential depression, x-ray scattering

(Some figures may appear in colour only in the online journal)

## 1. Introduction

The complex properties of dense plasmas play a crucial role in our understanding of celestial bodies like giant planets, brown dwarfs and stars as well as scientific and technological applications like intense laser-matter interaction (both optical and x-ray), fusion energy studies and radiation damage research [1]. For such systems, ionization is a key quantity that is very challenging to describe since in dense matter, the

bound states, and accordingly the ionization balance, are modified by the interaction with the surrounding medium. In particular, the influence of neighboring ions and screening due to continuum electrons results in reduced binding strength, which can be modeled by introducing effective (lower) ionization energies (ionization potential depression (IPD)). Although a precise description of the complex interaction of a dense plasma including bound states remains difficult, the Stewart and Pyatt model [2], which interpolates

between well-known low-density, high-temperature (Debye–Hueckel) and high-density, low-temperature (ion sphere) limits, is thought to catch most of the essential physics and is widely applied. However, several experiments have recently challenged this method, indicating a significantly larger depression of the ionization energy [3–5], whereas other results are in line with this description [6]. This controversy motivated several more advanced model approaches [7–10], but so far, no final convergence has been reached [11].

What is still lacking to resolve these discrepancies are experiments which can precisely measure electron temperature, free electron density (ionization) and IPD at the same time. In this article, we discuss dense carbon plasmas as an example and show very promising capabilities of high-precision spectrally resolved x-ray scattering, as now enabled by x-ray free electron lasers, to address present controversies on this topic. Dense plasma environments can be created by ultrafast isochoric heating with these highly brilliant x-ray sources [12]. Moreover, the x-rays can be applied to simultaneously characterize a set of plasma properties, particularly the IPD as well as ionization and electron temperature, with high-precision *in situ* spectrally resolved x-ray scattering. After introducing the basic concepts of this measurement technique focussing on the IPD measurement of dense carbon plasmas, we discuss a corresponding proof-of-principle experiment on isochorically heated graphite that was recently performed at the Linac Coherent Light Source (LCLS) of SLAC National Accelerator Laboratory and provides a first step forward towards future high-precision experiments.

## 2. Basic concepts

Carbon has the highest melting temperature of all elements and is of high interest for isochoric heating experiments. Cold carbon is dominated by electronic band structure and only the K-shell electrons can be treated in an atomic approximation. When entering the warm dense matter regime at a few thousand kelvins, significant bonding remains [13], while some of the L-shell electrons may be treated atomically before reaching significant ionization at higher temperatures. In the dense plasma regime with temperatures of several eV, plasma physics can be applied to describe the system via ionic bound states, free electrons and IPD. In addition, carbon is probably the most practical low- $Z$  material for experiments on dense plasma properties. In contrast to hydrogen, helium, nitrogen, oxygen, etc, it exists in a solid state at ambient conditions, which simplifies the experimental creation of solid-density plasmas. Moreover, its two ambient polymorphs, graphite ( $\rho = 2.2 \text{ g cm}^{-3}$ ) and diamond ( $\rho = 3.5 \text{ g cm}^{-3}$ ) allow for accessing conditions of different density by simply switching between these two forms of carbon as initial material. Furthermore, carbon is not as hazardous and not as difficult to handle as, e.g., beryllium or lithium. By using organic compounds as sample material, carbon can easily be studied in

mixtures with light elements that are of high relevance for astrophysics [14] and fusion applications [15].

Many of the characteristic properties of dense plasmas are accessible by spectrally resolved x-ray scattering. In general, the scattered radiation power spectrum per solid angle is given by [16]

$$\frac{dP}{d\Omega d\omega} = I_0 r_0^2 (1 - \cos^2 \phi \sin^2 \theta) N S(k, \omega), \quad (1)$$

where  $I_0$  is the initial probe intensity,  $r_0$  the classical electron radius,  $\phi$  the FEL polarization angle,  $\theta$  the scattering angle, and  $N$  the number of atoms in the probe volume.  $S(k, \omega)$  denotes the electron structure factor in dependence of scattering wave number  $k$  and frequency shift  $\omega$ , and contains the microscopic properties of the sample. In a plasma, x-rays either scatter from single electrons (non-collective scattering) or from collective fluctuations of many electrons (collective scattering), depending on the scattering geometry applied in the experiment [17]. The scattering parameter


$$\alpha = \frac{1}{k\lambda_s} \quad (2)$$

relates the scattering wave number  $k$  to the plasma screening length  $\lambda_s$  (e.g. Debye length for an ideal plasma). A scattering parameter of  $\alpha \ll 1$  defines the regime of non-collective scattering whereas  $\alpha \gtrsim 1$  results in collective scattering. For a non-collective scattering geometry, the structure factor is often split into three separate contributions (Chihara decomposition [18]):


$$S(k, \omega) = W_{el}(k)\delta(\omega) + W_{b-f}(k, \omega) + W_{f-f}(k, \omega). \quad (3)$$

Here, the first term describes elastic scattering, which is dominated by scattering from tightly bound electrons. The second term accounts for the inelastic scattering from weakly bound electrons and the third term describes inelastic scattering from free electrons. In general, design and analysis of spectrally resolved x-ray scattering experiments on dense plasmas need to be handled with care since a detailed data evaluation usually requires several model assumptions. For example, the Chihara decomposition can become problematic for plasmas of such extreme density that bound and free electrons can hardly be distinguished. Moreover, in experiments on extremely short timescales and with large gradients, one cannot apply local thermal equilibrium assumptions. Nevertheless, for a carefully chosen sample and scattering geometry, several physical quantities can be deduced from a single scattering spectrum. Particularly, these properties allow for deducing the IPD (bound-free feature), ionization  $Z$  (free-free feature in relation to elastic scattering and/or bound-free scattering) and electron temperature  $T_e$  (Doppler broadening of free-free feature) from a model fit to a measured spectrum. Furthermore, the ion correlations can be inferred from the ratio of elastic and inelastic scattering, which is highly sensitive to structural transitions, such as melting [19–21].

To model the IPD for the creation of synthetic x-ray scattering spectra, we apply two analytical models that have been extensively used in recent studies: Stewart–Pyatt [2] and Ecker–Kroell [22]. The Stewart–Pyatt approach interpolates



**Figure 1.** Synthetic scattering spectra for isochorically heated graphite varying  $Z$  at  $T_e = 20$  eV. For  $Z = 2$ , the free-free, bound-free and elastic features are shown as well.



**Figure 2.** Synthetic scattering spectra for isochorically heated graphite varying  $T_e$  at  $Z = 2$ .

between the Debye and ion sphere models. Smooth solutions are obtained on the level of electrostatic potential generated by the surrounding medium. The Debye model is always valid far away from the bound state and the ion sphere model close to the central atom/ion. The transition point is defined by requiring a smooth induced field. Another model was derived by Ecker and Kroell and has been discussed in a modified form in various recent publications [3, 6, 7]. Stewart and Pyatt however criticized this approach for violating charge neutrality. Nevertheless, since some recent experiments were matched by this model [3, 12], we include it for the generation of synthetic scattering spectra in our study. For high densities, the approach by Ecker and Kroell is very similar to the ion sphere model but is formed in terms of the total particle density. For low densities, the predicted IPD is again given by the Debye model. The Ecker–Kroell treatment includes a constant that can in principle be set with considerable freedom. Consistent with recent work, we set it to unity, which is usually called the ‘modified’ Ecker–Kroell model [3, 5].


### 3. Synthetic scattering spectra

Figures 1 and 2 show synthetic non-collective scattering spectra for 5.9 keV photon probe energy at a scattering angle of  $\theta = 160^\circ$  ( $k = 5.9 \text{ \AA}^{-1}$ ,  $\alpha \sim 0.2$ ) for varying ionization (figure 1) and electron temperature (figure 2) calculated by the XRS code [23] using an estimated instrument function combining the LCLS bandwidth in self-amplified spontaneous emission (SASE) mode and the resolution of the x-ray spectrometer. Since the frequency dependence of the elastic scattering is approximately a delta function on this energy scale, this scattering feature directly reflects the instrument function. Investigating the synthetic spectra on a logarithmic scale, particularly the blue-shifted part of the inelastic scattering on the high-energy side of the elastic scattering is very sensitive to ionization and electron temperature. Moreover, this feature is not obscured by remaining L-shell bound-free scattering since bound-free scattering with positive energy transfer is prevented by the corresponding ionization potential. Thus, an increase in  $Z$  at fixed electron temperature results in a linear increase of the scattering intensity on the high-energy side of the elastic scattering. On the other hand, an increase of electron temperature for a fixed  $Z$  results in a change in slope of the inelastic feature, since the electron energy distribution is broadened for higher electron temperatures. If the instrument function of the x-ray source and the spectrometer are well characterized over a dynamic range of three orders of magnitude in intensity, ionization and electron temperature of the plasma can therefore be inferred from blue-shifted inelastic scattering with high precision. Fitting the rest of the spectrum in addition can serve as a valuable cross-check of applied bound-free models and provide additional information, particularly the K-shell ionization energy and with that the IPD. It should be noted that for low ionization and low temperatures, remaining band structure [24] and chemistry [25] influences the L-shell wave functions and therefore the L-shell bound-free feature. This is not included in the presented single-ion calculations of the electronic structure factor and therefore, all illustrated curves for  $Z = 0$  and  $T_e = 0.1$  eV should only be interpreted as rough estimates of the corresponding scattering spectra.


Figure 3 shows the same synthetic spectra as in figure 1, but zoomed in on the carbon K-shell bound-free edge for varying  $Z$  at  $T_e = 20$  eV for two IPD models: Stewart–Pyatt, and modified Ecker–Kroell [3]. The effects of the different models on the K-edge feature are clearly visible. Figure 4 shows the effect of increasing the initial density, e.g. by using diamond instead of graphite, which results in a significant increase of the IPD when ionization is present.

### 4. Experiments


Experimental scattering spectra were obtained at the matter in extreme conditions (MEC) endstation of LCLS [26, 27]. A sketch of the experimental setup is shown in figure 5. Pyrolytic graphite samples (90  $\mu\text{m}$  thick, initial density  $\rho_0 = 2.21 \text{ g cm}^{-3}$ )



**Figure 3.** Synthetic spectra for isochorically heated graphite zoomed on the carbon K-shell bound-free edge and varying  $Z$  at  $T_e = 20$  eV.




**Figure 4.** Synthetic spectra zoomed on the carbon K-shell bound-free edge for varying the plasma density by using graphite and diamond samples. The Stewart–Pyatt model was applied to determine the IPD.



**Figure 5.** Sketch of the experimental setup at LCLS.


were irradiated by 5.9 keV x-ray pulses of 50 fs in duration and with pulse energies of  $\sim 3$  mJ. Using beryllium lenses, the x-ray beam was focussed down to spot sizes below  $5 \mu\text{m}$  in diameter, which results in an estimated peak fluence of up to  $3 \times 10^4 \text{ J cm}^{-2}$ . Around 20.2% of the incident x-ray flux is absorbed in the sample, which mainly produces K-shell holes and photo-electrons. The range of 5.6 keV photo-electrons inside graphite is around 500 nm [28], which is significantly smaller than the x-ray spot size and ensures that most of the absorbed energy is



**Figure 6.** Model fit to experimental data obtained from isochorically heated graphite at LCLS.

available to heat the sample inside the x-ray spot. For low- $Z$  elements like carbon, K-shell holes mainly relax by Auger decay which quickly heats the electron subsystem. For the smallest spot size, we estimate that an energy of  $\sim 40$  eV is absorbed per atom in the sample volume. Depending on the ionization state, this suggests peak temperatures of the free electrons on the order of 10 eV or more.

The scattered x-rays were collected by a highly annealed pyrolytic graphite crystal spectrometer at a scattering angle of  $160^\circ$ . Figure 5 shows a spectrum recorded from isochorically heated graphite, which was obtained by accumulating scattering data of 10 separate shots. The elastic and inelastic scattering features can clearly be distinguished. Moreover, the inelastic feature exhibits a clear edge at the low-energy end which originates from the onset of K-shell bound-free scattering and is therefore shifted by the energy required for the ionization of K-shell electrons. A model fit to the scattering spectrum using the Chihara decomposition and assuming local thermal equilibrium provides a stable fit giving  $T_e = 21.7$  eV,  $Z = 1.71$ , and an IPD of 24 eV. For the free-free scattering we apply random phase approximation (RPA) [23] and the bound-free term is calculated in the impulse approximation [29] using an effective ionization energy that is given by the atomic binding energy and the plasma-induced IPD. More precise analysis may require advanced quantum statistical approaches that go beyond RPA, which, however, is not within the scope of this proof-of-principle study. Moreover, the required high-precision characterization of the instrument function for accurate measurements of electron temperature and ionization from the blue-shifted part of the free-free scattering is unfortunately not available for the presented data set. It is only accurate within slightly more than two orders of magnitude and alternate instrument functions with broader wings are also possible (see figure 6). A fit using this modified instrument function provides  $Z = 1.1$  and  $T_e = 15$  eV, while the IPD remains at 24 eV. Therefore, due to the uncertainty of the instrument function, both  $Z$  and  $T_e$



**Figure 7.** Experimental scattering spectra zoomed onto the carbon K-shell bound-free edge for different x-ray focal spot sizes.

can only be stated with a systematic error around 50%, while the fit error of the IPD is around 5%. Moreover, the applied bound-free model does not seem to be able to exactly reproduce the shape of the inelastic scattering around its maximum at the Compton energy shift. Nonetheless, the K-shell binding energy and with that the IPD can already be fitted very accurately from the position of the carbon K-shell bound-free edge.

Figure 7 shows experimental scattering spectra zoomed on the carbon K-shell bound-free edge that were obtained in the described LCLS experiment. For the smallest spot size and thus maximum heating, there is a clear shift of the K-shell bound-free edge towards higher energies that may be induced by increased ionization. Comparing this shift to the trends shown in figure 3, where only the modified Ecker–Kroell approach shows a significant shift towards higher energies for increasing the ionization from 0 to 1 and 2, suggests that the Stewart–Pyatt model indeed underestimates the IPD which is in line with other studies [3–5]. However, looking at the absolute value of the IPD obtained from the fit (24 eV) is in very good agreement with the Stewart–Pyatt prediction for the best fit plasma parameters (25.3 eV) and does not agree with modified Ecker–Kroell (47.7 eV). For lower ionization, which is well within the error margin of the available data, the situation is different. For  $Z = 1$  at the same electron temperature, Stewart–Pyatt predicts 13.3 eV and modified Ecker–Kroell provides 24.6 eV. This underlines the importance of precisely characterizing the ionization state of the plasma as well as the electron temperature for accurate discrimination between models for IPD in dense plasmas.

## 5. Conclusions

We have presented an effective method to measure the IPD in dense carbon plasmas while simultaneously obtaining electron temperature and ionization. Our proof-of-principle experiment performed at LCLS shows that it is in principle possible to record scattering spectra of the required quality.

While the present data is not yet sufficient to provide precise values for electron temperature and ionization, the IPD can be well constrained by the carbon K-edge in the scattering spectrum. Future experiments will be able to accumulate more spectra for both instrument function characterization and scattering experiments, which will result in the desired accuracy for the blue-shifted free-free scattering and therefore a precise characterization of electron temperature as well as ionization. We have shown that such precision is required to reliably discriminate between IPD models for dense plasma environments.

Additional information may be obtained by adding collective x-ray scattering at relatively small scattering angles to the experiment. This method can observe electron plasma waves (plasmons) that follow a dispersion relation that is sensitive to density, electron temperature as well as the electron–ion collision frequency [30]. For high dynamic range and sufficiently high temperatures, the electron temperature can also be deduced from comparing the intensity ratio of the upshifted and downshifted plasmon features (detailed balance [31]). However, obtaining precise information on plasmons at dense plasma conditions usually requires very low bandwidth x-ray sources and the relatively broad SASE spectrum may not be sufficient for that. Using self-seeding [32] or monochromators [33] will significantly reduce the x-ray flux available for heating the samples. Moreover, collective scattering can result in ambiguities in the analysis since electron in cold solids or liquids can also be excited collectively (band structure) and corresponding features may obscure the signatures of the free electrons [24].

Finally, it has to be said that for simplicity, the presented analysis and resulting conclusions assume static equilibrium conditions and neglect the dynamic nature of the heating and simultaneous x-ray scattering processes. This simplification can lead to additional ambiguities [34]. However, the influence of heating dynamics can be probed in the experiment by varying the x-ray pulse duration, which can easily be realized at x-ray free electron laser facilities. Another very useful instrument could be the application of the described spectroscopy method to two-color x-ray pump-probe capabilities with varying time delay between the pulses [35].

## Acknowledgments

This work was performed at the MEC instrument of LCLS, supported by the US Department of Energy Office of Science, Fusion Energy Science under contract No. SF00515. DK, B Barb, and RWF acknowledge support by the US Department of Energy, Office of Science, Office of Fusion Energy Sciences and by the National Nuclear Security Administration under Award Numbers DE-FG52-10NA29649 and DE-NA0001859. DK and RWF acknowledge support of this work by the US Department of Energy, Office of Science, Office of Fusion Energy Sciences under Award Number DE-SC0018298, and the University of California Center for Frontiers in High Energy Density Science. DK was supported by the Helmholtz Association under VH-NG-1141. NJH was

supported by Kakenhi Grant No. 16K17846. SLAC HED is supported by DOE Office of Science, Fusion Energy Science under FWP 100182. SF and JH were supported by German Bundesministerium für Bildung und Forschung project Nos. 05P12RDFA1 and 06DA9043I. The work of B Bach and TD was performed under the auspices of the US Department of Energy by Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, and supported by Laboratory Directed Research and Development (LDRD) Grant No. 18-ERD-03.

## ORCID iDs

D Kraus  <https://orcid.org/0000-0002-6350-4180>

## References

- [1] Fortov V E 2011 *Extreme States of Matter on Earth and in the Cosmos* (Berlin: Springer)
- [2] Stewart J C and Pyatt K D 1966 *Astrophys. J.* **144** 1203
- [3] Ciricosta O *et al* 2012 *Phys. Rev. Lett.* **109** 065002
- [4] Fletcher L B *et al* 2014 *Phys. Rev. Lett.* **112** 145004
- [5] Kraus D *et al* 2016 *Phys. Rev. E* **94** 011202(R)
- [6] Hoarty D J *et al* 2013 *Phys. Rev. Lett.* **110** 265003
- [7] Vinko S M *et al* 2014 *Nat. Commun.* **5** 3533
- [8] Crowley B J B 2014 *High Energy Density Phys.* **13** 84–102
- [9] Son S-K *et al* 2014 *Phys. Rev. X* **4** 031004
- [10] Lin C *et al* 2017 *Phys. Rev. E* **96** 013202
- [11] Iglesias C A 2014 *High Energy Density Phys.* **12** 5–11
- [12] Vinko S M *et al* 2012 *Nature* **482** 59
- [13] Kraus D *et al* 2013 *Phys. Rev. Lett.* **111** 255501
- [14] Kraus D *et al* 2017 *Nat. Astron.* **1** 606–11
- [15] Hurricane O A *et al* 2014 *Nature* **506** 343
- [16] Sheffield J, Froula D, Glenzer S H and Luhmann J N C 2011 *Plasma Scattering of Electromagnetic Radiation* (New York: Academic)
- [17] Glenzer S H and Redmer R 2009 *Rev. Mod. Phys.* **81** 1625
- [18] Chihara J 2000 *J. Phys.: Condens. Matter* **12** 231–47
- [19] Hau-Riege S P *et al* 2012 *Phys. Rev. Lett.* **108** 217402
- [20] Helfrich J *et al* 2015 *High Energy Density Phys.* **14** 38–43
- [21] Kraus D *et al* 2015 *Phys. Plasmas* **22** 056307
- [22] Ecker G and Kröll W 1963 *Phys. Fluids* **6** 62
- [23] Gregori G *et al* 2003 *Phys. Rev. E* **67** 026412
- [24] Gamboa E J *et al* 2015 *Phys. Plasmas* **22** 056319
- [25] Kraus D *et al* 2018 *Phys. Plasmas* **25** 056313
- [26] Nagler B *et al* 2015 *J. Synchrotron Radiat.* **22** 520–5
- [27] Glenzer S H *et al* 2016 *J. Phys. B: At. Mol. Opt. Phys.* **49** 092001
- [28] Berger M J, Coursey J S, Zucker M A, Chang J and ESTAR PSTAR ASTAR 2018 Computer programs for calculating stopping-power and range tables for electrons, protons, and helium ions (version 2.0.1) National Institute of Standards and Technology
- [29] Schumacher M *et al* 1975 *J. Phys. B: At. Mol. Phys.* **8** 1428–39
- [30] Sperling P *et al* 2015 *Phys. Rev. Lett.* **115** 115001
- [31] Glenzer S H *et al* 2007 *Phys. Rev. Lett.* **98** 065002
- [32] Fletcher L B *et al* 2015 *Nat. Photon.* **9** 274
- [33] Shu D *et al* 2013 *J. Phys.: Conf. Ser.* **425** 052004
- [34] Chapman D A and Gericke D O 2011 *Phys. Rev. Lett.* **107** 165004
- [35] Inoue I *et al* 2016 *Proc. Natl Acad. Sci.* **113** 1492–7