

Lawrence Berkeley National Laboratory

LBL Publications

Title

The Heat of Formation of Thorium Tetrachloride

Permalink

<https://escholarship.org/uc/item/9bh6f8kf>

Authors

Eyring, LeRoy

Westrum, Edgar F

Publication Date

1950-03-01

UCRL 623
~~9/1/8~~
C.2

DECLASSIFIED

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-623
C2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

CONFIDENTIAL

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

DECLASSIFIED

THE HEAT OF FORMATION OF THORIUM TETRACHLORIDE

LeRoy Eyring and Edgar F. Westrum, Jr.

March 8, 1950

CAUTION

This document contains information affecting the National Defense of the United States. Its transmission or the disclosure of its contents in any manner to an unauthorized person is prohibited and may result in severe criminal penalties under applicable Federal Laws.

Berkeley, California

DECLASSIFIED

CHEMISTRY-GENERAL

Standard Distribution: Series B

Copy Nos.

Argonne National Laboratory	1-8
Armed Forces Special Weapons Project	9
Atomic Energy Commission, Washington	10-11
Battelle Memorial Institute	12
Brookhaven National Laboratory	13-16
Bureau of Ships	17
Carbide and Carbon Chemicals Corporation (K-25 Plant)	18-21
Carbide and Carbon Chemicals Corporation (Y-12 Plant)	22-25
Columbia University (Failla)	26
Dow Chemical Company	27
General Electric Company, Richland	28-33
Hanford Operations Office	34
Idaho Operations Office	35-36
Iowa State College	37
Knolls Atomic Power Laboratory	38-41
Los Alamos	42-44
Mallinckrodt Chemical Works	45
Massachusetts Institute of Technology (Gaudin)	46
Massachusetts Institute of Technology (Kaufmann)	47
Mound Laboratory	48-50
National Advisory Committee for Aeronautics	51
National Bureau of Standards	52-53
Naval Radiological Defense Laboratory	54
NEPA Project	55
New Brunswick Laboratory	56
New York Operations Office	57-58
North American Aviation, Inc.	59
Oak Ridge National Laboratory	60-71
Patent Branch, Washington	72
Sylvania Electric Products, Inc.	73
Technical Information Branch, ORE	74-88
UCLA Medical Research Laboratory (Warren)	89
University of California Radiation Laboratory	90-93
University of Rochester	94-95
Western Reserve University (Friedell)	96
Westinghouse Electric Corporation	97-98



THE HEAT OF FORMATION OF THORIUM TETRACHLORIDE(*)

LeRoy Eyring(**) and Edgar F. Westrum, Jr.(***)
Department of Chemistry and Radiation Laboratory
University of California, Berkeley, California
March 8, 1950

DECLASSIFIED

ABSTRACT

The heats of solution of thorium metal and of thorium tetrachloride in six molar hydrochloric acid have been measured. The addition of a very small concentration of sodium fluosilicate was required to obtain quantitative solution of the metal. The heat of formation of $\text{ThCl}_4(\text{c})$ is calculated to be -284.5 kcal/mole at 25°C . and is markedly lower than the value reported in the literature.

(*) This work was performed under Manhattan District Contract W-31-109-Eng-38 in 1946 at the Radiation Laboratory of the University of California at Berkeley.

(**) Present address: Department of Chemistry, The State University of Iowa, Iowa City, Iowa.

(***) Present address: Department of Chemistry, University of Michigan, Ann Arbor, Michigan.

Current interest in the elements of the actinide series implies a scientific and technical requirement of accurate thermodynamic data on the compounds of members of this series. Some measurements by Westrum and Robinson⁽¹⁾

- (1) E. F. Westrum, Jr. and H. P. Robinson, National Nuclear Energy Series, Plutonium Project Record, Vol. 14B, "The Transuranium Elements: Research Papers," Paper No. 6.50 (McGraw-Hill Book Co., Inc., New York, 1949).
-

on the heat of solution of thorium tetrachloride in several concentrations of hydrochloric acid indicated that as a result of neglecting the concentration of the acid used to dissolve the thorium metal in the evaluation and combination of the experimental data, the heat of formation of thorium tetrachloride, reported by Chauvenet⁽²⁾ is about nine kilocalories too large. Even after the

- (2) E. Chauvenet, Annales de Chimie et de Physique, Series 8, Volume 23, 425 (1911).
-

application of this correction, the reported heat of formation appeared surprisingly large. Furthermore it is unlikely, on the basis of recent information,⁽³⁾

- (3) R. W. Nottorf, A. S. Wilson, R. E. Rundle, A. S. Newton, J. E. Powell, reported in Manhattan Project Document CC-2722 March 8, 1945.
-

that pure thorium metal would have been obtained by the process described by Chauvenet.

The present measurements were made on two different very pure samples of thorium metal prepared by the Manhattan District Laboratory at Ames, Iowa, and some microscopic crystals of anhydrous thorium tetrachloride obtained indirectly from Lindsay Light and Chemical Co.

Thorium metal does not appear to dissolve completely in 6M HCl at room temperature even on long standing. The bulk of the metal (ca. 75%) dissolves quite readily in a short time giving rise to a voluminous black residue the exact composition of which has not been determined. This residue can be dissolved by more stringent treatment such as the addition of fluosilicate, etc., leaving only a relatively small white precipitate. The addition of a very small concentration of sodium fluosilicate (Na_2SiF_6) to the acid effects solution of more than 98% of the sample leaving only the white crystalline residue of thorium dioxide known to be present in the original sample.

PURITY OF THE MATERIALS

Samples of pure thorium metal prepared by the reduction of thorium tetrachloride with zinc and subsequent removal of the zinc by vacuum distillation were made available to us. A section cut from the middle portion of the ingot identified as H-3-15 was the source of our samples and the analytical samples from which were obtained the following information relative to the purity of the metal by the Analytical Section of the Metallurgical Laboratory. Duplicate assays yielded the following information: Th (99.5%, 99.7%); C (600 ppm, 560 ppm); N (773 ppm, 795 ppm), and H (133 ppm, 135 ppm). Spectrochemical analyses set limits on the maximum amounts of metallic impurities as less than one part per million of B, Bi, Cd, K, and Li; less than ten ppm of Ag, As, Ca, Na, Sb, and Sn; less than 20 ppm of Co, Cr, Ni, and P; less than 100 ppm of Al and Mo; less than 200 ppm of Ti. Elements detected were 10 ppm of Be, 20 ppm of Cu, 40 ppm of Pb, 70 ppm of Mn, 200 ppm of Si, 800 ppm of Mg, and 2000 ppm of Zn. Survey experiments upon another sample of metal identified as AEC-21 with a similar typical analysis showed entirely similar behavior. Separate measurements of the hydrogen evolved upon dissolution of

metal in HCl solutions containing fluosilicate confirmed the purity of the metal and the completeness of dissolution under the calorimetric conditions. The metal increases in weight 13.86% on combustion in air corresponding to 100.3% of free metal if the light impurities are neglected in the calculation.

Crystals about 2 mm. in length of anhydrous thorium tetrachloride prepared by Lindsay Light and Chemical Company carried an analysis of 62.8% Th and 37.8% Cl (theoretical: 62.07% Th and 37.93% Cl).

Sodium fluosilicate of C.P. grade was used. The aqueous HCl was standardized against mercuric oxide.

EXPERIMENTAL METHOD AND DATA

The heat of solution measurements were made in a calorimeter previously described.⁽⁴⁾ Calibrations of the electrical resistances and the heat

(4) Westrum and Robinson, op. cit., Paper No. 6.51.

capacity of the system were confirmed after shipment of the instrument.

The heat of solution of thorium tetrachloride data are presented in Table I. The data for 6.0M HCl are in excellent agreement with the value 44.21 ± 0.1 kcal/mole obtained by Westrum and Robinson⁽¹⁾ on finely divided, crystalline vacuum sublimed anhydrous ThCl_4 . The difference in the observed heat of solution in the presence of the Na_2SiF_6 is definitely beyond experimental error. It was noted that the time required for practical completion of the thermal effect was 2, 3, and 4 minutes in runs A, B, and C, respectively; approximately a minute was required in the presence of Na_2SiF_6 .

The amount of thorium dissolved was determined by precipitation of thorium oxalate and gravimetric determination of the dioxide. The interference of the small concentration of fluosilicate was noted, but by carrying out the analyses in a reproducible manner an empirically determined correction was applicable.

Table I

The Heat of Solution of Thorium Tetrachloride

In 6.004 Molar Hydrochloric Acid at 25°C.

Molecular weight $\text{ThCl}_4 = 373.95$

Weight of acid = 212.25 grams (vacuo)

Run	Millimoles ThCl_4 (vacuo weight of samples)	Millimoles ThCl_4 (by oxalate ppt)	Observed Heat Evolved (calories)	Heat of Solution (kcal/mole)
A	0.4423	0.4424	19.59	-44.29
B	0.8247	0.8254	36.49	-44.25
C	1.0450	1.0445	46.29	-44.30
			Average	-44.28 \pm 0.06
<u>With 0.00500 Molar Sodium Fluosilicate Added</u>				
D	1.0627	1.0638	47.38	-44.58
E	0.5633	0.5630	25.11	-44.58
			Average	-44.58 \pm 0.08

Table II

The Heat of Solution of Thorium Metal in

6.004 Molar Hydrochloric Acid

with 0.00500 Molar Sodium Fluosilicate Added at 25°C.

Atomic weight of Th = 232.12


Weight of acid = 212.25 grams (vacuo)

Run	Millimoles Th (vacuo weight of samples)	Millimoles Dissolved (by oxalate precipitation)	Percent in Soln.	Observed Heat Evolved (calories)	Observed Heat of Solution k/cal/mol	*Corrected heat of solution kcal/mol
J	0.4515	0.4449	98.5	80.70	181.6	181.8
K	0.4446	0.4379	98.5	79.39	181.4	181.6

*Correction applied for evaporation of solution by hydrogen evolution, dissolution of light metals and reaction of thorium compounds present as impurities.

THE HEAT OF FORMATION

The following cycle is used to evaluate the heat of formation:



Typical values are: $a = 2400$, $b/a = 8.105$

Reactions (1) and (2) correspond to the experimental measurements of the heat of solution of thorium metal and thorium tetrachloride, respectively. Reaction (3) is evaluated as the apparent heat of formation of hydrochloric acid in the 6.684 molal aqueous solution from the heat of formation of the infinitely dilute aqueous solution of hydrogen chloride,⁽⁵⁾⁽⁶⁾ and the relative

(5) F. D. Rossini, J. Research Natl. Bur. Standards 9:679 (1932).

(6) Selected Values of Chemical Thermodynamic Properties, National Bureau of Standards, Washington (1947).

molal heat content of the 6 molar acid. The heat of this reaction is assumed to be uninfluenced by the small concentration of the thorium salt since the ionic strengths of the solutions will be nearly unchanged by its removal.

The heat of formation of ThCl_4 at 25°C . is, therefore:

$$\begin{aligned}
 \Delta H_4 &= \Delta H_1 - \Delta H_2 + \Delta H_3 \\
 &= -181.7 + 44.6 - 4(36.84) = 284.5 \text{ kcal/mole.}
 \end{aligned}$$

A linear extrapolation of the heat of solution of thorium tetrachloride from one molar⁽¹⁾ concentration of hydrogen chloride to infinite dilution

yields a value of -60.0 for the heat of solution in the infinity dilute acid. The heat of formation of the thorium ion $\text{Th} + 4\text{H}^+ (\text{aq}) = \text{Th}^{+4} (\text{aq}) + 2\text{H}_2 (\text{g})$ is calculated to be -184.4 kcal/mole by combining the above value with the heat of formation of the infinitely dilute aqueous solution of hydrogen chloride⁽⁶⁾ and the heat of formation of ThCl_4 .

ACKNOWLEDGMENT

Credit for the performance of this work is gratefully shared with those who supplied the materials and the analyses of them. This applies especially to the Manhattan Project at Ames for the thorium metal, the Analytical Section of the Metallurgical Laboratory for chemical and spectrographic analysis of the thorium metal and the thorium tetrachloride, and to Winifred Heppler for technical assistance.



DECLASSIFIED