UC Irvine

UC Irvine Previously Published Works

Title

Genetic diversity of the Plasmodium falciparum apical membrane antigen I gene in parasite population from the China-Myanmar border area.

Permalink

<https://escholarship.org/uc/item/9bw4s3n3>

Authors

Zhu, Xiaotong
Zhao, Zhenjun
Feng, Yonghui
et al.

Publication Date

2016-04-01

DOI

10.1016/j.meegid.2016.01.021

Peer reviewed

Research paper

Genetic diversity of the *Plasmodium falciparum* apical membrane antigen I gene in parasite population from the China–Myanmar border area Xiaotong Zhu^a, Zhenjun Zhao^b, Yonghui Feng^c, Peipei Li^b, Fei Liu^a, Jun Liu^a, Zhaoqing Yang^d, Guiyun Yan^e, Qi Fan^b, Yaming Cao^{a,*}, Liwang Cui^{a,f,**}^a Department of Immunology, College of Basic Medical Science, China Medical University, Shenyang, Liaoning 110122, China^b Dalian Institute of Biotechnology, Dalian, Liaoning, China^c Department of Laboratory Medicine, The First Hospital of China Medical University, Shenyang, Liaoning 110001, China^d Department of Pathogen Biology and Immunology, Kunming Medical University, Kunming, China^e Program in Public Health, University of California, Irvine, CA, USA^f Department of Entomology, The Pennsylvania State University, University Park, PA 16802, USA

ARTICLE INFO

Article history:

Received 16 October 2015

Received in revised form 20 January 2016

Accepted 23 January 2016

Available online 26 January 2016

Keyword:

Plasmodium falciparum

PfAMA1

Genetic diversity

China–Myanmar border

Malaria

ABSTRACT

To investigate the genetic diversity of the *Plasmodium falciparum* apical membrane antigen 1 (PfAMA1) gene in Southeast Asia, we determined PfAMA1 sequences from 135 field isolates collected from the China–Myanmar border area and compared them with 956 publically available PfAMA1 sequences from seven global *P. falciparum* populations. This analysis revealed high genetic diversity of PfAMA1 in global *P. falciparum* populations with a total of 229 haplotypes identified. The genetic diversity of PfAMA1 gene from the China–Myanmar border is not evenly distributed in the different domains of this gene. Sequence diversity in PfAMA1 from the China–Myanmar border is lower than that observed in Thai, African and Oceanian populations, but higher than that in the South American population. This appeared to correlate well with the levels of endemicity of different malaria-endemic regions, where hyperendemic regions favor genetic cross of the parasite isolates and generation of higher genetic diversity. Neutrality tests show significant departure from neutrality in the entire ectodomain and Domain I of PfAMA1 in the China–Myanmar border parasite population. We found evidence supporting a substantial continent-wise genetic structure among *P. falciparum* populations, with the highest genetic differentiation detected between the China–Myanmar border and the South American populations. Whereas no alleles were unique to a specific region, there were considerable geographical differences in major alleles and their frequencies, highlighting further necessity to include more PfAMA1 alleles in vaccine designs.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Plasmodium falciparum is the most deadly of five *Plasmodium* species that infect humans, with an estimated 584,000 deaths (90% from sub-Saharan Africa and 78% children <5 years of age) in 2013 (WHO, 2014). Increasing drug resistance of the parasites and insecticide resistance of the vector mosquitoes have made malaria control difficult, and there is a strong demand for an effective vaccine to contain this deadly disease. However, antigen polymorphism of many asexual stage vaccine candidates such as merozoite surface protein 1 (MSP1), MSP2, and MSP3 (Jordan et al., 2009; Koukouikila-Koussounda et al., 2012; Sakihama et al., 2001) hamper the development of vaccines effective against all parasite populations. Thus, successful interventions

targeting different developmental stages will require better understanding of genetic variations of target antigens within and between parasite populations.

Apical membrane antigen 1 (AMA1) is conserved in apicomplexans, and *P. falciparum* apical membrane antigen-1 (PfAMA1) is one of the leading blood stage vaccine candidates in human trials (Dutta et al., 2003; Kocken et al., 2002; Mitchell et al., 2004). PfAMA1 is an 83 kDa antigen synthesized by mature blood stages of the parasite and is initially localized in the micronemes of the merozoite, an apical organelle that plays a key role in erythrocyte invasion (Healer et al., 2002; Narum and Thomas, 1994; Peterson et al., 1989). Prior to merozoite invasion, PfAMA1 is processed into a 66 kDa product and released onto the merozoite surface, where they participate in the formation of tight junctions by interacting with the RON proteins and transducing the force generated by the parasite motor during internalization (Bargieri et al., 2012; Cao et al., 2009; Richard et al., 2010). The complete 1686 bp PfAMA1 coding region contains an ectodomain with three sub-domains (Domains I–III) (Silvie et al., 2004). A higher rate of mutations and level of diversifying selection have been shown in Domain I (Cortes

* Corresponding author.

** Correspondence to: L. Cui, Department of Immunology, College of Basic Medical Science, China Medical University, Shenyang, Liaoning 110122, China.

E-mail addresses: ymcao@mail.cmu.edu.cn (Y. Cao), luc2@psu.edu (L. Cui).

et al., 2003; Figtree et al., 2000; Garg et al., 2007; Thakur et al., 2008). However, Domain II presents a high-degree of amino acid sequence conservation. A loop region within this domain has been shown to contain an epitope recognized by an invasion-inhibitory monoclonal antibody (Chesne-Seck et al., 2005).

The ectodomain of AMA1 are highly immunogenic. Antibodies raised against AMA1 inhibit merozoite invasion in vitro (Crewther et al., 1996; Healer et al., 2004; Hodder et al., 2001; Kennedy et al., 2002; Kocken et al., 2002; Kusi et al., 2009; Remarque et al., 2008). Immunization with recombinant AMA1 has been shown to elicit protective antibody response against homologous strain challenges in both rodent and primate models, but the response is less protective against a heterologous parasite line (Healer et al., 2004; Remarque et al., 2008). Antibodies to AMA1 are typically highly prevalent among malaria endemic populations (Thomas et al., 1994). A systematic meta-analysis of data that met rigorous quality criteria detected a tendency towards a protective association among studies of AMA1 (Fowkes et al., 2010). These results support the development of AMA1 as a malaria vaccine, but also highlight the need to better understand genetic diversity of *PfAMA1*.

The Greater Mekong Subregion (GMS) is one of the most threatening foci of malaria in Southeast Asia (Cui et al., 2012). The recent emergence of artemisinin resistant *P. falciparum* parasites has posed a serious problem for both regional and global malaria control (Dondorp et al., 2011). Within the GMS, more than half of the malaria cases and an estimated 75% of the malaria deaths occurred in Myanmar (Wang et al., 2015). Moreover, international border regions, such as the one shared by China and Myanmar have the highest malaria incidence in the GMS (Cui et al., 2012). Border malaria is extremely difficult to monitor and control, largely due to frequent human population movements across the porous international borders. In the present study, we investigated the genetic diversity in the ectodomain of *PfAMA1* gene in the China–Myanmar border area. By comparing with global *P. falciparum* populations, we identified important differences in the *PfAMA1* alleles and their frequencies, which bear important implications for the design of *PfAMA1* based vaccines.

2. Materials and methods

2.1. Sample collection

A total of 171 field isolates were collected by finger-prick from patients attending malaria clinics in northwest Myanmar along the China–Myanmar border between April 2011 and October 2013. Blood samples were confirmed for *P. falciparum* infections based on microscopy of Giemsa-stained thick smears (Li et al., 2013). Use of the samples for this study was approved by the Biomedical Research Ethics Review Board of China Medical University.

2.2. DNA isolation, PCR amplification and sequencing of the *PfAMA1* gene

Plasmodium DNA was extracted from filter papers or whole blood using QIAamp DNA Blood Mini kit (QIAGEN, Hilden, Germany) according to the manufacturer's protocol. Parasites were genotyped at three polymorphic loci to exclude multi-clonal infections (Meng et al., 2010). Extracellular coding region contain Domains I–III (DI–III) of *PfAMA1* were amplified using the following primer pairs: *Pfama1_F* (5′-GAAGTTCATGGTTTCAGGTATAAG-3′), *Pfama1_R* (5′-GTATGGTTTTCATCAGAACTGG-3′), *Pfama1_NF* (5′-GATGCTGAAGTAGCTGGAAGTC-3′), and *Pfama1_NR* (5′-GTGATGCTCTTTTCTTCCCCC-3′). The PCR reaction contained 2 µl of 10× KOD-Plus-Neo buffer, 2 µl of 2 mM dNTPs, 0.8 µl of 25 mM MgSO₄, 0.5 µl of 10 µM of each primer, 0.5 units of KOD-Plus-Neo DNA polymerase (Toyobo, Osaka, Japan), and 1.0 µl genomic DNA template in a final volume of 20 µl. The cycling parameters for the primary PCR were as follows: initial denaturation at 94 °C for 2 min, 45 cycles of 94 °C for 15 s, 56 °C for 15 s, and 68 °C for 90 s, and a final extension at 68 °C for 5 min. The amplification conditions for

nested PCR were the same as those for primary PCR except that the number of cycles was 35 cycles. Nested PCR products were agarose gel purified using the QIAquick Gel Extraction Kit (QIAGEN, CA, USA), as per manufacturer protocol for preparation of templates for sequencing. Purified PCR products were sequenced with primers *Pfama1_NF* and *Pfama1_NR* in both directions using the ABI Prism® BigDye™ cycle sequencing kit (Applied Biosystems, Foster City, CA, USA).

2.3. Sequence alignment, polymorphism and statistical analyses

Of the 141 *P. falciparum* infections amplified by PCR, *PfAMA1* was successfully sequenced from 135 samples (sequences submitted to GenBank under accession nos.: KT897327–KT897461). To evaluate the polymorphism of *PfAMA1*, the *AMA1* gene (PF3D7_1133400) from the *P. falciparum* 3D7 strain was used as the reference gene. A single contiguous 1347 bp of *PfAMA1* (nucleotide region 445–1791 and codons 149–597) was derived for each of the 135 *PfAMA1* sequences, which include DI (codons 149–302), DII (codons 320–418) and DIII (codons 443–509). These sequences were aligned using the CLUSTAL W program in MEGA6.0 (Tamura et al., 2013) and exported as a FASTA alignment for statistical analysis using the DnaSP5.10.01 software (Librado and Rozas, 2009). The number of segregating (polymorphic) sites (*S*), nucleotide diversity (π), the average number of nucleotide differences (*k*), the number of haplotypes (*H*) and haplotype diversity (*H_d*) were calculated by DnaSP (Librado and Rozas, 2009). The distribution of nucleotide diversity (π) across the DI–III of *AMA1* gene was analyzed using the sliding window method with a window size of 90 bp and a step size of 3 bp.

2.4. Haplotype network construction, linkage disequilibrium (LD) and *F_{st}* analysis

The haplotype network was constructed using the program NETWORK Version 4.6.1.3 with the Median-joining method (Bandelt et al., 1999). A total of 956 publically available *PfAMA1* sequences representing seven *P. falciparum* populations were retrieved from GenBank: Thailand (Polley et al., 2003), Papua New Guinea (Arnott et al., 2014), Gambia (Tetteh et al., 2009), Nigeria (Polley and Conway, 2001), Kenya (Osier et al., 2010), Mali (Takala et al., 2009), and Venezuela (Ord et al., 2008). Analysis of the minimum number of recombination events (*R_M*) (Hudson and Kaplan, 1985), recombination parameter *C* ($C = 4Nr$, where *N* is the effective population size and *r* is the probability of recombination between adjacent nucleotides per generation) (Hudson, 1987), as well as linkage disequilibrium (LD) using the indices *D'* (Lewontin, 1964) and *R²* (Hill and Robertson, 1968) were calculated by DnaSP (Librado and Rozas, 2009). To assess the proportion of genetic variance due to population subdivision, the inter-population variance in allele frequency at the *PfAMA1* locus was compared among populations by calculating the Wright's *F_{st}* using DnaSP (Librado and Rozas, 2009).

2.5. Molecular evolutionary analysis

To test deviation from neutral evolution, the ratio of nonsynonymous substitutions to synonymous substitutions (*d_N/d_S*) were calculated using the Nei and Gojobori method (Nei and Gojobori, 1986) with the Jukes and Cantor correction and were compared with the Z-test of selection ($P < 0.05$) in MEGA6.0 (Tamura et al., 2013). A significant excess of *d_N* relative to *d_S* indicates positive natural selection, whereas negative values indicate negative or purifying selection (Nei and Gojobori, 1986). Departure from neutrality was estimated by Tajima's *D* test, Fu and Li's *D** and *F** tests and McDonald–Kreitman test. In Tajima's *D* test, departure from neutrality in the nucleotide frequency distributions is determined by divergence in the values of π (observed average pairwise nucleotide diversity) and θ (expected nucleotide diversity under neutrality derived from the number of segregating sites (*S*) (Tajima, 1989). Fu and Li's *D** and *F** tests reflect

the same trends as Tajima's D test and evaluate departure from neutrality by comparing the number of mutations in the external (considered to be "new" mutations) and internal (considered to be "older" mutations) branches of the genealogy. The number of mutations in external phylogeny branches was estimated from singleton sites (S) and the total number of mutations giving the D^* index, or between S and η giving the F^* index (Fu and Li, 1993). A positive value of Tajima's D and Fu and Li's D^* and F^* tests corresponds to positive natural selection, whereas a negative value indicating population size expansion and/or negative/purifying selection. Sliding window plots with a window size of 90 bp and a step size of 3 bp were also performed for Tajima's D and Fu and Li's D^* and F^* tests by DnaSP5.10.01 (Librado and Rozas, 2009). McDonald–Kreitman test (McDonald and Kreitman, 1991) was applied by taking a single *Plasmodium reichenowi* AMA1 sequence as the outgroup (accession no. AJ252087; Kocken et al., 2000) using DnaSP. The ratio of interspecies fixed synonymous substitutions to nonsynonymous substitutions is compared to that of intraspecific synonymous substitutions to nonsynonymous substitutions. An excess of the ratio between species over within species suggests purifying selection. Two-tailed Fisher's exact test was computed to determine the statistical significance ($P < 0.05$).

3. Results

3.1. Sequence diversity of the *PfAMA1* gene among the China–Myanmar border isolates

For the 135 *PfAMA1* sequences obtained from the China–Myanmar border isolates, the average number of nucleotide differences (k) for the entire 1347 bp sequenced region, DI, DII, and DIII were 15.15, 8.16, 1.55, and 2.06, respectively (Table 1). A total of 34 distinct haplotypes (H) were identified. The haplotype diversity (Hd) was relatively high, with an average of 0.859 in the entire sequenced region of all isolates (Table 1). Within the entire sequenced region (codons 149–597), there were 57 polymorphic sites. These polymorphic sites are distributed unevenly within the sequenced region, with 31, 7 and 8 being located in DI, DII, and DIII, respectively (Table 1). This is further illustrated in a sliding window plot of π across the sequenced region (Fig. 1A). Likewise, the pairwise nucleotide diversity (π) is highest in DI ($\pi = 0.018$), and lower in DII ($\pi = 0.005$) and DIII ($\pi = 0.010$). Also illustrated by the locations of the substitutions (Fig. 1B), DI (nt 445–906 and aa 149–302) is more polymorphic than DII (nt 958–1254 and aa 320–418) and DIII (nt 1327–1527 and aa 443–509).

3.2. Recombination and LD

We compared the recombination and LD of *PfAMA1* between populations from the China–Myanmar border area and other geographic regions. A minimum of 16 recombination events were detected in the China–Myanmar border and in Thai samples (Polley et al., 2003),

Fig. 1. Sliding window plot of nucleotide diversity and amino acid polymorphism of *PfAMA1* ectodomain in China–Myanmar border isolates. Nucleotide diversity is plotted with a window length of 90 bp and step size of 3 bp (A) and the number of the amino acid residues at each amino acid position is plotted, also to visualize the location sites showing high diversity, a scheme of the ectodomain of *ama1* is shown below (B). Black arrow indicates analyzed regions. A total of 135 sequences from China–Myanmar border are used. Nucleotide and amino acid positions are after the 3D7 line sequences.

compared to over 20 recombinant events detected in Papua New Guinea (Arnott et al., 2014) and African populations (Osier et al., 2010; Polley and Conway, 2001; Takala et al., 2009; Tetteh et al., 2009) (Table 2). The recombination parameter (C) between adjacent nucleotide sites and for the whole sequence had a value of 0.003 and 3.8, respectively in the China–Myanmar border population. This value was much lower than that from other geographic areas analyzed in this study but higher than that of the Venezuelan population (Ord

Table 1

Estimation of nucleotide diversity and summary statistics of *PfAMA1* in 135 *P. falciparum* isolates from the China–Myanmar border area^a.

	S	k	$\pi \pm SD$	H	$Hd \pm SD$	d_N/d_S	D	D^*	F^*	MK
Total	57	15.157	0.011 $\pm$ 0.001	34	0.859 $\pm$ 0.025	16.459***	1.198	1.911**	1.929*	0.000***
DI	31	8.167	0.018 $\pm$ 0.001	20	0.752 $\pm$ 0.036	18.909**	1.058	2.030**	1.964*	0.008**
DII	7	1.550	0.005 $\pm$ 0.000	9	0.705 $\pm$ 0.037	–	0.477	1.171	1.107	0.417
DIII	8	2.058	0.010 $\pm$ 0.000	11	0.699 $\pm$ 0.037	–	0.605	–0.271	0.045	0.308

^a The total sequenced region includes codons 149 to 597, Domain I codons 149 to 302, Domain II codons 320 to 418, and Domain III codons 443 to 509 (Hodder et al., 1996). S , number of polymorphic (segregating) sites; k , the average number of nucleotide differences, π , pairwise nucleotide diversity; H , number of haplotypes; Hd , haplotype diversity; d_N/d_S , the ratio of non-synonymous to synonymous mutations; D , Tajima's D test; D^* , Fu and Li's D^* value; F^* , Fu and Li's F^* value; MK , McDonald–Kreitman Test. SD = standard deviation.

* $P < 0.05$.

** $P < 0.02$.

*** $P < 0.001$.

Table 2

Comparison of different estimates of recombination events in *PfAMA1* among global isolates.

Locality (no.)	R^a	R^b	Rm
Myanmar (135)	0.003	3.8	16
Thailand (50)	0.051	66.5	16
PNG (76)	0.050	58.7	21
Gambia (114)	0.089	106	26
Nigeria (51)	0.160	207	25
Kenya (129)	0.123	146	26
Mali (506)	0.112	133	28
Venezuela (30)	0.000	0.001	1

R^a , recombinant parameter between adjacent sites; R^b , recombinant parameter for the whole gene; Rm, minimum number of recombination events.

et al., 2008) (Table 2). The LD index R^2 decreased rapidly with increasing nucleotide distance for the China–Myanmar border samples, indicating a high meiotic recombination rate (Fig. 2).

3.3. *Fst* analysis of *PfAMA1* in the China–Myanmar border area and other countries

In order to understand the distribution of diversity across geographically different populations, *Fst* values of the China–Myanmar border population and seven worldwide populations with full-length ectodomain sequences were evaluated. Pairwise population comparisons showed a high level of genetic differentiation ($Fst = 0.47$) between the South American population (Venezuela) with the China–Myanmar border population, and other worldwide populations (range from 0.23–0.35). A moderate range of *Fst* value (0.13–0.16) was detected

Table 3

Genetic differentiation (*Fst*) of the *PfAMA1* gene among 10 geographic different populations.

Locality (no.)	Myanmar	Thailand	PNG	Gambia	Nigeria	Kenya	Mali
Myanmar (135)							
Thailand (50)	0.14						
PNG (76)	0.02	0.05					
Gambia (114)	0.13	0.03	0.05				
Nigeria (51)	0.16	0.04	0.07	0.01			
Kenya (129)	0.15	0.03	0.07	0.01	0.01		
Mali (506)	0.15	0.02	0.06	0.00	0.01	0.01	
Venezuela (30)	0.47	0.27	0.35	0.25	0.23	0.26	0.25

Note: Myanmar, China–Myanmar border population; PNG, Papua New Guinea population.

when comparing the China–Myanmar border populations with the Thai and African populations, but a much lower genetic difference was revealed when the China–Myanmar border population was compared with the Oceania population ($Fst = 0.02$) (Table 3).

3.4. Positive diversifying selection on *PfAMA1* gene

Because of the observed high genetic diversity of *PfAMA1*, we evaluated signatures of selection on *PfAMA1* in the China–Myanmar border isolates (Table 1). A significant excess of nonsynonymous substitutions over synonymous substitutions was detected when the entire sequenced region was evaluated ($P < 0.001$). The same analysis performed against DI, DII, and DIII detected a significant excess of nonsynonymous substitutions over synonymous substitutions in DI, but not DII or DIII (Table 1). The results suggest that positive selection is acting on DI of *PfAMA1*. Tajima's *D* test shows positive values of *D* for the region as a whole and each domain separately (Table 1). Although not significant, sliding window plot depicted positive *D* values in DI and linker regions of each domain, suggesting natural selection in these regions (Fig. 3). Furthermore, Table 1 also shows significant values of Fu and Li's D^* (1.911) and F^* (1.929) for the entire sequenced region (Fu and Li's D^* ; $P < 0.02$ and F^* ; $P < 0.05$). When the three domains were separately assessed, significant deviations greater than zero were detected for DI ($D^* = 2.030$, $P < 0.02$ and $F^* = 1.964$; $P < 0.05$), but not DII or DIII. Sliding window plot analysis of Fu and Li's D^* and F^* also revealed a significant larger deviation than in DI (Fig. 3). Finally, the McDonald–Kreitman test showed a significant excess of intraspecific nonsynonymous substitutions over synonymous substitutions in the entire sequenced region and DI as compared with interspecies fixed differences of nonsynonymous and synonymous changes, suggesting positive selection ($P < 0.001$ and $P < 0.02$). Although not statistically significant, DII and III show the same trend as DI, suggesting that positive selection occurred across the ectodomain of *PfAMA1*. Collectively, all evolutionary tests for neutrality detected a signature of positive diversifying selection on the whole sequenced region and DI at the 98% confidence level.

3.5. Haplotype network reconstruction

A haplotype network was constructed to establish the relationships among the *PfAMA1* haplotypes from global *P. falciparum* populations. A total of 229 haplotypes were identified in 1091 sequences, of which 45.4% was singleton. Haplotype prevalence ranged from 0.4 to 38.9%. Haplotypes with sequences matching the *P. falciparum* vaccine strains 3D7 and FVO (GenBank accession no. AJ277646.1) were 9.6% and 7.8% respectively. Haplotypes 3, 5, 8, 15, 16 and 21 are shared between the China–Myanmar border population and African populations. Among which, haplotype 3 has an observed frequency of 38.9%, and it is the dominant haplotype of the China–Myanmar border population with a frequency of 60.4%. Haplotype 15 is the only haplotype shared among populations from the four continents (Asia, Africa, Oceania, and South America), with a frequency of 23.1% in total sequences analyzed. This haplotype occurred at 0.4% in the China–Myanmar border, 18.3% in

Fig. 2. Linkage disequilibrium (LD) across the *PfAMA1* gene of China–Myanmar border isolates was calculated by using the (A) R^2 and (B) D' index. Those pairs of sites that show significant linkage disequilibrium as calculated by Fisher's exact test is shown as red circles, while all others are shown by black circles. Trace line represents the regression line.

Fig. 3. Sliding window plot of Tajima's *D* test (A), Fu and Li's *D** (B) and *F** (C) tests for *PfAMA1* isolates from China–Myanmar border area. Nucleotide numbers are those of 3D7. Window length is 90 bp, and step size is 3 bp. The three domains of AMA1 are represented within the first plot as lines labeled Domain I, Domain II, and Domain III, respectively. Regions outside of the dash lines indicate the region with a significant departure from neutrality ($P < 0.05$, one-tailed). Regions outside the solid lines indicate regions with positive values ($P < 0.1$, one-tailed).

African, 2.2% in Oceanian, and 2.2% in South American populations, respectively. To obtain a haplotype network of major haplotypes, we excluded the 104 singletons and frequency $\leq 2\%$ haplotypes from analysis. When only the network torso is displayed, a clustered distribution of haplotypes correlated with the continents of origin and clearly separated the South American population from the rest of worldwide populations (Fig. 4).

4. Discussion

The current study analyzed the extent of genetic polymorphism in and diversifying selection on *PfAMA1* in the China–Myanmar border *P. falciparum* population. In this first attempt to determine *PfAMA1* diversity from this region, we detected high levels of genetic diversity in the *PfAMA1* ectodomain from 135 parasite isolates. This observation is comparable with results published on global isolates (Arnott et al., 2014; Ord et al., 2008; Osier et al., 2010; Polley et al., 2003; Polley and

Conway, 2001; Takala et al., 2009; Tetteh et al., 2009). Yet, the level of genetic diversity of *PfAMA1* from the China–Myanmar border population is lower than that from the African populations ($\pi = 0.026$ – 0.028) (Osier et al., 2010; Polley and Conway, 2001; Takala et al., 2009; Tetteh et al., 2009), the Southeast Asian population (Thai, $\pi = 0.025$ (Polley et al., 2003)), and the Oceania population (Papua New Guinea, $\pi = 0.026$ (Arnott et al., 2014)), but higher than that from the South American population ($\pi = 0.012$ (Ord et al., 2008)). The lower genetic diversity in *P. falciparum* population at the China–Myanmar border isolates is consistent with the epidemiologic characteristics of the study area, where malaria transmission is seasonal and unstable (Li et al., 2013) as compared to the holoendemic African regions. Though the number of analyzed sequences varied in parasite populations used in the comparison, the majority of mutation sites are shared among these populations, suggesting that similar selective forces (host immunity) act in various geographical regions.

Recombination and positive selection are possibly two major factors responsible for the evolution and genetic variation in *PfAMA1* (Eisen et al., 1999). In several populations, recombination is frequently observed, as shown by a high estimate of the recombination parameter, *C*, and very rapid decline in linkage disequilibrium with increasing distance between nucleotide sites (Osier et al., 2010; Polley et al., 2003; Polley and Conway, 2001; Takala et al., 2009; Tetteh et al., 2009). Though higher than that in the South American population (Ord et al., 2008), the estimated recombinant parameters of our study population are much lower than those of the African populations (Osier et al., 2010; Polley and Conway, 2001; Takala et al., 2009; Tetteh et al., 2009). Within Southeast Asia, the recombination rate of the China–Myanmar border population is lower than that in the Thai–Myanmar border population (Polley et al., 2003). Considering the average recombination rate in the *P. falciparum* genome as 1 cM per 17 kb of sequence (Su et al., 1999), which equates to $r = 6 \times 10^{-7}$, the estimated genetically effective population size *N* in the current study would be 1.3×10^3 , which is much lower than the estimate for African populations at $\sim 6.6 \times 10^4$ (Polley et al., 2003). This again may be due to the lower malaria endemicity in the China–Myanmar border area, which restricts the opportunity of multiclonal infections, subsequent cross-fertilization, and recombination in mosquitoes (Walliker, 2000). Our recent epidemiological studies at the international borders of the GMS showed much lower *P. falciparum* endemicity at the China–Myanmar border than at the Thai–Myanmar border (Baum et al., 2015; Zhou et al., 2014). We also obtained evidence indicating limited genetic differentiation of *PfAMA1* (overall *F_{st}* value 0.09) among parasite populations. However, the *F_{st}* value between the China–Myanmar border and the Venezuelan population (*F_{st}* = 0.47) showed evidence of strong population division (Ord et al., 2008). In addition, *F_{st}* value for the China–Myanmar border population in comparison with other global populations showed moderate levels of genetic differentiation, which is largely due to geographic area-specific SNPs (*F_{st}* = 0.13–0.16 (Osier et al., 2010; Polley et al., 2003; Polley and Conway, 2001; Takala et al., 2009; Tetteh et al., 2009)). However, it is intriguing that the China–Myanmar border population showed little genetic differentiation from the Papua New Guinea population (Arnott et al., 2014). While this may result from similar host immune selection forces imposed on *PfAMA1* in these geographically separated parasite populations, it nonetheless raised hope that the final *PfAMA1*-based vaccine design targeting a limited number of *PfAMA1*-haplotypes may be effective for different endemic regions (Drew et al., 2012).

The significant ratio of nonsynonymous to synonymous substitutions (d_N/d_S) observed in our study and the other 7 *P. falciparum* populations suggest that amino acid replacements are generally favored in *PfAMA1* and this adaptive evolution is presumably due to host immune pressure. Additional neutrality tests provided corroborating findings indicative of positive diversifying selection operating at *PfAMA1* DI and possibly the entire ectodomain. Though the value of Tajima's *D* test

In conclusion, our study presented baseline data of the *PfAMA1* polymorphism in 135 *P. falciparum* isolates from the China–Myanmar border area. Understanding polymorphism and host immune response is an important aspect for malaria vaccine development in

Arnott, A., Wapling, J., Mueller, I., Ramsland, P.A., Siba, P.M., Reeder, J.C., Barry, A.E., 2014. Distinct patterns of diversity, population structure and evolution in the AMA1 genes of sympatric *Plasmodium falciparum* and *Plasmodium vivax* populations of Papua New Guinea from an area of similarly high transmission. *Malar. J.* 13, 233.

Bandelt, H.J., Forster, P., Rohlf, A., 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* 16, 37–48.

Bargieri, D., Lagal, V., Tardieux, I., Menard, R., 2012. Host cell invasion by apicomplexans: what do we know? *Trends Parasitol.* 28, 131–135.

Baum, E., Sattabongkot, J., Sirichaisinthop, J., Kiattibutr, K., Davies, D.H., Jain, A., Lo, E., Lee, M.C., Randall, A.Z., Molina, D.M., Liang, X., Cui, L., Felgner, P.L., Yan, G., 2015. Submicroscopic and asymptomatic *Plasmodium falciparum* and *Plasmodium vivax* infections are common in western Thailand – molecular and serological evidence. *Malar. J.* 14, 95.

Cao, J., Kaneko, O., Thongkukiattkul, A., Tachibana, M., Otsuki, H., Gao, Q., Tsuboi, T., Torii, M., 2009. Rhoptry neck protein RON2 forms a complex with microneme protein AMA1 in *Plasmodium falciparum* merozoites. *Parasitol. Int.* 58, 29–35.

Chesne-Seck, M.L., Pizarro, J.C., Vulliez-le Normand, B., Collins, R.A., Blackman, M.J., Faber, B.W., Remarque, E.J., Kocken, C.H., Thomas, A.W., Bentley, G.A., 2005. Structural comparison of apical membrane antigen 1 orthologues and paralogues in apicomplexan parasites. *Mol. Biochem. Parasitol.* 144, 55–67.

Cortes, A., Mellombo, M., Mueller, I., Benet, A., Reeder, J.C., Anders, R.F., 2003. Geographical structure of diversity and differences between symptomatic and asymptomatic infections for *Plasmodium falciparum* vaccine candidate AMA1. *Infect. Immun.* 71, 1416–1426.

Crewther, P.E., Matthew, M.L., Flegg, R.H., Anders, R.F., 1996. Protective immune responses to apical membrane antigen 1 of *Plasmodium chabaudi* involve recognition of strain-specific epitopes. *Infect. Immun.* 64, 3310–3317.

Cui, L., Yan, G., Sattabongkot, J., Cao, Y., Chen, B., Chen, X., Fan, Q., Fang, Q., Jongwutiwes, S., Parker, D., Sirichaisinthop, I., Kwaw, M.P., Su, X.Z., Yang, H., Yang, Z., Wang, B., Xu, L.

- Zheng, B., Zhong, D., Zhou, G., 2012. Malaria in the Greater Mekong Subregion: heterogeneity and complexity. *Acta Trop.* 121, 227–239.
- Dondorp, A.M., Fairhurst, R.M., Slutsker, L., Macarthur, J.R., Breman, J.G., Guerin, P.J., Wellems, T.E., Ringwald, P., Newman, R.D., Plowe, C.V., 2011. The threat of artemisinin-resistant malaria. *N. Engl. J. Med.* 365, 1073–1075.
- Drew, D.R., Hodder, A.N., Wilson, D.W., Foley, M., Mueller, I., Siba, P.M., Dent, A.E., Cowman, A.F., Beeson, J.G., 2012. Defining the antigenic diversity of *Plasmodium falciparum* apical membrane antigen 1 and the requirements for a multi-allele vaccine against malaria. *PLoS ONE* 7, e51023.
- Dutta, S., Haynes, J.D., Moch, J.K., Barbosa, A., Lanar, D.E., 2003. Invasion-inhibitory antibodies inhibit proteolytic processing of apical membrane antigen 1 of *Plasmodium falciparum* merozoites. *Proc. Natl. Acad. Sci. U. S. A.* 100, 12295–12300.
- Eisen, D.P., Marshall, V.M., Billman-Jacobe, H., Coppel, R.L., 1999. A *Plasmodium falciparum* apical membrane antigen-1 (AMA-1) gene apparently generated by intragenic recombination. *Mol. Biochem. Parasitol.* 100, 243–246.
- Ellis, R.D., Wu, Y., Martin, L.B., Shaffer, D., Miura, K., Aebig, J., Orcutt, A., Rausch, K., Zhu, D., Mogensen, A., Fay, M.P., Narum, D.L., Long, C., Miller, L., Durbin, A.P., 2012. Phase 1 study in malaria naive adults of BSAM2/Alhydrogel(R) + CPG 7909, a blood stage vaccine against *P. falciparum* malaria. *PLoS ONE* 7, e46094.
- Figtree, M., Pasay, C.J., Slade, R., Cheng, Q., Cloonan, N., Walker, J., Saul, A., 2000. *Plasmodium vivax* synonymous substitution frequencies, evolution and population structure deduced from diversity in AMA 1 and MSP 1 genes. *Mol. Biochem. Parasitol.* 108, 53–66.
- Fowkes, F.J., Richards, J.S., Simpson, J.A., Beeson, J.G., 2010. The relationship between anti-merozoite antibodies and incidence of *Plasmodium falciparum* malaria: a systematic review and meta-analysis. *PLoS Med.* 7, e1000218.
- Fu, Y.X., Li, W.H., 1993. Statistical tests of neutrality of mutations. *Genetics* 133, 693–709.
- Garg, S., Alam, M.T., Das, M.K., Dev, V., Kumar, A., Dash, A.P., Sharma, Y.D., 2007. Sequence diversity and natural selection at domain I of the apical membrane antigen 1 among Indian *Plasmodium falciparum* populations. *Malar. J.* 6, 154.
- Healer, J., Crawford, S., Ralph, S., McFadden, G., Cowman, A.F., 2002. Independent translocation of two micronemal proteins in developing *Plasmodium falciparum* merozoites. *Infect. Immun.* 70, 5751–5758.
- Healer, J., Murphy, V., Hodder, A.N., Masciantonio, R., Gemmill, A.W., Anders, R.F., Cowman, A.F., Batchelor, A., 2004. Allelic polymorphisms in apical membrane antigen-1 are responsible for evasion of antibody-mediated inhibition in *Plasmodium falciparum*. *Mol. Microbiol.* 52, 159–168.
- Hill, W.G., Robertson, A., 1968. Linkage disequilibrium in finite populations. *TAG. Theor. Appl. Genet.* 38, 226–231.
- Hodder, A.N., Crewther, P.E., Anders, R.F., 2001. Specificity of the protective antibody response to apical membrane antigen 1. *Infect. Immun.* 69, 3286–3294.
- Hodder, A.N., Crewther, P.E., Matthew, M.L., Reid, G.E., Moritz, R.L., Simpson, R.J., Anders, R.F., 1996. The disulfide bond structure of *Plasmodium* apical membrane antigen-1. *J. Biol. Chem.* 271, 29446–29452.
- Hudson, R.R., 1987. Estimating the recombination parameter of a finite population model without selection. *Genet. Res.* 50, 245–250.
- Hudson, R.R., Kaplan, N.L., 1985. Statistical properties of the number of recombination events in the history of a sample of DNA sequences. *Genetics* 111, 147–164.
- Jordan, S.J., Branch, O.H., Castro, J.C., Oster, R.A., Rayner, J.C., 2009. Genetic diversity of the malaria vaccine candidate *Plasmodium falciparum* merozoite surface protein-3 in a hypoendemic transmission environment. *Am. J. Trop. Med. Hyg.* 80, 479–486.
- Kennedy, M.C., Wang, J., Zhang, Y., Miles, A.P., Chitsaz, F., Saul, A., Long, C.A., Miller, L.H., Stowers, A.W., 2002. In vitro studies with recombinant *Plasmodium falciparum* apical membrane antigen 1 (AMA1): production and activity of an AMA1 vaccine and generation of a multiallelic response. *Infect. Immun.* 70, 6948–6960.
- Kocken, C.H., Narum, D.L., Massougbodji, A., Ayivi, B., Dubbeld, M.A., van der Wel, A., Conway, D.J., Sanni, A., Thomas, A.W., 2000. Molecular characterisation of *Plasmodium reichenowi* apical membrane antigen-1 (AMA-1), comparison with *P. falciparum* AMA-1, and antibody-mediated inhibition of red cell invasion. *Mol. Biochem. Parasitol.* 109, 147–156.
- Kocken, C.H., Withers-Martinez, C., Dubbeld, M.A., van der Wel, A., Hackett, F., Valderrama, A., Blackman, M.J., Thomas, A.W., 2002. High-level expression of the malaria blood-stage vaccine candidate *Plasmodium falciparum* apical membrane antigen 1 and induction of antibodies that inhibit erythrocyte invasion. *Infect. Immun.* 70, 4471–4476.
- Koukouikila-Koussounda, F., Malonga, V., Mayengue, P.I., Ndounga, M., Vouvougui, C.J., Ntouni, F., 2012. Genetic polymorphism of merozoite surface protein 2 and prevalence of K76T pfCRT mutation in *Plasmodium falciparum* field isolates from Congolese children with asymptomatic infections. *Malar. J.* 11, 105.
- Kusi, K.A., Faber, B.W., Thomas, A.W., Remarque, E.J., 2009. Humoral immune response to mixed PfAMA1 alleles; multivalent PfAMA1 vaccines induce broad specificity. *PLoS ONE* 4, e8110.
- Laurens, M.B., Thera, M.A., Coulibaly, D., Ouattara, A., Kone, A.K., Guindo, A.B., Traore, K., Traore, I., Kouriba, B., Diallo, D.A., Diarra, I., Daou, M., Dolo, A., Tolo, Y., Sissoko, M.S., Niangaly, A., Sissoko, M., Takala-Harrison, S., Lyke, K.E., Wu, Y., Blackwelder, W.C., Godeaux, O., Vekemans, J., Dubois, M.C., Ballou, W.R., Cohen, J., Dube, T., Soisson, L., Diggs, C.L., House, B., Bennett, J.W., Lanar, D.E., Dutta, S., Heppner, D.G., Plowe, C.V., Doumbo, O.K., 2013. Extended safety, immunogenicity and efficacy of a blood-stage malaria vaccine in malian children: 24-month follow-up of a randomized, double-blinded phase 2 trial. *PLoS ONE* 8, e79323.
- Lewontin, R.C., 1964. The interaction of selection and linkage. I. General considerations; heterotic models. *Genetics* 49, 49–67.
- Li, N., Parker, D.M., Yang, Z., Fan, Q., Zhou, G., Ai, G., Duan, J., Lee, M.C., Yan, G., Matthews, S.A., Cui, L., Wang, Y., 2013. Risk factors associated with slide positivity among febrile patients in a conflict zone of north-eastern Myanmar along the China–Myanmar border. *Malar. J.* 12, 361.
- Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451–1452.
- McDonald, J.H., Kreitman, M., 1991. Adaptive protein evolution at the Adh locus in *Drosophila*. *Nature* 351, 652–654.
- Meng, H., Zhang, R., Yang, H., Fan, Q., Su, X., Miao, J., Cui, L., Yang, Z., 2010. In vitro sensitivity of *Plasmodium falciparum* clinical isolates from the China–Myanmar border area to quinine and association with polymorphism in the Na⁺/H⁺ exchanger. *Antimicrob. Agents Chemother.* 54, 4306–4313.
- Mitchell, G.H., Thomas, A.W., Margos, G., Dluzewski, A.R., Bannister, L.H., 2004. Apical membrane antigen 1, a major malaria vaccine candidate, mediates the close attachment of invasive merozoites to host red blood cells. *Infect. Immun.* 72, 154–158.
- Narum, D.L., Thomas, A.W., 1994. Differential localization of full-length and processed forms of PF83/AMA-1 an apical membrane antigen of *Plasmodium falciparum* merozoites. *Mol. Biochem. Parasitol.* 67, 59–68.
- Nei, M., Gojobori, T., 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* 3, 418–426.
- Ord, R.L., Tami, A., Sutherland, C.J., 2008. Ama1 genes of sympatric *Plasmodium vivax* and *P. falciparum* from Venezuela differ significantly in genetic diversity and recombination frequency. *PLoS ONE* 3, e3366.
- Osier, F.H., Weedall, G.D., Verra, F., Murungi, L., Tetteh, K.K., Bull, P., Faber, B.W., Remarque, E., Thomas, A., Marsh, K., Conway, D.J., 2010. Allelic diversity and naturally acquired allele-specific antibody responses to *Plasmodium falciparum* apical membrane antigen 1 in Kenya. *Infect. Immun.* 78, 4625–4633.
- Peterson, M.G., Marshall, V.M., Smythe, J.A., Crewther, P.E., Lew, A., Silva, A., Anders, R.F., Kemp, D.J., 1989. Integral membrane protein located in the apical complex of *Plasmodium falciparum*. *Mol. Cell. Biol.* 9, 3151–3154.
- Polley, S.D., Conway, D.J., 2001. Strong diversifying selection on domains of the *Plasmodium falciparum* apical membrane antigen 1 gene. *Genetics* 158, 1505–1512.
- Polley, S.D., Chokejindachai, W., Conway, D.J., 2003. Allele frequency-based analyses robustly map sequence sites under balancing selection in a malaria vaccine candidate antigen. *Genetics* 165, 555–561.
- Remarque, E.J., Faber, B.W., Kocken, C.H., Thomas, A.W., 2008. Apical membrane antigen 1: a malaria vaccine candidate in review. *Trends Parasitol.* 24, 74–84.
- Richard, D., MacRaild, C.A., Riglar, D.T., Chan, J.A., Foley, M., Baum, J., Ralph, S.A., Norton, R.S., Cowman, A.F., 2010. Interaction between *Plasmodium falciparum* apical membrane antigen 1 and the rhoptry neck protein complex defines a key step in the erythrocyte invasion process of malaria parasites. *J. Biol. Chem.* 285, 14815–14822.
- Sakihama, N., Kaneko, A., Hattori, T., Tanabe, K., 2001. Limited recombination events in merozoite surface protein-1 alleles of *Plasmodium falciparum* on islands. *Gene* 279, 41–48.
- Silvie, O., Franetich, J.F., Charrin, S., Mueller, M.S., Siau, A., Bodescot, M., Rubinstein, E., Hannoun, L., Charoenvit, Y., Kocken, C.H., Thomas, A.W., Van Gemert, G.J., Sauerwein, R.W., Blackman, M.J., Anders, R.F., Pluschke, G., Mazier, D., 2004. A role for apical membrane antigen 1 during invasion of hepatocytes by *Plasmodium falciparum* sporozoites. *J. Biol. Chem.* 279, 9490–9496.
- Su, X., Ferdig, M.T., Huang, Y., Huynh, C.Q., Liu, A., You, J., Wootton, J.C., Wellems, T.E., 1999. A genetic map and recombination parameters of the human malaria parasite *Plasmodium falciparum*. *Science* 286, 1351–1353.
- Tajima, F., 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123, 585–595.
- Takala, S.L., Coulibaly, D., Thera, M.A., Batchelor, A.H., Cummings, M.P., Escalante, A.A., Ouattara, A., Traore, K., Niangaly, A., Djimde, A.A., Doumbo, O.K., Plowe, C.V., 2009. Extreme polymorphism in a vaccine antigen and risk of clinical malaria: implications for vaccine development. *Sci. Transl. Med.* 1, 2ra5.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729.
- Terheggen, U., Drew, D.R., Hodder, A.N., Cross, N.J., Mugenyi, C.K., Barry, A.E., Anders, R.F., Dutta, S., Osier, F.H., Elliott, S.R., Senn, N., Stanisic, D.I., Marsh, K., Siba, P.M., Mueller, I., Richards, J.S., Beeson, J.G., 2014. Limited antigenic diversity of *Plasmodium falciparum* apical membrane antigen 1 supports the development of effective multi-allele vaccines. *BMC Med.* 12, 183.
- Tetteh, K.K., Stewart, L.B., Ochola, L.I., Amambua-Ngwa, A., Thomas, A.W., Marsh, K., Weedall, G.D., Conway, D.J., 2009. Prospective identification of malaria parasite genes under balancing selection. *PLoS ONE* 4, e5568.
- Thakur, A., Alam, M.T., Bora, H., Kaur, P., Sharma, Y.D., 2008. *Plasmodium vivax*: sequence polymorphism and effect of natural selection at apical membrane antigen 1 (PvAMA1) among Indian population. *Gene* 419, 35–42.
- Thera, M.A., Doumbo, O.K., Coulibaly, D., Laurens, M.B., Kone, A.K., Guindo, A.B., Traore, K., Sissoko, M., Diallo, D.A., Diarra, I., Kouriba, B., Daou, M., Dolo, A., Baby, M., Sissoko, M.S., Sagara, I., Niangaly, A., Traore, I., Olotu, A., Godeaux, O., Leach, A., Dubois, M.C., Ballou, W.R., Cohen, J., Thompson, D., Dube, T., Soisson, L., Diggs, C.L., Takala, S.L., Lyke, K.E., House, B., Lanar, D.E., Dutta, S., Heppner, D.G., Plowe, C.V., 2010. Safety and immunogenicity of an AMA1 malaria vaccine in Malian children: results of a phase 1 randomized controlled trial. *PLoS ONE* 5, e9041.
- Thera, M.A., Doumbo, O.K., Coulibaly, D., Laurens, M.B., Ouattara, A., Kone, A.K., Guindo, A.B., Traore, K., Traore, I., Kouriba, B., Diallo, D.A., Diarra, I., Daou, M., Dolo, A., Tolo, Y., Sissoko, M.S., Niangaly, A., Sissoko, M., Takala-Harrison, S., Lyke, K.E., Wu, Y., Blackwelder, W.C., Godeaux, O., Vekemans, J., Dubois, M.C., Ballou, W.R., Cohen, J., Thompson, D., Dube, T., Soisson, L., Diggs, C.L., House, B., Lanar, D.E., Dutta, S., Heppner Jr., D.G., Plowe, C.V., 2011. A field trial to assess a blood-stage malaria vaccine. *N. Engl. J. Med.* 365, 1004–1013.
- Thomas, A.W., Trape, J.F., Rogier, C., Gonçalves, A., Rosario, V.E., Narum, D.L., 1994. High prevalence of natural antibodies against *Plasmodium falciparum* 83-kilodalton apical membrane antigen (PF83/AMA-1) as detected by capture-enzyme-linked

- immunosorbent assay using full-length baculovirus recombinant PF83/AMA-1. *Am.J.Trop. Med. Hyg.* 51, 730–740.
- Walliker, D., 2000. Malaria.
- Wang, Y., Zhong, D., Cui, L., Lee, M.C., Yang, Z., Yan, G., Zhou, G., 2015. Population dynamics and community structure of anopheles mosquitoes along the China–Myanmar border. *Parasit. Vectors* 8, 445.
- WHO, 2014. World Malaria Report 2014. World Health Organ. 242.
- Zhou, G., Sun, L., Xia, R., Duan, Y., Xu, J., Yang, H., Wang, Y., Lee, M.C., Xiang, Z., Yan, G., Cui, L., Yang, Z., 2014. Clinical malaria along the China–Myanmar border, Yunnan Province, China, January 2011–August 2012. *Emerg. Infect. Dis.* 20, 675–678.