

Lawrence Berkeley National Laboratory

LBL Publications

Title

The Ion-Exchange Separation of Zirconium and Hafnium

Permalink

<https://escholarship.org/uc/item/9bz24489>

Authors

Street, Kenneth, Jr.

Seaborg, Glenn T

Publication Date

1943-10-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-191
0.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

~~CONFIDENTIAL~~
~~CONFIDENTIAL~~

CLASSIFICATION CANCELLED BY AUTHORITY
OF THE DISTRICT ENGINEER
BY THE DECLASSIFICATION COMMITTEE
UNIVERSITY OF CALIFORNIA

Radiation Laboratory

THE ION-EXCHANGE SEPARATION OF ZIRCONIUM AND HAFNIUM

Kenneth Street, Jr. and Glenn T. Seaborg

October 11, 1943

Special Review of Declassified Reports
Authorized by USDOE JX E-1001
Unclassified TWX P182206Z May 79

REPORT PROPERLY DECLASSIFIED

<u>J N Green</u>	<u>8/16/79</u>
Authorized Derivative Classifier	Date
<u>R K Hunt</u>	<u>8/17/79</u>
By	Date

Berkeley, California

STANDARD DISTRIBUTION: Series A

Argonne National Laboratory	9
Armed Forces Special Weapons Project	10-11
Atomic Energy Commission, Washington	12
Battelle Memorial Institute	13-22
Brookhaven National Laboratory	23-26
Carbide & Carbon Chemicals Corporation (K-25 Area)	27-30
Carbide & Carbon Chemicals Corporation (Y-12 Area)	31
Columbia University (Failla)	32-35
General Electric Company	36-42
Hanford Directed Operations	43
Iowa State College	44-45
Kellex Corporation	46-48
Los Alamos	49
Massachusetts Institute of Technology	50-51
Monsanto Chemical Company, Dayton	52-53
National Bureau of Standards	54
Naval Radiological Defense Laboratory	55
NEPA	56-57
New York Directed Operations	58-69
Oak Ridge National Laboratory	70
Patent Advisor	71-85
Technical Information Division	86
UCLA Medical Research Laboratory (Warren)	87-89
University of California, Radiation Laboratory	90
Information Division	91-92
Chemistry Dept.	93
University of Rochester	94
Western Reserve University (Friedell)	
Office of Chicago Directed Operations	

DECLASSIFICATION PROCEDURE:

Declassification Officer	95-98
Publication Officer	99
Patent Department	100-101
Area Manager	102
E. O. Lawrence	103
Information Div.	104

CLASSIFICATION CONTROLLED BY AUTHORITY
 OF THE DISTRICT ENGINEER
 BY THE DECLASSIFICATION COMMITTEE

The Ion-Exchange Separation of Zirconium and Hafnium

Kenneth Street, Jr. and Glenn T. Seaborg
Department of Chemistry and Radiation Laboratory
University of California, Berkeley, California

CLASSIFICATION CANCELLED BY AUTHORITY
OF DISTRICT ENGINEER
CALIFORNIA CLASSIFICATION COMMITTEE

October 11, 1948

In the course of a rather cursory examination of the elution of tetra-positive ions from the cation exchange resin Dowex 50 with hydrochloric acid solutions, we have discovered a very effective method of separating zirconium from hafnium. In view of the great labor involved in preparing even reasonably pure hafnium compounds by existing methods, we feel that this procedure will prove very valuable to those interested in obtaining hafnium compounds free of zirconium.

Although the conditions which give satisfactory separation were first worked out using microgram amounts of material and the radioactive tracer technique, the run described below, involving milligrams of material, illustrates the applicability of the method to the production of significant amounts of pure hafnium and zirconium.

35 mg. of zirconium oxide 15 mg. of hafnium oxide were dissolved in sulfuric and hydrofluoric acids, hafnium and zirconium tracer added, and the mixture fumed to dryness. The residue was taken up in concentrated hydrochloric acid and the hydroxides precipitated with ammonium hydroxide and washed. The hydroxides were again dissolved in hydrochloric acid and the oxychlorides crystallized by evaporation. 1 cc. of 250 to 500 mesh Dowex 50 spheres, in the ammonium form, were suspended in 30 cc. of 2 molal perchloric acid and the oxychlorides added a few mg. at a time over a period of 15 minutes, the mixture being continually agitated by bubbling

air through it. Under these conditions, i.e. $\sim 0.01M$ zirconium and hafnium in 2M perchloric acid, the zirconium and hafnium are not appreciably polymerized¹ and about 80 percent of each goes on the resin. The slurry of resin was placed on the top of an ion-exchange column 1 sq. cm. in area and 30 cm. long which had been packed with the same resin and washed with 6 molal hydrochloric acid to convert it to the acid form. On elution with 6 molal hydrochloric acid the curve shown in Fig. 1 was obtained. The outline of the curve was obtained by counting the tracers (Zr^{95} and Hf^{181}) and the dotted portions by optical spectrographic analysis.

Thus it can be seen that ~ 66 percent of the starting hafnium oxide, i.e. 10 mg., is obtained containing ~ 0.1 percent zirconium oxide by weight. The column used in these experiments was relatively small and thus gram amounts of material should be easily handled on columns of only moderate size.

This work was done under the auspices of the Atomic Energy Commission at the Radiation Laboratory, University of California, Berkeley, California.

1. Connick, R. E., and W. H. McVey, Private Communication.




FIG. 1
ELUTION OF Zr AND Hf WITH 6.0M HCl

CLASSIFICATION
OF THE DISTRICT
BY THE DEPARTMENT

CLASSIFICATION CANCELLED
OF THE DISTRICT
BY THE DECLASSIFICATION COMMITTEE

RESTRICTED