

Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

Evidence for Subshell at $Z = 96$

Permalink

<https://escholarship.org/uc/item/9d65b7hh>

Author

Seaborg, Glenn T.

Publication Date

1953-09-03

UCRL-2335

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

EVIDENCE FOR SUBSHELL AT $Z = 96$

Glenn T. Seaborg

September 3, 1953

Berkeley, California

EVIDENCE FOR SUBSHELL AT $Z = 96$

Glenn T. Seaborg
 Radiation Laboratory and Department of Chemistry
 University of California, Berkeley, California

September 3, 1953

The evidence is decisive that major nuclear shells are completed at 82 protons and 126 neutrons (both represented by the nuclide Pb^{208}) and these, along with major shells at 82 neutrons and certain lower nucleon numbers (N or $Z = 20, 28, 50$), are well explained by the strong spin orbit coupling model of Mayer¹ and Haxel, Jensen, and Suess.² This model suggests the filling of quantum states at certain intermediate points, and there is an accumulating amount of evidence that such "subshells" are also discernible, for example, at $Z = 58$ ^{3, 4, 5} and $Z = 64$.^{6, 7}

The evidence from alpha radioactivity, both (1) the effect of the nuclear radius shrinkage on the relationship between energy and half-life and (2) the discontinuities in the plots of energy vs. mass number at constant Z , gives a striking indication⁸ of the closing of major shells at $Z = 82$ and $N = 126$. Application of these sensitive criteria as tests for the much smaller "subshell" effects in the regions $Z > 82$ and $N > 126$ leads to some evidence for such a subshell at $Z = 96$ (curium).

Since it has been shown that the Gamow-Gurney-Condon type of formula for alpha decay applies very well to the ground state to ground state transition for even-even alpha emitters,⁸ the known⁹ alpha energies and partial half-lives were used in the Preston¹⁰ form of the formula to calculate the nuclear radii of a number of even-even nuclides in the range $Z = \overset{84}{\cancel{96}}$ to $\overset{98}{\cancel{96}}$. Using these radii a value of r_0 was

calculated for each nuclide from the relationship $\text{radius} = r_0 A^{1/3}$ and the plot in Fig. 1 of the average value of r_0 (r_0 generally decreasing with increasing Z) for each element indicates a just discernible minimum or plateau at $Z = 96$. The average value of r_0 for each element was plotted because there is no discernible regular variation of r_0 with A at constant Z .

The stable shell of 82 protons is attained upon completion of the $h_{11/2}$ level and the spin of Bi^{209} ($9/2$) indicates that the 83rd proton begins the filling of the $h_{9/2}$ level as might be expected. However, if the $h_{9/2}$ level is raised in energy as more protons are added, so that the $f_{7/2}$ and $f_{5/2}$ are filled before the $h_{9/2}$ levels, one might expect subshell effects at $Z = 90$ and 96 . If the quantum states are filled in this order, the variation of r_0 with Z should perhaps also indicate an effect $Z = 90$. A careful consideration of the values of r_0 in the region on both sides of $Z = 90$ points to a barely discernible plateau in the variation of r_0 with Z at this atomic number.

In the case of $Z = 96$ there is an additional argument which points to the completion of a subshell here. The known odd-even isotopes of berkelium, Bk^{243} and Bk^{245} , are highly "hindered"⁸ in their most prominent modes of alpha decay, i. e., they decay much slower than the simple formula would indicate. This exceptional degree of hindrance is not observed for similar (odd-even) isotopes of any other odd Z alpha particle emitter with the exception of bismuth ($Z = 83$) where the slowed rates of decay are presumably to be associated with the closed proton shell (and consequent shrunken radius) at $Z = 82$.

There are other lines of evidence which may also point to the filling of the $f_{7/2}$ and $f_{5/2}$ before the $h_{9/2}$ proton states. The spins of Np^{237}_{11} and Am^{241}_{12} are both $5/2$ as expected on this basis. On the other hand, the spins of Ac^{227}_{13} and Pa^{231}_{14} have both been reported as $3/2$ indicating perhaps that the odd proton occupies the $p_{3/2}$ state, whereas spins of $7/2$ and $5/2$ corresponding to $f_{7/2}$ and $f_{5/2}$ states, respectively, would be expected. Whether or not this indicates a breakdown of the single particle model it does seem to indicate that arguments based on spin values cannot be conclusive here. It is interesting to add that a consideration of the systematics of beta radioactivity in this region also leads to the assignment of spectroscopic states in agreement with the suggested higher position of the $h_{9/2}$ level energetically.

It is interesting to note that arguments based on spin values¹⁵ indicate that in the case of neutrons the $f_{7/2}$ level fills before the $h_{9/2}$ level just after the completion of the major shell at 82 neutrons. Thus, the situation is analogous to that postulated for protons although the evidence is not clear on the relative position of the $f_{5/2}$ and $h_{9/2}$ neutron levels.

It seems likely that in the region following the major closed shell at 126 neutrons, which is completed with the filling of the $i_{13/2}$ states, the $i_{11/2}$ level is not occupied until the $g_{9/2}$ and possibly also the $g_{7/2}$ and $d_{5/2}$, etc., states are filled. Attempts to apply the above mentioned criteria of alpha decay data do not lead to any discernible evidence for subshells up to $N = 148$. Successful application of these criteria here probably awaits both a larger quantity of and

more accurate alpha decay data in this region. There is only one spin value known in this region, that of U^{235} , which has been reported as $5/2$ or $7/2$ with the former value the more probable.¹⁶ Since U^{235} has 143 neutrons either spin value would fit nicely with the view that the $i_{11/2}$ lies above the $g_{9/2}$, $g_{7/2}$, and $d_{5/2}$ levels.

The generous help of F. Asaro and D. C. Dunlavey in making numerous calculations is gratefully acknowledged.

FIGURE CAPTION

Figure 1. Plot of average values of r vs. Z . Following isotopes included: Po^{212} , Po^{214} , Po^{216} , Po^{218} for Po; Em^{220} , Em^{222} for Em; Ra^{222} , Ra^{224} , Ra^{226} for Ra; Th^{226} , Th^{228} , Th^{230} , Th^{232} for Th; U^{230} , U^{232} , U^{234} , U^{236} , U^{238} for U; Pu^{236} , Pu^{238} , Pu^{240} , Pu^{242} for Pu; Cm^{242} , Cm^{244} for Cm; Cf^{246} , Cf^{248} for Cf.

REFERENCES

1. M. G. Mayer, Phys. Rev. 75, 1969 (1949); *ibid.*, 78, 16,22 (1950).
2. Haxel, Jensen, and Suess, Phys. Rev. 75, 1766 (1949);
ibid., Z. Physik 128, 295 (1950).
3. H. E. Duckworth and R. S. Preston, Phys. Rev. 82, 468 (1951).
4. Duckworth, Kegley, Olson, and Stanford, Phys. Rev. 83, 1114
(1941).
5. G. P. Dube and S. Jha, Phys. Rev. 85, 1042 (1952).
6. Coryell, Brightsen, and Pappas, Phys. Rev. 85, 732 (1952);
Coryell, Ann. Rev. Nuc. Sci. II, 305 (1953).
7. Rasmussen, Thompson, and Ghiorso, Phys. Rev. 89, 33 (1953).
8. Perlman, Ghiorso, and Seaborg, Phys. Rev. 77, 26 (1950).
9. Hollander, Perlman, and Seaborg, Revs. Modern Phys. 25,
469 (1953); additional data: Cf²⁴⁸, partial half-life 260 d,
energy 6.26 Mev (E. K. Hulet, Ph.D. thesis, University of
California, UCRL-2283, August 1953); U²³⁰, energy 5.886 Mev,
Th²²⁶, energy 6.342 Mev, Ra²²², energy 6.567 Mev, U²³²,
energy 5.318 Mev (F. Asaro, private communication); U²³⁸,
energy 4.187 Mev, Th²³², energy 3.998 Mev (B. G. Harvey,
private communication); Cf²⁴⁶, partial half-life 40.6 hrs
(D. C. Dunlavey, private communication).
10. M. Preston, Phys. Rev. 71, 865 (1947).
11. F. S. Tompkins and M. Fred, J. Opt. Soc. Amer. 39, 357
(1949).
12. M. Fred and F. S. Tompkins, Phys. Rev. 89, 318 (1953).
13. Tompkins, Fred, and Meggers, Phys. Rev. 84, 168 (1951).
14. J. E. Mack, Revs. Modern Phys. 22, 64 (1950).

15. P. F. A. Klinkenberg, *Revs. Modern Phys.* 24, 63 (1952).
16. G. L. Stukenbroeker and J. R. McNally, *J. Opt. Soc. Amer.* 40, 336 (1950).

