UC Irvine

UC Irvine Previously Published Works

Title

Effects of fungicides and bactericides on orchid seed germination and shoot tip cultures in vitro

Permalink

<https://escholarship.org/uc/item/9f26x5vr>

Journal

Plant Cell, Tissue and Organ Culture, 1(1)

ISSN

0167-6857

Authors

Brown, DM
Groom, CL
Cvitanik, M
et al.

Publication Date

1981

DOI

10.1007/BF02318914

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at
<https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Effects of fungicides and bactericides on orchid seed germination and shoot tip cultures in vitro

D.M. BROWN, C.L. GROOM, M.CVITANIK, M. BROWN, J.L. COOPER, and J. ARDITTI

Developmental and Cell Biology, University of California, Irvine, California 92717, USA

(Received 15 January 1982; in revised form 23 April 1982)

Key words: antibiotics, bactericides, contamination, fungicides, in vitro, orchids, shoot tip, seed germination, tissue culture

Abstract. Amphotericin B, benomyl, gentamycin, nystatin, quintozone penicillin G, sodium omadine, and vancomycin singly and in several combinations have no deleterious effects on the germination of orchid seeds, but inhibit the growth in vitro of shoot tip explants.

Introduction

Contamination by fungi and bacteria from the air, water, tools, glassware, and other sources may occur in orchid seedling and tissue cultures. Surface sterilization of the seeds or tissues can prevent initial contamination, but microorganisms often find their way into cultures at a later date. Incorporation of amphotericin B, benomyl (as the pure compound or the commercial preparation Benlate), gentamycin, nystatin (Mycostatin), quintozone (formerly known as PCNB), penicillin G, sodium omadine, and vancomycin in several combinations can prevent or inhibit contamination of *Cattleya aurantiaca* and *Stanhopea oculata* seedlings in vitro and *Phalaenopsis* (all three are orchids) flower stalk node cultures without adversely affecting the plants [20, 23, 24]. However, the effects of these compounds on the germination of orchid seeds, growth of excised shoot tips, and plantlet development of a larger number of species is not known. The work described here was undertaken to obtain information on the effects of bactericides and fungicides on seed germination and the subsequent growth of seedlings as well as shoot tip development in vitro. As in our previous work [20, 23, 24], the expectations are that combinations of these compounds can prevent or inhibit introduced contamination.

Correspondence should be addressed to J. Arditti.

Table 1. Bactericides and fungicides screened for their effects on orchid seed germination and excised shoot tips in culture

		Concentrations screened (ppm)				Stock solution in 70% ethanol ^a	Volume of stock solution per liter of medium (ml)	Source
Compound ^a	Property	Seeds		Shoot tips				
		Single compounds	In mixtures ^b	Single compounds	In mixtures ^c			
Amphotericin B (92.7% purity)	Fungicide	1, 5, 10	3	10	10	107.9 mg/10 ml	0.1, 0.3, 0.5, 1	SJ Luciana, Squibb Institute for Medical Research, PO Box 400, Princeton, NJ
Benomyl (Benlate, ^d 50% purity)	Fungicide	5, 25, 50 ^b	15	50	50	500 mg/100 ml	1, 3, 5, 10	HJ Thome, Agrichemicals, Dupont Co, Wilmington DE 48640
Dowicide (97% purity)	Fungicide and bactericide	1, 5		5		51.5 mg/20 ml	0.2, 1	Dow Chemical Co, PO Box 1847 2045 Dow Center, Midland, MI 48640
Ethirimol (100% purity)	Fungicide	25, 50, 100				1 g/10 ml	0.25, 0.5, 1	R Fox, Imperial Chemical Industries Ltd, Jealott's Hill Reserach Station, Bracknell, Berkshire RG12 GEY, UK
Gentamicin (purchased in sterile solution in ampule)	Bactericide	5, 25, 50	15	50	50		0.1, 0.3, 0.5, 1	Walter Prozman and TS Schaffer, Schering Corp, 60 Orange Street, Bloomfield, NJ 07003 ^a

Nystatin (4020 units/mg)	Fungicide	5, 25	5	25	25	250/ mg/10 ml	0.2, 1	See amphotericin B ^a
Quintozene (PCNB, 99% purity)	Fungicide	25, 50, 100	25	100	100	101 mg/10 ml	2.5, 5, 10	LR Faulkner, Olin Corp, PO Box 991, Little Rock, AR 72203
Penicillin G (1595 units/mg)	Bactericide	25, 50, 100	25	100	100	1000 mg/ml	0.25, 0.5, 1	See amphotericin B ^a
Sodium omadine (90% purity)	Fungicide and bactericide	1, 5	5	5	5	55.5 mg/10 ml	0.2, 1	See quintozene
Vancomycin (97.8% purity)	Bactericide	5, 25, 50	5	50	50	511 mg/10 ml	0.1, 0.5, 1	RJ Hosley, Lilly Research Laboratories, Indianapolis, IN 46206 ^a

^aAlso purchased from Sigma Chemical Corporation, PO Box 14508, St Louis, MO 63178

^bSee Tables 4 and 5

^cSee Table 6

^dAvailable from retail nurseries as systemic fungicides under a variety of trade names. These fungicides contain 50% active ingredient; therefore, the amounts used should be 10, 50, 100 mg/liter

Materials and Methods

Seeds

Mature seeds of *Cattleya elongata*, *Laelia tenebrosa*, and *Vanda tricolor* were used for the screening of single compounds (Table 1). *Brassavola nodosa*, *Cattleya aurantiaca*, *Cymbidium madidum*, *Dendrobium speciosum* var. *hillii*, *Epidendrum nocturnum*, *Grammatophyllum elegans*, *Laelia lobata*, *Phajus tankervilliae* × *P. maculatus*, *Odontoglossum citrosmum*, and *Zygopetalum mackay* were employed in germination experiments with mixtures of fungicides and bactericides (Table 1). The seeds were produced in the University of California, Irvine Orchid Collection (UCIOC), and stored over desiccant (CaCl_2) at 4°C. Availability of seeds determined their use in the screening of single compounds or mixtures.

The seeds were surface sterilized with saturated calcium hypochlorite (7 g/100 ml water) for 15 min in glass tubes and washed with sterile distilled water before placing them in culture [11].

Knudson C solution containing 0.2% graphite [20, 23, 24] was used as the basal culture medium. It was sterilized by autoclaving and the fungicides and bactericides were added in stock solutions (Table 1) to the medium before it solidified and were mixed by swirling. Prescription bottles, 100-ml capacity, containing 20 ml medium, were used as culture vessels.

Germination was evaluated five months after placing the seeds in culture on media containing single compounds and 10–12 weeks for mixtures by the growth index method [12, 21]. According to this method, seedlings are classified into six developmental stages (which are numbered one through six as in Fig. 1). Random samples of 100 seedlings are observed under a dissecting microscope and classified into the six stages (for example, 10% stage 1; 20% stage 2; 20% stage 3; 10% stage 4; 20% stage 5; and 20% stage 6). The percentage of seedlings in each stage is then multiplied by its number and the totals are added to produce the growth index $[(10 \times 1) + (20 \times 2) + (20 \times 3) + (10 \times 4) + (20 \times 5) + (20 \times 6) = 370]$. This growth index measures normal development rather than mere increases in size (i.e., appearance of meristems, leaves, and roots and not only increases in diameter, length, or weight of protocorms, for example), and can be subjected to statistical analysis [12, 21].

Shoot tips

New growths were taken from *Cymbidium* hybrids in the UCIOC, rinsed with tap water, and freed of dead, damaged, or excessively hard external tissues and parts. The shoots were then immersed in a mixture of household bleach–distilled water (1:1; vol/vol). Excision of explants was carried out under a dissecting microscope on an open laboratory bench washed with 95% ethanol. External leaves and leaf primordia were removed to expose

Figure 1. Stages in the germination of a *Cattleya* orchid [2].

the shoot tip which was excised by making four incisions [2]. Explants were cultured on modified Tsuchiya [25, 27; also listed in 2] medium (Table 2). The fungicides and bactericides used were the same as those employed with seeds (Table 1). Erlenmeyer flasks, 25- or 50-ml capacity containing 5 or 12.5 ml medium, respectively, were used as culture vessels for shoot tips. Flasks containing explants in liquid media were placed on a reciprocating shaker (60 oscillations/min). Shoot tip cultures were evaluated visually and are described in subjective terms.

All cultures were maintained under $22^{\circ} \pm 2^{\circ}\text{C}$, $0.8\text{mW}/\text{cm}^2$ and 18-h photoperiods provided by a mixture of Wide Spectrum Gro Lux tubes and incandescent lamps.

Results

In experiments with single compounds, there were no major differences between the growth indices of controls and of seeds germinated on fungicide and bactericide containing media (Table 3). Only the highest concentrations of benomyl, nystatin, quintozone, and penicillin G prevented germination of *Cattleya elongata*. All concentrations of amphotericin B and the highest level of gentamycin reduced germination of *Vanda tricolor* (Table 3). The solvent (70% ethanol) had no deleterious effects on germination (Table 3).

Table 2. Tsuchiya medium as used for the culture of *Cymbidium* shoot tips [2, 25, 27]

Component	Amount per liter
<i>Macrolements</i>	
Potassium nitrate, KNO_3	525 mg
Dicalcium phosphate, CaHPO_4	200 mg
Potassium phosphate, KH_2PO_4	250 mg
Ammonium sulfate, $(\text{NH}_4)_2\text{SO}_4$	500 mg
Magnesium sulfate, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	250 mg
Ferric tartrate, $\text{Fe}_2(\text{C}_4\text{H}_4\text{O}_6)_3$	30 mg
Tryptone	2 g
Sucrose	20 g
Water, distilled	to 1000 ml

Combinations of compounds had no negative effects on the germination of *Brassavola nodosa*, *Cattleya aurantiaca*, *Laelia lobata*, *Phajus tankervilleae* × *P. maculatus*, and *Zygopetalum mackay* (Table 4). In these experiments, too, 2 ml 70% ethanol per liter of medium had no inhibitory effects except on the germination of *Odontoglossum citrosimum* (Table 4). Appearance of roots in seedlings of *Brassavola nodosa* and *Cattleya aurantiaca* was enhanced by all combinations. Most of these combinations also enhanced the percentage of seedlings of these species and *Laelia lobata* which developed leaves (Table 5). These three species belong to the subtribe Laeliinae. Only combination 5 increased the growth indices of *Phajus* (subfamily Epidendroideae) and *Zygopetalum* (subfamily Vandoideae). All other combinations were without effect on the growth indices of these orchids (Table 4). There were no seedlings of *Odontoglossum citrosimum* past stage 1 (Fig. 1-1) on any of the combinations of bactericides and fungicides. On the basal medium, seedlings of *O. citrosimum* did not develop past stage 2 (Fig. 1-2). There were no seedlings of *Zygopetalum mackayi* past stage 3 (Fig. 1-3) on combinations 1, 4, and 5, and the two controls. On media 2 and 3, the seedlings did not develop past stage 2 (Fig. 1-2).

All combinations accelerated the appearance of roots and leaves in *Brassavola nodosa*. Only combinations 3–5 had a similar effect on *Cattleya aurantiaca*. In *Laelia lobata*, the appearance, but not the expansion, of leaves was accelerated by combinations 1–3 and 5 (Table 5).

None of the combinations screened could prevent contamination of the cultures of *Dendrobium speciosum* var. *hillii*. Only one flask (of three) on combination 5 (Table 4) remained uncontaminated and its growth index was 336. Of the seedlings, 1% had roots (average length 0.9 mm) and 18% had leaves (average length of the first leaf 2.21 ± 0.68 mm; of the second 1.34 ± 0.6 mm).

When used singly, the compounds we screened either delayed or entirely prevented the development of excised *Cymbidium* shoot tips (Table 6). All combinations had deleterious effects on these explants.

Table 3. Effects of separate bactericides and fungicides on the germination of orchid seeds

Compounds	Concentration (ppm)	Growth index ^a		
		<i>Cattelya elongata</i> ^b	<i>Laelia tenebrosa</i> ^b	<i>Vanda tricolor</i> ^c
Amphotericin B	1		344/11	330
	5		382/51	278
	10		321/51	324/55
Benomyl	5	370/75	410/26	480
	25	346/23	412/22	443/94
	50	NG ^d	NG ^d	NG ^d
Dowicide	1		381/42	375/66
	5	408/101	405/40	532/25
Ethirimol	25	450	403/45	430/158
	50		346/30	
	100	300/87	384/41	398/140
Gentamicin	5	421/18	350/21	326/44
	25	317/29	336/17	381/139
	50	313	346/76	302/2
Nystatin	5	298/259	412/9	402/16
	25	NG ^d	343/22	357/80
Quintozene	25	367/29	399/35	465/148
	50	361/75	364/57	428/111
	100	NG ^d	362/20	600
Penicillin G	25	377/41	382/9	456/63
	50	335/14	319/9	326/29
	100	NG ^d	366/38	355
Sodium omadine	1	350/50	339/15	447/150
	5	372/43	435/33	562/23
Vancomycin	5	392/12	378/24	484/30
	25	382/31	330/32	537/37
	50	341/21	373/24	494/57
Basal control		397/5	365/9	442/56
Solvent control		338/33	393/22	467/67

^a The values given are grow index/standard deviation^b Subfamily Epidendroideae, tribe Epidendreae, subtribe Laeliinae [7]^c Subfamily Vandoideae, Tribe Vandaeae, Subtribe Sarcanthinae [7]^d NG, no germination

Discussion

Orchid embryos lack cotyledons, leaf initials, and rudimentary roots. As a result, orchid seeds do not germinate like those of other plants. The dry seed swells (Fig. 1-1–2), forming a spherical body called a protocorm (Fig. 1-3), a term first used in 1890 by Melchior Treub to describe bodies formed when lycopods germinate [3]. Apical meristems on the tops of protocorms form leaves (Fig. 1-4) which elongate (Fig. 1-5) generally prior to, but at

Table 4. Effects of combination of fungicides and bactericides on the germination of orchid seeds

No.	Combinations ^b	Growth index ^a					
		<i>Brassavola nodosa</i> ^c	<i>Cattleya aurantiaca</i> ^c	<i>Laelia lobata</i> ^c	<i>Phajus tankervilleae</i> ^d × <i>P. maculatus</i> ^d	<i>Odontoglossum citrosum</i> ^e	<i>Zygopetqlum mackayi</i> ^f
1	Benomyl, nystatin penicillin G, gentamycin	415/30	274/43	195/69	114/5	100/0	200/70
2	Benomyl, nystatin, penicillin G, gentamycin, sodium omadine	390/0	269/20	225/23	111/14	100/0	137/4
3	Benomyl, nystatin, penicillin G, gentamycin, amphotericin B, vancomycin	436/27	211/69	137/35	126/10	100/0	151/2
4	Benomyl, quintozone penicillin G, amphotericin B, sodium omadine	436/39	271/64	125/19	114/4	100/0	213/13
5	Benomyl, nystatin, amphotericin B, penicillin G, sodium omadine	408/40	246/55	199/125	148/23	100/0	259/16
<i>Controls</i>							
	Basal medium only		259/9	191/37	118/15	125	169/39
	Basal medium plus 2.0 ml 70% ethanol/liter	230	173/25	169/45	153/23	100/0	193/34

^aThe values given are growth index/standard deviation^bSee Table 1 for concentrations^cSubfamily Epidendroideae, tribe Epidendroideae, subtribe Laeliinae [7]^dSubfamily Epidendroideae, tribe Epidendroideae, subtribe Blentiniinae [7]^eSubfamily Vandoideae, tribe Cymbidiaceae, subtribe Oncidiinae [7]^fSubfamily Vandoideae, tribe Maxillariaceae, subtribe Zygopetalinae [7]

times together with, the appearance of roots (Fig. 1-6). Consequently, substances which affect seedlings may do so at any one of several stages, including swelling of the seeds, formation of an apical meristem, as well as initiation and elongation of leaves and roots. Because of this, germinating orchid seeds are very suitable for studying the effects of fungicides and bactericides in culture media.

The orchid species used in these experiments were affected differently by the compounds screened. These differences did not always correlate with the systematic affinities [7] of the orchids (Tables 3-5). Growth of *Cattleya aurantiaca* and *Stanhopea occulata* seedlings was not affected deleteriously by 70% ethanol [23, 24], but germination of *Odontoglossum citrosimum* was inhibited. The very low growth index of the latter on basal medium, however, suggested that the seeds may have been of low viability and therefore very susceptible to inhibition (the amount of work and time factors associated with germinating orchid seeds are such that viability tests are not practical). Previous reports regarding the use of bactericides and fungicides on seed germination indicated that both the effects of the compounds and the tolerance of different plant species may vary. For example, benomyl was less toxic to cottonseeds than 5-ethoxy-3-(trichloromethyl)-1, 2, 4-thiadiazole and carboxin [6]. On the other hand, 'benomyl gave a slightly stunting action to cotton seedlings at 160 ppm' [1] and inhibited germination, length of radicles, and allantoinase activity in seeds of *Arachis hypogaea*, *Glycine max.*, *Phaseolus aureus*, and *P. mungo* [16]. Benomyl [methyl-1(butylcarbamoyl)-2-benzimidazole carbamate] and BAS 3460F (methyl-2-benzimidazole carbamate), but not thiabendazole, can break the heat-induced seed dormancy in celery seeds [22].

Temik had a deleterious effect on the germination of Egyptian cotton, but vitavax/captan and disyston or a combination of both frequently enhanced it [10]. Benomyl, thiram, and captan either enhanced or had no deleterious effects on the germination of wheat [8] and peas [13]. Combinations of penicillin G, nystatin (Mycostatin, Squibb), and amphotericin B (Fungizone, Squibb) prevented in vitro contamination of seedlings of *Hibiscus esculentus*, *Vigna sinensis*, and *Dolichos lablab* [9]. Our findings with *Cattleya aurantiaca* and *Stanhopea occulata* seedlings and cultures of *Phalaenopsis* flower stalk nodes also demonstrated that some bactericides and fungicides may enhance growth somewhat or at least have no deleterious effects, whereas others may be inhibitory [20, 23, 24]. Thus our findings are similar in principle to previous reports.

Benomyl is a systemic fungicide which is taken up and translocated by plant cells and organs [19]. This means that in the presence of benomyl not only the culture medium but also the plant material will be protected from fungal contamination. In addition, benomyl has been reported to have a cytokinin-like effect in wheat leaf [26], and soybean callus and radish cotyledon [18] bioassays, as well as a growth-enhancing effect on shoot and

Table 5. The effects of combinations of fungicides and bactericides on root and leaf formation and growth^a

		<i>Brassavola nodosa</i>				
Mixtures		Roots ^b		Leaves ^c		
		Seedlings with roots (%)	Length (mm)	Seedlings with leaves (%)	Length (mm) ^d	
Combinations ^b	No.				1st leaf	2nd leaf
Benomyl, nystatin penicillin G, gentamycin	1	1.7/0.6	0.56/0.13	20.0/13.9	1.32/0.49	0.75/0.26
Benomyl, nystatin penicillin G, gentamycin, sodium omadine	2	1.0/1.7	0.62/0.33	31.3/11.2	1.91/11.2	0.95/0.46
Benomyl, nystatin, penicillin G, gentamycin, amphotericin B, vancomycin	3	2.3/2.5	0.39/0.15	47.0/21.5	1.88/0.63	0.93/0.44
Benomyl, quintozone penicillin G, amphotericin B, sodium omadine	4	1.0/1.0	0.58/0.14	51.3/18.6	1.93/0.90	0.87/0.49
Benomyl, nystatin, amphotericin B, penicillin G, sodium omadine	5	5.7/8.1	0.45/0.07	57.7/1.5	2.72/1.48	1.38/0.91
<i>Controls</i>						
Basal medium only		—	—	—	—	—
Basal medium plus 2.0 ml 70% ethanol/liter		0 ^e	—	15	1.50/0.41	0.94/0.30

<i>Cattleya aurantiaca</i> ^e			<i>Laelia lobata</i> ^e			<i>Phagus tankervilleae</i> × <i>P. maculatus</i>
Leaves ^c			Leaves ^c			
Seedlings with leaves (%)	Length (mm)		Seedlings with leaves ^f (%)	Length (mm) ^d		Seedlings with leaves (%)
	1st leaf	2nd leaf		1st leaf	2nd leaf	
0 ^g	—	—	11.30/16.3	—	—	There were no seedlings past stage 3
3.30/4.0	1.36/0.29	0.92/0.40	10.70/3.1	—	—	There were no seedlings past stage 3
14.00/11.5	2.34/1.22	1.12/0.39	9.00/6.1	—	—	1.0/1.0 ^t
16.00/21.0	2.27/1.01	1.03/0.37	1.30/2.3	2.89/0.73	1.36/0.57	1.0/1.7 ^f
22.00/13.0	2.85/1.56	1.33/0.86	13.0/18.4	—	—	There were no seedlings past stage 2
2.70/3.8	1.69/0.54	1.23/0.14	1.0/1.7	1.5/0.62	1.17/0.63	There were no seedlings past stage 2
0.23/0.40 ^g	—	—	There were no seedlings past stage 3 ^h			0.3/0.6 ^g

^a Average of all replicas/standard deviation; see Table 4 for systematic affiliations of the orchids

^b By definition, roots are preset only on stage 6 seedlings (Fig. 1)

^c Stages 5 and/or 6 have leaves (Fig. 1)

^d The 1st (lower) leaf is usually longer

^e None of the seedlings formed roots

^f There were no seedlings past stage 4 (Fig. 1), which means that none had expanded leaves

Table 6. Effects of bactericides and fungicides on the growth and development of excised *Cymbidium* (subfamily Vandoideae, tribe Cymbideae, subtribe Cyrtopodiinae [7]) shoot tips

No.	Compound	Time for explants to turn green (days)	Remarks
	Control	12 ± 2	Normal growth, proliferation
1	<i>Single compounds</i> Amphotericin B	17 ± 2	Except for a delay, normal growth, proliferation, and plantlet formation
2	Benlate	42 ± 3	Weak and inhibited growth and development; no shoot formation
3	Dowicide	23 ± 2	Weak and inhibited growth and development; no shoot formation
4	Gentamycin	57 ± 5	Weak and inhibited growth and development; no shoot formation
5	Nystatin	23 ± 2	Except for a delay, normal growth, development and plantlet formation
6	Quintozene		No growth
7	Penicillin G	25 ± 2	Normal callus growth; no plantlet formation
8	Sodium omadine	23 ± 2	Except for a delay, normal growth and development
9	Vancomycin		No growth
	<i>Mixtures</i>		
	a Benomyl		
	Gentamycin		
	Nystatin		
	Penicillin G		No growth

b	Benomyl Gentamycin Nystatin Penicillin G Sodium omadine	30 ± 5	A small callus formed within 3 weeks, but died 14 days later
c	Amphotericin Benomyl Gentamycin Nystatin Penicillin G Vancomycin	25 ± 3	A small callus formed within 3 weeks, but died 14 days later
d	Amphotericin B Benomyl Quintozene Penicillin G Sodium omadine		No growth
e	Amphotericin B Benomyl Nystatin Penicillin G Sodium omadine		No growth
f	Amphotericin B Benomyl Sodium omadine		No growth

root development in *Asparagus officinalis* tissue cultures [28]. These facts render benomyl an especially attractive fungicide for use in tissue culture. However, caution is necessary in such use since benomyl had an inhibitory effect on shoot tip cultures of *Cymbidium* (Table 6) and *Colocasia esculenta* (Arditti and Gonzales, unpublished results). These deleterious effects may be due to metabolic and/or enzyme inhibition [16] as well as the induction of chromosome aberrations [29].

A number of antibiotics have been used in tissue cultures with varying results. Bacitracin (50–100 ppm), griseofulvin (2–5 ppm), oxytetracycline (2.5 and 5 ppm), and streptomycin (2.5 ppm or less) are not toxic to *Catharanthus roseus* tissue cultures [4]. Rifampicin (50 µg/ml) was shown to be effective against contaminating bacteria in cultured explants of *Helianthus tuberosus* without affecting differentiation of tracheary elements, rates of cell division, or DNA synthesis [17]. Low levels of kanamycin enhanced shoot differentiation in callus cultures of tobacco and carrots [15]. Therefore, our findings with shoot tips of *Cymbidium* (Table 5) and *Colocasia esculenta* (unpublished results) suggest that these plants are particularly sensitive to the fungicides and bactericides we screened, or that these compounds are more toxic to explants than bacitracin, griseofulvin, oxytetracycline, streptomycin, rifampicin, and kanamycin.

Our findings (Tables 4 and 5) suggest that (a) on occasion seeds may be so heavily laden with spores that nonphytotoxic concentrations of fungicides and/or bactericides cannot prevent contamination, or (b) some contaminants may be unaffected by the compounds used. Support for the latter comes from reports that benomyl (a) was not as effective as other compounds [14] in the control of *Alternaria* spp., *Botrytis* sp., *Cladosporium* spp., *Cochliobolus sativus*, *Penicillium* spp., *Rhizopus* sp. and *Streptomyces* spp., (b) did not eliminate pathogens [13] from pea seeds, (c) could not control [5] *Aphanomyces cochlioides* in sugar-beet seedling, and (d) controlled only temporarily the contamination of *Hibiscus esculentus*, *Vigna sinensis*, and *Dolichos lablab* seedlings on nutrient agar [9].

The need for, and importance of, agents which can combat contamination can be expected to increase as the use of tissue culture and in vitro seed germination becomes more widespread. Our findings suggest that useful compounds may be identified and appropriate combinations can be formulated for general use.

Acknowledgments. This research has been supported in part by a grant from the American Orchid Society Fund for Research and Education, and University of California Undergraduate Research Scholarships for which we thank Diana Sherwood. We also thank Alison F. Oliva for assistance and for reading and commenting on the manuscript, as well as the manufacturers (Table 1) for samples of the compounds screened.

References

1. Al-Beldawi AS, Pinckard JA (1970) Control of *Rhizoctonia solani* on cotton seedlings by means of Benomyl. *Plant Dis Rep* 54:76–80
2. Arditti J (1977) Clonal propagation of orchids by means of tissue culture – a manual. In: Arditti J (ed) *Orchid biology, reviews and perspectives*, 1. Ithaca: Cornell University Press, pp 203–293
3. Arditti J (1979) Origin of protocorm. *Am Orchid Soc Bull* 48:228
4. Carew DP, Patterson BD (1970) The effect of antibiotics on the growth of *Catharanthus roseus* tissue cultures. *Lloydia* 33:275–277
5. Byford WJ, Prince J (1976) Experiments with fungicides to control *Aphanomyces coelicoides* in sugar beet. *Ann Appl Biol* 83:69–77
6. Davis RD, Pinckard JA (1971) Comparative systemic fungicidal activity and phytotoxicity of certain seed and soil fungicides potentially useful for control of cotton seedling diseases. *Plant Dis Rep* 55:1111–1115
7. Dressler RL (1981) *The orchids, natural history and classification*. Cambridge: Harvard University Press
8. El-Zayyat MM, Sawhney JS, Ablack M, Nashnoosh I (1974) Effect of some seed dressings on germination of wheat grain. *Libyan J Agric* 3:85–88
9. Griffiths DA (1965) Temporary control of interstitial contamination during growth of seedlings on nutrient agar. *Can J Microbiol* 11:1025–1027
10. Hamawi H, Mahdi MT, El Ghandour MA, Eid A, Sheltawi E (1977) Effect of seed treatment with pesticides on germination of some Egyptian cotton cultivars. *Seed Sci Technol* 5:97–103
11. Harrison C (1970) A simple method for flasking orchid seeds. *Am Orchid Soc Bull* 39:715–716
12. Mariat F (1952) Recherches sur la physiologie des embryons des orchidées. *Rev Gen Bot* 59:324–377
13. Maude RB, Kyle AM (1970) Seed treatment with benomyl and other fungicides for the control of *Ascochyta pisi* on peas. *Ann Appl Biol* 66:37–41
14. Mills JT, Wallace HAH (1972) Differential action of fungicides upon fungi occurring on wheat, barley, buckwheat, and oil seeds. *Can J Plant Sci* 52:281–290
15. Owens LD (1979) Kanamycin promotes morphogenesis of plant tissues. *Plant Sci Lett* 16:225–230
16. Paul B, Thapar VK, Singh R (1973) Effect of systemic fungitoxics on ureide metabolism in germinating seeds. *Indian J Agric Sci* 43:662–666
17. Phillips R, Arnott SM, Kaplan SE (1981) Antibiotics in plant tissue culture: rifampicin effectively controls bacterial contaminants without affecting the growth of short-term explant cultures of *Helianthus tuberosus*. *Plant Sci Lett* 21:235–240
18. Skene KGM (1972) Cytokinin-like properties of the systemic fungicide benomyl. *J Hort Sci* 47:179–182
19. Solel Z, Schooley JM, Edgington LV (1973) Uptake and translocation of benomyl and carbendazim (methyl benzimidazol-2-yl carbamate) in the symplast. *Pestic Sci* 4:713–718
20. Spencer SJ, Thurston KC, Johnson JA, Perera RG, Arditti J (1979–1980) Media which do not require sterilization for the culture of orchid seedlings and Phalaenopsis flower stalk nodes. *Malay Orchid Rev* 14:44–53
21. Spoerl E (1948) Amino acids as sources of nitrogen for orchid embryos. *Am J Bot* 35:88–95
22. Thomas TH (1973) Growth regulatory effects of three benzimidazole fungicides on the germination of celery (*Apium graveolens*) seeds. *Ann Appl Biol* 74:233–238
23. Thurston KC, Spencer SJ, Arditti J (1979) Phytotoxicity of bactericides and fungicides in orchid culture media. *Am J Bot* 66:825–835
24. Thurston KC, Spencer SJ, Arditti J (1980) Incorporation of fungicides and bactericides in orchid seed and seedling culture media. *Proc 9th World Orchid Conf, Bangkok, 1978*, pp 179–190

25. Tsuchiya I (1954) Possibility of germination of orchid seeds from immature fruits. Na Pua Okika O Hawaii Nei—Orchids Hawaii 4:11–16
26. Williams PF (1975) Cytokinin activity of benomyl. Aust Plant Pathol Soc Newslett 4:12–13
27. Wimber DE (1963) Clonal multiplication of cymbidiums through tissue culture of the shoot meristem. Am Orchid Soc Bull 32:105–107
28. Yang H-J (1976) Effect of benomyl on *Asparagus officinalis* L. shoot and root development in culture media. HortScience 11:473–474
29. Zutshi U, Kaul BL (1975) Studies on the cytogenetic activity of some common fungicides in higher plants. Cytobios 12:61–67