UC Irvine

UC Irvine Previously Published Works

Title

Exercise Science and Child Health: A Tale of Many Journeys.

Permalink

<https://escholarship.org/uc/item/9f48v361>

Journal

Pediatric exercise science, 31(2)

ISSN

0899-8493

Author

Cooper, Dan M

Publication Date

2019-05-01

DOI

10.1123/pes.2018-0259

Peer reviewed

Exercise Science and Child Health: A Tale of Many Journeys

Dan M. Cooper

University of California, Irvine

Children are the most naturally physically active human beings; reduced physical activity is a cardinal sign of childhood disease, and exercise testing provides mechanistic insights into health and disease that are often hidden when the child is at rest. The physical inactivity epidemic is leading to increased disease risk in children and, eventually, in adults in unprecedented ways. Cardiopulmonary exercise testing (CPET) biomarkers are used to assess disease severity, progress, and response to therapy across an expanding range of childhood diseases and conditions. There is mounting data that fitness in children tracks across the life span and may prove to be an early, modifiable indicator of cardiovascular disease risk later in life. Despite these factors, CPET has failed to fulfill its promise in child health research and clinical practice. A major barrier to more accurate and effective clinical use of CPET in children is that data analytics and testing protocols have failed to keep pace with enabling technologies and computing capacity. As a consequence, biomarkers of fitness and physical activity have yet to be widely incorporated into translational research and clinical practice in child health. In this review, the author re-examines some of the long-held assumptions that mold CPET in children. In particular, the author suggests that current testing strategies that rely predominantly on maximal exercise may, inadvertently, obfuscate novel and clinically useful insights that can be gleaned from more comprehensive data analytics. New pathways to discovery may emanate from the simple recognition that the physiological journey that human beings undertake in response to the challenge of exercise may be far more important than the elusive destination of maximal or peak effort.

The purpose of this review is to focus on several intersecting journeys, new pathways necessary to advance the field of exercise medicine and exercise science in child health. We will cover the journey of concept and discovery in which novel technologies are reshaping how we measure and gauge fitness in critical periods of growth and development in children, and the physiological journey that integrates thermodynamic, biologic, and cognitive mechanisms permitting children to engage in physical activities that vary widely in intensity, duration, and skill. Collectively, these journeys are bringing us to a new and exciting era of discovery for all of child health research and clinical application.

The Nobel Laureate A.V. Hill was the pioneering British exercise scientist who in the 1920s accessed the novel technologies and conceptualized the existence of a maximum level of oxygen intake. In describing a series of exercise tests in several healthy volunteers, Hill et al (48) wrote, “The oxygen intake rises steadily as the [running] speed is increased, attaining a maximum, however, beyond which no bodily effort can drive it.” Hill and his group were inquisitive experimentalists and also noted that the so-called maximal levels of $\dot{V}O_2$ in a given individual intake were dependent on many conditions such as the type of exercise and the concentration of inspired oxygen.

Hill was acutely aware that his laboratory’s focus on exercise in the context of athletics limited the generalizability of the nascent discipline of exercise physiology. In his introduction to a series of lectures given at Cornell University in 1926–1927, he (47) wrote, “The complaint has been made to me—‘Why investigate athletics, why not study the process of industry or of disease?’ The answer is twofold. (i) The processes of athletics are simple and measureable and carried out to a constant degree, namely to the utmost of a man’s powers: those of industry are not; and (ii) athletes themselves, being in a state of health and dynamic equilibrium, can be experimented on without danger and can repeat their performances exactly again and again.”

Hill's prescience in grasping that the approach to exercise science depended on the context in which physical activity (PA) was considered (eg, athletics, presence of disease) was not shared by the majority of physiologists and clinicians over the past century. Historical realities, the cultural zeitgeist, and politics have always shaped scientific research (83), and exercise science has not been an exception. A century ago, the Russian revolution prompted scientists to better understand the physiological limits of physical labor as productivity at that time depended in no small measure on the physical work capacity of laborers (87). In the United States, entry into World War I and institution of mandatory conscription to military service revealed an alarming percentage of young Americans simply unfit for service. It is not surprising that the Harvard Fatigue Laboratory, established in 1927 and housed in the basement of the Harvard School of Business, led the nation in understanding the limits of physical capacity in human beings with a particular focus on industry and the military (94).

The focus on athletic and sports performance has remained the predominant theme in pediatric exercise science. As shown in Figure 1, an exponential increase in research in publications focused on exercise testing in the pediatric age range in the context of sports and athletics has occurred over the past 5 to 6 decades. A far smaller increase has occurred over the same period examining exercise testing and performance in the context of disease or illness. In Figure 2, I present a 10-year snapshot (2007–2017) of the relative focus of research publications in pediatric exercise on relevant diseases and conditions.

Figure 1 — A bibliometrics search examining the use of exercise testing to assess functional aspects of fitness in children in both health and disease.

Figure 2 — Distribution of publications in exercise testing and childhood conditions, and diseases over the past 10 years.

Physical Activity: An Essential Element of Growth and Development in Children and Adolescents

Exercise in children and adolescents is not merely play but is an essential component of growth and development (39,57,91). Children are among the most spontaneously physically active human beings (66). It is not surprising that participation in PA is a major determinant of health across the life span and health-related quality of life in both healthy children and in children with chronic diseases (46,64,68,76). Despite this essential biologic role for PA, children have not been spared the relentless reduction in levels of PA, that is, creating a crisis in health care in our nation and throughout the world (58). Recognition of the enormous morbidity and cost of physical inactivity-related diseases, such as atherosclerosis, type 2 diabetes, and osteoporosis, has spurred new policy initiatives targeting preventive medicine early in life (18,23,74). The concept of pediatric origins of adult health and disease is gaining scientific merit (10,13,44,86), highlighting the need to transform existing notions of how to evaluate health in a growing child. A physically inactive (even normal weight) child may have no symptoms of disease, but evidence of deterioration in vascular health may already be present (20,51,92).

Equally worrisome is that the deleterious health effects of physical inactivity and poor fitness are exacerbated in children with chronic disease and/or disabilities (1,49,50,62,67,99) or with environmental-lifestyle conditions like obesity (53). Children with diseases or conditions previously associated with excessive mortality and morbidity during the first 2 decades of life (eg, cystic fibrosis, pulmonary hypertension, sickle cell disease, congenital heart disease, malignancies, etc) are living longer because of remarkable advances in research and care (35,37,43,55,59) but are often unable to achieve levels of PA and fitness associated with health benefits in otherwise healthy children (2,60,89,95). Not surprisingly, the health span (the period of life free from serious chronic diseases and disability [90]) of children with chronic diseases is threatened not only by the underlying disease but also by the compounding effects of insufficient PA and sedentary behavior (15,19,33,40,81). Increasing PA and fitness is feasible but has proven quite challenging to implement in a systematic manner (38).

Once a pattern of physical inactivity and a sedentary lifestyle is established, a vicious cycle ensues (Figure 3), in which constraints on PA harm immediate health and contribute to lifelong health impairment ranging from cardiovascular and metabolic disease to osteoporosis (16,24,32,57,61,63). Exactly what constitutes ideal physical fitness in a child with a chronic condition remains unknown. Finding beneficial levels of PA in children with chronic disease or disability is challenging because the optimal range of exercise is much narrower than in a healthy child (Figure 4).

The Physiological Journey of the Exercise Response: Rethinking the Guideposts

As previously noted, a major driving concept in exercise science research in both children and adults over the past century has been maximal oxygen intake ($\dot{V}O_{2\text{max}}$). Substantial controversy still surrounds this term, and many researchers prefer to employ $\dot{V}O_{2\text{peak}}$, particularly when there is no clear plateau of $\dot{V}O_2$ as work rate (WR) increases. There is no doubt that this approach has enormously advanced our understanding of the mechanisms involved in human PA. Maximal exercise tests in children are highly dependent on the willingness of each participant to continue exercise at relatively high WRs when dyspnea, muscle fatigue, and other stress sensations are commonly experienced. These tests involve exercise intensities that are infrequently attained in natural PA (7,12,41) and are not particularly enjoyable for many children (Figure 5). As a result, a fair amount of cheerleading by the laboratory personnel is often necessary to help the child achieve a maximal effort in cardiopulmonary exercise testing (CPET). Despite its proven clinical and research utility, true $\dot{V}O_{2\text{max}}$ (finding an unambiguous plateau in $\dot{V}O_2$)

only occurs in a minority of exercise tests even in healthy children and adults and maybe less so in children with chronic disease or disability (17,29,34).

Figure 3 — Physical inactivity in childhood, be it imposed by environmental factors or chronic diseases or conditions, leads to a vicious cycle that impairs health across the life span.

Figure 4 — Health benefits of exercise are determined, in part, by the energy expenditure associated with physical activity. Both too much (excessive) and too little (sedentarism) exercise can impair health. As shown, the range of healthy exercise is narrower in the child with chronic disease or disability. CPET and physical fitness testing must be designed to identify biological mechanisms that can be translated into finding the mode, frequency, duration, and intensity of exercise that can benefit health during critical periods of growth. CPET indicates cardiopulmonary exercise testing.

The conundrum of $\dot{V}O_2\text{max}$ testing is highlighted in the current controversy surrounding the idea that in order to “verify” $\dot{V}O_2\text{max}$, one must perform traditional progressive exercise tests to the limit of the participant’s tolerance followed within minutes by a constant WR “supramaximal” test at a higher WR, again, to the limit of the participant’s tolerance. The very name of the additional test, *supramaximal*, highlights the conceptual difficulty. A maximum is, after all, the greatest or highest amount possible or attained. “Supramax” does not exist. The data clearly indicate that $\dot{V}O_2$ in the supramaximal constant workload test is often higher than the “ $\dot{V}O_2\text{max}$ ” obtained moments before in the progressive test. This is a fascinating observation but in no way indicates the discovery of the real $\dot{V}O_2\text{max}$. A much more reasonable explanation is that the elusive $\dot{V}O_2\text{max}$ is dynamic that the prior exercise through well-

established impact on time constants, blood flow, lung function, cognitive factors, and so on, (9,14,65,103) altered physiological processes and permitted a greater capacity to uptake atmospheric O₂.

One of the transformative discoveries over the past several decades has been the ability of exercise to stimulate systemic immune, stress, and both proinflammatory and anti-inflammatory mediators and cytokines. The seminal observations by Ostrowski et al (75) in adults have been corroborated in children and adolescents (72,73). In retrospect, of course, a stress, “danger” type signaling by exercise is not surprising; even brief bouts of sufficiently intense PA lead to a metabolic acidosis, an outpouring of catecholamines, and a profound perturbation of cellular homeostasis (28,71). The initial acute increase in circulating leukocytes and cytokines such as interleukin-6 is often balanced by anti-inflammatory mechanisms, and exercise training, such as other short-term stresses, may ultimately prove to be, in balance, anti-inflammatory. Nonetheless, exercise stimulus of stress and inflammation can be dangerous. A number of studies have demonstrated that high-intensity exercise is aversive or unpleasant in both children and adults (36,88), and one is reminded of the popular mantra about exercise, “no pain, no gain.”

Figure 5 — “Found Art” in the pediatric exercise laboratory. Maximal exercise testing is uncomfortable and unpleasant for many children. A 9-year-old girl left this note for us. It is time to rethink exercise testing in children.

When a child performs standard CPET but does not reach an exercise level that can be classified as maximal, the whole test may be deemed a failure despite the wealth of data successfully collected. For example, in a large study of children and adolescents (mean age: 12.3 y) who had undergone the Fontan correction for congenital heart disease during childhood, only 40% achieved an acceptable $\dot{V}O_2\text{max}$ using current criteria (77). Many clinically oriented researchers are hesitant about pushing certain patient populations during high-intensity exercise when acidosis can ensue and stress mediators are elevated. Salvadego et al (85) studied exercise in a group of obese, otherwise healthy, adolescents and stopped exercise when the participant achieved a heart rate (HR) of 180 bpm. The authors noted, “A true maximal test was not performed to avoid the cardiovascular risks associated with maximal exercise in obese subjects.”

The value of alternatives to maximal testing strategies have been recognized for many years, but predominantly focused on using data obtained from submaximal portions of exercise tests to estimate a predicted $\dot{V}O_2\text{max}$ rather than on the value of the submaximal data itself. Åstrand and Rhyming (6) recognized over 50 years ago that $\dot{V}O_2\text{max}$ in adults could be predicted by HR measured during submaximal exercise. Fitness tests that require HR data alone are more feasible for large cohorts than studies in which gas exchange is measured in each participant. For example, Pate et al (78) used HR data collected in the National Health and Nutrition Examination Survey (70) to characterize fitness in 3287 children and adolescents (aged 12–19 y). One must balance the loss of accuracy introduced when $\dot{V}O_2$ is estimated from HR (often through complex formulas whose parameters are based on many assumptions)

with the increased cost and complexity associated with actual breath-by-breath measurements of gas exchange.

New Technology and New Pathways: Impact of Breath-By-Breath Measurement of Gas Exchange

The breakthrough maximal exercise studies in the early 1900s were made possible by technological advances in measurement of oxygen concentration. However, large volumes of gas were required, limiting the number of data points in which variables such as $\dot{V}O_2$ or $\dot{V}CO_2$ could be obtained during an exercise protocol. Progress in gas concentration and gas flow analytics proceeded rapidly with the advent of mass spectrometry and rapid detection of flow rates and vectors from small quantities of gas, and by the 1970s, breath-by-breath measurement of gas exchange became possible (104). The early devices were limited by computing capacity; for example, the then most advanced HP-1000 computer had only 64kb of memory and required clever programming and data manipulation to facilitate the calculations necessary for breath-to-breath analysis. Breath-by-breath CPET became commercially available in the late 1980s and permitted quantification of gas exchange response time over minutes and in adult studies led to a revolution in research focused on “gas exchange kinetics,” an engineering and integrative systems approach to the adaptive biology of PA. *Between 1976–1981 and 2011–2015, a PubMed search revealed an increase from 5 to 71 publications in adults using breath-by-breath exercise analytics in clinical trials. Over the same interval, studies in children increased only from 2 to 11. The time is opportune to apply enabling and new technologies to PA biomarkers in pediatric health.*

In a study of 169 children and adolescents (27), we demonstrated a set of remarkably strong relationships among submaximal slopes (obtained from the “forgotten” data of progressive exercise CPET), body weight, lean body mass (obtained from dual x-ray absorptiometry [DXA]), and the traditional peak $\dot{V}O_2$ (Figures 6 and 7). These observations hold the promise of greatly expanding the clinical and translational research utility of CPET. Earlier studies from our group have demonstrated the power of slope analysis in children with specific diseases to identify unique pathophysiological biomarkers associated with the underlying condition. For example, we found abnormal slopes of $\Delta\dot{V}_E/\Delta\dot{V}CO_2$ (the change in ventilations relative to the change in CO_2 output) in CF, consistent with increased ventilation/perfusion mismatching and dead space (67). In children who had undergone the Fontan surgical correction for a variety of congenital heart diseases, we noted reduced slopes of $\Delta\dot{V}O_2/\Delta WR$ quantifying the degree of impaired oxygen delivery (99). More recently, abnormal $\dot{V}O_2/WR$, $\dot{V}_E/\dot{V}O_2$, $\dot{V}O_2/HR$, and $\dot{V}_E/\dot{V}CO_2$ slopes were observed in patients with sickle cell disease (60). These observations were made during submaximal phases of maximal work CPET in sickle cell disease patients in whom caution should be exercised when encouraging PA at high intensities, characterized by elevated catecholamines and stress and inflammatory factors.

Taking full advantage of enabling technologies and computing capacity, new testing strategies have emerged over the past several decades. As valuable as traditional progressive-exercise CPET has proven, it tends to overlook one of the critical components of successful PA in children, namely the ability to rapidly respond to and recover from brief bouts of exercise. The evolutionary pressures of successfully gathering or hunting food and/or escaping predation led to integrated physiological systems in which (in response to exercise) cardiac output increased rapidly, oxygen was delivered quickly to working muscles, and blood flow maintained to key nonexercising organs, such as the brain, so that executive function could be maintained and wise decisions (where to run; when to hide) made. Testing strategies have emerged that accentuate the ability to gauge impairment in the critical onset and recovery phases of the exercise response.

Figure 6 — Enhancing the value of CPET. Shown here is a schematic of the type of data derived from progressive exercise in children and adolescents. Traditional CPET relies on maximal or peak values. We propose to better capture dynamic relationships of breath-by-breath data during submaximal phases of progressive CPET and transform how we measure the physiologic response to acute exercise. CPET indicates cardiopulmonary exercise testing; WR, work rate; HR, heart rate; $\dot{V}O_2$, oxygen uptake; $\dot{V}O_{2max}$, maximal $\dot{V}O_2$ uptake.

Figure 7 — Scatterplots of CPET variables predicted to be relatively body size dependent versus TBM determined by standard scale (A, B, and C) and LBM determined by DXA (D, E, and F). Females are shown as open circles and males as closed circles. The correlation and the corresponding 95% CI are

shown for each plot. In all cases, the correlations with TBM were high but were significantly improved with LBM. CPET indicates cardiopulmonary exercise testing; CI, confidence interval; TBM, total body mass; LBM, lean body mass; $\dot{V}O_2$, oxygen uptake.

Interestingly, one of A.V. Hill's most important contributions was the concept of oxygen debt and deficit, an idea that arose from observations that $\dot{V}O_2$ increased exponentially, not immediately, in response to a constant WR perturbation. Hill et al surmised the existence of stores and energy mechanisms that provided the immediate requirements of muscular work and were not immediately dependent on oxygen uptake from the atmosphere. The deficit incurred was then "paid back" as evidenced by the delayed (exponential) return of $\dot{V}O_2$ to baseline after the termination of physical work.

Pioneering work from a number of investigators is emerging to examine gas exchange dynamics in children in health and disease (3,4,26). For example, Barker et al (9) combined breath-by-breath measurements of gas exchange with near-infrared spectroscopy to simultaneously measure muscle O_2 delivery, O_2 utilization, and muscle activity following priming exercise to better understand the factors limiting $\dot{V}O_2$ kinetics during high-intensity exercise tolerance in youth. Their findings were intriguing (Figure 8); despite an enhanced aerobic energy provision following the priming intervention, exercise tolerance was reduced between exercise bouts due, perhaps, to inadequate recovery of the muscle metabolic status before the subsequent exercise perturbation commenced. Earlier work in otherwise healthy children has demonstrated key differences in gas exchange kinetics between children and adults, most notably in the dynamics of ventilation and $\dot{V}CO_2$ (26,69).

Figure 8 — Mean HHb (top panel) and TOI (bottom panel) dynamics obtained from near-infrared spectroscopy during an initial exercise bout (dotted line) rapidly followed by a repeat bout (solid line). The vertical dotted line signifies the onset of exercise. Note that in the second bout, the TOI is elevated at baseline and throughout the exercise transition. Despite this, exercise tolerance was not improved by the priming bout. Reprinted with permission from Barker et al (9). TOI indicates tissue oxidation index; HHb, deoxyhemoglobin.

Dynamic exercise testing has also been done to elucidate disease mechanisms in childhood illnesses. For example, Hebestreit et al (45) used oxygen uptake kinetics to demonstrate an impairment of oxygen delivery in children with cystic fibrosis. A number of years ago, our group was able to demonstrate large differences in gas exchange responses to exercise in children who had undergone the Fontan surgical correction for congenital heart diseases. We used a relatively simple approach, the recovery from a 1-minute bout of constant WR exercise (99) (Figure 9).

Figure 9 — (A) HR response before, during, and after 1 minute of exercise in a 14-year-old Fontan group subject. The recovery kinetics were quantified using a single exponential, as shown. (B) HR and $\dot{V}O_2$ recovery times for control and Fontan group subjects. In control subjects, recovery times were longer after the higher work rate protocols (* $P < .05$). In Fontan group subjects, recovery times were prolonged compared with the same absolute (2 W/kg) and relative (3.5 W/kg) protocols in control subjects (** $P < .001$). HR indicates heart rate; $\dot{V}O_2$, oxygen consumption.

The Journey From “Noise” to “Signal”: Modern Approaches to CPET and Habitual PA Data Analysis Using Biologic Variability and Machine Learning

Cardiopulmonary exercise testing produces large data sets comprised of a variety of measures of breath-by-breath gas exchange and heart rate. Advanced computing has revolutionized analysis of large data sets, and many physiological data sets previously considered too noisy for predictive analysis can now yield valuable signals in biomedicine with statistics and machine learning expertise. In CPET, for example, Beltrame et al (11) recently found that it is possible to predict $\dot{V}O_{2\max}$ from steady-state and exercise transitions based on easy-to-obtain inputs and machine learning paradigms. Variability (and its related volatility and entropy) is increasingly used as biomarkers in assessing physiologic data in exercise and in

adult diseases (8,21,30,100,105). Early work in children has proved promising as well. Reybrouck et al (82) found that increased interbreath variability with oscillatory changes of $\dot{V}O_2$ during exercise in children with heart disease was predictive of peak $\dot{V}O_2$. In our own data (unpublished), as shown in Figure 10, we found increased CPET variability in younger compared with older children and as the intensity of exercise progressed.

One of the key challenges in assessing aerobic function in children and adolescents in health and disease is to link seemingly diverse testing strategies: formal CPET and measures of habitual PA (HPA) in the real lives of human beings. Surprisingly, in many studies, the correlation between CPET results and HPA metrics in both children and adults is poor. Either we are measuring phenomenon that is biologically minimally related, or alternatively, the way we are measuring these phenotypes is suboptimal. Earlier studies from this and other laboratories had suggested, for example, that the pattern of PA (eg, its tempo and intensity) might be useful in characterizing real-life PA in children (7,12), but these studies involved labor-intensive techniques, such as direct observation and manual recording of PA in field settings. Technological advances in wearable monitoring has led to an outpouring of new research on HPA in child health, and the actual measurement of HPA using techniques such as accelerometry, in contrast to recall or survey methodologies, may lead to a better understanding of the mechanisms that link laboratory CPET testing and HPA (106). Specific patterning of sedentary behaviors may also prove of benefit in linking HPA to cardiovascular disease risk in children and adolescents (31).

Figure 10 — A series of constant work rate exercise bouts (closed squares) reveals striking maturational differences in the relationship between $\dot{V}_E$ and $\dot{V}O_2$. This ratio (related to the oxygen uptake efficiency slope in standard CPET) is proving useful as a biomarker in a variety of cardiac and pulmonary diseases. Note the value of a test that uncovers data during both the onset and recovery from a brief exercise bout. CPET indicates cardiopulmonary exercise testing; $\dot{V}O_2$, oxygen consumption.

A number of researchers are rethinking how to analyze data obtained from wearable accelerometers in the quantification of HPA. Although the application of cut points continues to be common research practice, there is growing recognition that the relationship between proprietary accelerometer counts and energy expenditure is highly activity dependent and that a single regression equation cannot accurately determine energy expenditure across a wide range of activities. Validation studies involving independent samples indicate that regression-based cut-point approaches misclassify the

true intensity of HPA 35% to 45% of the time (93,97). An alternative approach to accelerometer data reduction that has significantly improved sensor-based measurement of HPA and sedentary behavior is pattern recognition through machine learning. A number of researchers have demonstrated that machine learning algorithms, such as artificial neural networks (22,42,52,80,98), provide highly accurate predictions of activity type and intensity from accelerometer data collected in children and adolescents. While the potential for data science advanced analytics in quantifying HPA in adults and children is great, much work needs to be done to achieve consensus and harmonization before this promise can be realized (56).

Physical Fitness as a Right of Children: The Journey Toward Robust Metrics

Physical inactivity as a leading cause of chronic illness is a global phenomenon, distributed across both low- and high-income nations (25,54,79). In a remarkable report card on the PA of children and youth in 38 countries from 6 continents (representing 60% of the world's population), Tremblay et al (96) recently noted, "The wide distribution of [PA for children] grades results in global average grades for all indicators being D or C [and] shows that the challenge of enhancing PA behaviors and opportunities for children and youth around the world remains unresolved."

Two transformative concepts highlight the need to provide clinicians with metrics of physical fitness in children that are accurate, cost-effective, and reproducible. First, there is growing recognition that "social determinants of health," defined by the World Health Organization (108) as "the conditions in which people are born, grow, live, work and age," must be identified if we are to mitigate health disparities. While it is evident that physical fitness and PA interact with social determinants of child and adolescent health in many ways (102), research is needed to identify, prioritize, and develop interventions that most effectively use exercise as medicine to address health disparities during critical periods of growth and development.

Second, the notion that children have unique rights was codified in the 1989 United Nations Convention on the Rights of the Child (101). The history of the United Nations Convention indicates the strong connection to child health (107). Among, the rights specified are as follows:

- The right of the child to the enjoyment of the highest attainable standard of health and to facilities for the treatment of illness and rehabilitation of health;
- The right of the child to develop the fullest potential of mental and physical abilities.

Although it is clear that the adoption of the Convention on the Rights of the Child has benefited the lives of children and adolescents (84), no mention is made of the profound role that physical fitness during childhood has on health. It would seem timely that the community of pediatric exercise scientists and clinicians promotes a clear role for physical fitness as a right of children throughout the world. A key challenge facing the community of pediatric exercise clinicians and scientists is the development of testing strategies that can be used to gauge physical fitness robustly, repeatedly, and cost effectively.

Much work needs to be done to achieve this goal. Over a 3-year period, a group of key stakeholders in pediatric exercise partners convened a working group to identify challenges facing optimal utilization of exercise testing in child health research (5).

The major recommendations of that group were as follows:

- (1) Build a formal framework for data harmonization and terminology interoperability in child health exercise science. This is essential to support the hoped-for expansion of clinical trials using CPET.

- (2) Create a network of child health–focused clinical and research exercise laboratories with the ultimate goal of establishing a data consortium. The network will
 - (a) Work to establish robust reference values in child health CPET;
 - (b) Begin to more precisely define the impact of disease and therapy on exercise responses during growth and development in children.
- (3) Engage and collaborate with existing child health– and adult exercise–focused groups and organizations to support life span research, global health initiatives, and create economies of scale in the data harmonization efforts.
- (4) Promote formal programs to train child health care professionals in essential areas of exercise physiology, PA assessment, and relevant concepts of fitness in children.
- (5) Utilize the network to establish common protocols for CPET to ensure, in particular, that data obtained from multicenter trials are truly comparable.
- (6) Empower the network to become a resource for review, data sharing, and innovation across the broad spectrum of PA research and clinical application in child health.

These challenges can guide our ultimate journey: To improve child health and health across the life span through PA research and its clinical application.

Acknowledgments

This work was supported in part by National Institutes of Health P01HD-n048721 grant, Pediatric Exercise and Genomic Research Center Systems Biology Research Fund, National Center for Advancing Translational Sciences (NCATS) UL1TR001414 grant and NCATS CTSA Collaboration Innovation Award, Project REACH (Revamping Exercise Assessments in Child Health), UL1TR002004.

References

1. Akber A, Portale AA, Johansen KL. Pedometer-assessed physical activity in children and young adults with CKD. *Clin J AmSoc Nephrol*. 2012;7:720–26. PubMed ID: 22422539 doi:10.2215/CJN.06330611
2. Alvarado AM, Ward KM, Muntz DS, et al. Heart rate recovery is impaired after maximal exercise testing in children with sickle cell anemia. *J Pediatr*. 2015;166:389–93.e1. PubMed ID: 25477159 doi:10.1016/j.jpeds.2014.10.064
3. Armon Y, Cooper DM, Zanconato S. Maturation of ventilator responses to 1-minute exercise. *Pediatr Res*. 1991;29:362–68. PubMed ID: 1906595 doi:10.1203/00006450-199104000-00007
4. Armstrong N, Barker AR. Oxygen uptake kinetics in children and adolescents: a review. *Pediatr Exerc Sci*. 2009;21:130–47. PubMed ID: 19556620 doi:10.1123/pes.21.2.130
5. Ashish N, Bamman MM, Cerny FJ, et al. The clinical translation gap in child health exercise research: a call for disruptive innovation. *Clin Transl Sci*. 2015;8:67–76. PubMed ID: 25109386 doi:10.1111/cts.12194
6. Åstrand PO, Rhyming I. Anomogram for calculation of aerobic capacity from pulse rate during sub-maximal work. *J Appl Physiol*. 1954; 7:218–21. PubMed ID: 13211501 doi:10.1152/jappl.1954.7.2.218
7. Bailey RC, Olson J, Pepper SL, Barstow TJ, Porszasz J, Cooper DM. The level and tempo of children's physical activities: an observational study. *Med Sci Sports Exerc*. 1995;27:1033–41. PubMed ID: 7564970 doi:10.1249/00005768-199507000-00012

8. Bansal T, Haji GS, Rossiter HB, Polkey MI, Hull JH. Exercise ventilatory irregularity can be quantified by approximate entropy to detect breathing pattern disorder. *Respir Physiol Neurobiol.* 2018; 255:1–6. PubMed ID: 29730423 doi:10.1016/j.resp.2018.05.002
9. Barker AR, Trebilcock E, Breese B, Jones AM, Armstrong N. The effect of priming exercise on O₂ uptake kinetics, muscle O₂ delivery and utilization, muscle activity, and exercise tolerance in boys. *Appl Physiol Nutr Metab.* 2014;39:308–17. PubMed ID: 24552371 doi:10.1139/apnm-2013-0174
10. Barker DJ, Osmond C, Forsén TJ, Thornburg KL, Kajantie E, Eriksson JG. Foetal and childhood growth and asthma in adult life. *Acta Paediatr.* 2013;102:732–38. PubMed ID: 23560734 doi:10.1111/apa.12257
11. Beltrame T, Amelard R, Villar R, Shafiee MJ, Wong A, Hughson RL. Estimating oxygen uptake and energy expenditure during treadmill walking by neural network analysis of easy-to-obtain inputs. *J Appl Physiol.* 2016;121:1226–33. PubMed ID: 27687561 doi:10.1152/jappphysiol.00600.2016
12. Berman N, Bailey RC, Barstow TJ, Cooper DM. Spectral and bout detection analysis of physical activity patterns in healthy, perpubertal boys and girls. *Am J Hum Biol.* 1998;10:289–97. PubMed ID: 28561394 doi:10.1002/(SICI)1520-6300(1998)10:3<289::AIDAJHB4>3.0.CO;2-E
13. Blaisdell CJ, Weinmann GG. NHLBI viewpoint: lung health and disease prevention research starting in childhood. *Pediatr Pulmonol.* 2015;50:604–6. PubMed ID: 25857257 doi:10.1002/ppul.23198
14. Blanchfield AW, Hardy J, De Morree HM, Staiano W, Marcora SM. Talking yourself out of exhaustion. *Med Sci Sports Exerc.* 2014;46:998–1007. PubMed ID: 24121242 doi:10.1249/MSS.0000000000000184
15. Bohr AH, Fuhlbrigge RC, Pedersen FK, de Ferranti SD, Müller K. Premature subclinical atherosclerosis in children and young adults with juvenile idiopathic arthritis. A review considering preventive measures. *Pediatr Rheumatol Online J.* 2016;14:3. PubMed ID: 26738563 doi:10.1186/s12969-015-0061-5
16. Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol.* 2012;2:1143–211. PubMed ID: 23798298
17. Breithaupt P, Adamo KB, Colley RC. The HALO submaximal treadmill protocol to measure cardiorespiratory fitness in obese children and youth: a proof of principle study. *Appl Physiol Nutr Metab.* 2012;37:308–14. PubMed ID: 22452609 doi:10.1139/h2012-003
18. Briggs AM, Cross MJ, Hoy DG, et al. Musculoskeletal health conditions represent a global threat to healthy aging: a report for the 2015 World Health Organization world report on ageing and health. *Gerontologist.* 2016;56(suppl 2):S243–55. PubMed ID: 26994264 doi:10.1093/geront/gnw002
19. Caughey MC, Loehr LR, Key NS, et al. Sick cell trait and incident ischemic stroke in the Atherosclerosis Risk in Communities study. *Stroke.* 2014;45:2863–7. PubMed ID: 25139879 doi:10.1161/STROKEAHA.114.006110
20. Cayres SU, de Lira FS, Machado-Rodrigues AM, Freitas Junior IF, Barbosa MF, Fernandes RA. The mediating role of physical inactivity on the relationship between inflammation and artery thickness in prepubertal adolescents. *J Pediatr.* 2015;166:924–9. PubMed ID: 25661410 doi:10.1016/j.jpeds.2014.12.057
21. Chang JS, Kim EY, Jung D, et al. Altered cardiorespiratory coupling in young male adults with excessive online gaming. *Biol Psychol.* 2015;110:159–66. PubMed ID: 26253868 doi:10.1016/j.biopsycho.2015.07.016

22. Chowdhury AK, Tjondronegoro D, Chandran V, Trost SG. Ensemble methods for classification of physical activities from wrist accelerometry. *Med Sci Sports Exerc.* 2017;49:1965–73. PubMed ID: 28419025 doi:10.1249/MSS.0000000000001291
23. Chu P, Pandya A, Salomon JA, Goldie SJ, Hunink MG. Comparative effectiveness of personalized lifestyle management strategies for cardiovascular disease risk reduction. *J Am Heart Assoc.* 2016;4: e002737. PubMed ID: 27025969
24. Chung A, Backholer K, Wong E, Palermo C, Keating C, Peeters A. Trends in child and adolescent obesity prevalence in economically advanced countries according to socioeconomic position: a systematic review. *Obes Rev.* 2015;17:276–95. PubMed ID: 26693831 doi:10.1111/obr.12360
25. Cominato L, Di Biagio GF, Lellis D, Franco RR, Mancini MC, de Melo ME. Obesity prevention: strategies and challenges in Latin America. *Curr Obes Rep.* 2018;7:97–104. PubMed ID: 29737493 doi:10.1007/s13679-018-0311-1
26. Cooper DM, Kaplan MR, Baumgarten L, Weiler-Ravell D, Whipp BJ, Wasserman K. Coupling of ventilation and CO₂ production during exercise in children. *Pediatr Res.* 1987;21:568–72. PubMed ID: 3110725 doi:10.1203/00006450-198706000-00012
27. Cooper DM, Leu SY, Galassetti P, Radom-Aizik S. Dynamic interactions of gas exchange, body mass, and progressive exercise in children. *Med Sci Sports Exerc.* 2014;46:877–86. PubMed ID: 24091992 doi:10.1249/MSS.0000000000000180
28. Cooper DM, Radom-Aizik S, Schwindt C, Zaldivar F Jr. Dangerous exercise: lessons learned from dysregulated inflammatory responses to physical activity. *J Appl Physiol.* 2007;103:700–9. PubMed ID: 17495117 doi:10.1152/jappphysiol.00225.2007
29. Cooper DM, Weiler-Ravell D, Whipp BJ, Wasserman K. Aerobic parameters of exercise as a function of body size during growth in children. *J Appl Physiol.* 1984;56:628–34. PubMed ID: 6706770 doi:10.1152/jappl.1984.56.3.628
30. Costa M, Goldberger A. Generalized multiscale entropy analysis: application to quantifying the complex volatility of human heartbeat time series. *Entropy.* 2015;17:1197–203. PubMed ID: 27099455 doi:10.3390/e17031197
31. Cristi-Montero C, Chillo'n P, Labayen I, et al. Cardiometabolic risk through an integrative classification combining physical activity and sedentary behavior in European adolescents: HELENA study. *J Sport Heal Sci.* 2019;8:55–62. PubMed ID: 30719384 doi:10.1016/j.jshs.2018.03.004
32. Cunningham SA, Kramer MR, Narayan KM. Incidence of childhood obesity in the United States. *N Engl J Med.* 2014;370:403–11. PubMed ID: 24476431 doi:10.1056/NEJMoa1309753
33. Dahhan A. Coronary artery ectasia in atherosclerotic coronary artery disease, inflammatory disorders, and sickle cell disease. *Cardiovasc Ther.* 2015;33:79–88. PubMed ID: 25677643 doi:10.1111/1755-5922.12106
34. Day JR, Rossiter HB, Coats EM, Skasick A, Whipp BJ. The maximally attainable VO₂ during exercise in humans: the peak vs. maximum issue. *J Appl Physiol.* 2003;95:1901–7. PubMed ID: 12857763 doi:10.1152/jappphysiol.00024.2003
35. Dodge JA. A millennial view of cystic fibrosis. *Dev period Med.* 2015;19:9–13. PubMed ID: 26003065
36. Ekkekakis P, Parfitt G, Petruzzello SJ. The pleasure and displeasure people feel when they exercise at different intensities: decennial update and progress towards a tripartite rationale for exercise intensity prescription. *Sports Med.* 2011;41:641–71. Available from: <http://link.springer.com/10.2165/11590680-000000000-00000>.

37. Ephrem G, Hebson C, John A, et al. Frontiers in Fontan failure: innovation and improving outcomes: a conference summary. *Congenit Heart Dis*. 2019;14:128–137. PubMed ID: 30343507
38. Foster GD, Linder B, Baranowski T, et al. A school-based intervention for diabetes risk reduction. *N Engl J Med*. 2010;363:443–53. PubMed ID: 20581420 doi:10.1056/NEJMoa1001933
39. Gates PE, Strain WD, Shore AC. Human endothelial function and microvascular ageing. *Exp Physiol*. 2009;94:311–6. PubMed ID: 19042980 doi:10.1113/expphysiol.2008.043349
40. Gibson TM, Ehrhardt MJ, Ness KK. Obesity and metabolic syndrome among adult survivors of childhood leukemia. *Curr Treat Options Oncol*. 2016;17:17. PubMed ID: 26951206 doi:10.1007/s11864-016-0393-5
41. Gilliam TB, Freedson PS, Geenen DL, Shahraray B. Physical activity patterns determined by heart rate monitoring in 6-7 year-old children. *Med Sci Sports Exerc*. 1981;13:65–7. PubMed ID: 7219138
42. Hagenbuchner M, Cliff DP, Trost SG, Van Tuc N, Peoples GE. Prediction of activity type in preschool children using machine learning techniques. *J Sci Med Sport*. 2015;18:426–31. PubMed ID: 25088983 doi:10.1016/j.jsams.2014.06.003
43. Hansmann G. Pulmonary hypertension in infants, children, and young adults. *J Am Coll Cardiol*. 2017;69:2551–69. PubMed ID: 28521893 doi:10.1016/j.jacc.2017.03.575
44. Hardy R, Lawlor DA, Kuh D. A life course approach to cardiovascular aging. *Future Cardiol*. 2015;11:101–13. PubMed ID: 25606706 doi:10.2217/fca.14.67
45. Hebestreit H, Hebestreit A, Trusen A, Hughson RL. Oxygen uptake kinetics are slowed in cystic fibrosis. *Med Sci Sports Exerc*. 2005;37:10–7. PubMed ID: 15632661
46. Herman KM, Hopman WM, Sabiston CM. Physical activity, screen time and self-rated health and mental health in Canadian adolescents. *Prev Med*. 2015;73:112–6. PubMed ID: 25660484 doi:10.1016/j.ypmed.2015.01.030
47. Hill AV. *Muscular Movement in Man*. New York, NY: McGraw-Hill; 1927.
48. Hill AV, Long CNH, Lupton H. Muscular exercise, lactic acid and the supply and utilisation of oxygen. *Proc R Soc B Biol Sci*. 1924;97:155–76. <http://rspb.royalsocietypublishing.org/cgi/doi/10.1098/rspb.1924.0048>.
49. Hoffman MC, Mulrooney DA, Steinberger J, Lee J, Baker KS, Ness KK. Deficits in physical function among young childhood cancer survivors. *J Clin Oncol*. 2013;31:2799–805. PubMed ID: 23796992 doi:10.1200/JCO.2012.47.8081
50. Houghton KM, Tucker LB, Potts JE, McKenzie DC. Fitness, fatigue, disease activity, and quality of life in pediatric lupus. *Arthritis Rheum*. 2008;59:537–45. PubMed ID: 18383417 doi:10.1002/art.23534
51. Idler N, Teuner CM, Hunger M, et al; GINIplus and LISApplus Study Groups. The association between physical activity and healthcare costs in children—results from the GINIplus and LISApplus cohort studies. *BMC Public Health*. 2015;15:437. PubMed ID: 25925399 doi:10.1186/s12889-015-1721-6
52. Jang Y, Kim S, Kim K, Lee D. Deep learning-based classification with improved time resolution for physical activities of children. *PeerJ*. 2018;6:e5764. PubMed ID: 30364555 doi:10.7717/peerj.5764
53. Jiménez-Pavón D, Kelly J, Reilly JJ. Associations between objectively measured habitual physical activity and adiposity in children and adolescents: systematic review. *Int J Pediatr Obes*. 2010;5:3–18. PubMed ID: 19562608 doi:10.3109/17477160903067601

54. Juma PA, Mohamed SF, Matanje Mwangomba BL, et al. Noncommunicable disease prevention policy process in five African countries authors. *BMC Public Health*. 2018;18:961. PubMed ID: 30168393 doi:10.1186/s12889-018-5825-7
55. Keegan TH, Ries LA, Barr RD, et al. Comparison of cancer survival trends in the United States of adolescents and young adults with those in children and older adults. *Cancer*. 2016;122:1009–16. PubMed ID: 26848927 doi:10.1002/cncr.29869
56. Kerr J, Marinac CR, Ellis K, et al. Comparison of accelerometry methods for estimating physical activity. *Med Sci Sports Exerc*. 2017;49:617–24. PubMed ID: 27755355 doi:10.1249/MSS.0000000000001124
57. Klentrou P. Influence of exercise and training during critical stages of bone growth and development. *Pediatr Exerc Sci*. 2016;28(2):178–86. PubMed ID: 26884506 doi:10.1123/pes.2015-0265
58. Kohl HW, Craig CL, Lambert EV, et al. The pandemic of physical inactivity: global action for public health. *Lancet*. 2012;380:294–305. <http://www.sciencedirect.com/science/article/pii/S0140673612608988>.
59. Lanzkron S, Carroll CP, Haywood C. Mortality rates and age at death from sickle cell disease: U.S., 1979–2005. *Public Health Rep*. 2013;128:110–116. PubMed ID: 23450875 doi:10.1177/003335491312800206
60. Liem RI, Reddy M, Pelligra SA, et al. Reduced fitness and abnormal cardiopulmonary responses to maximal exercise testing in children and young adults with sickle cell anemia. *Physiol Rep*. 2015;3:pii: e12338. PubMed ID: 25847915 doi:10.14814/phy2.12338
61. Lobstein T, Jackson-Leach R, Moodie ML, et al. Child and adolescent obesity: part of a bigger picture. *Lancet*. 2015;385:2510–20. PubMed ID: 25703114 doi:10.1016/S0140-6736(14)61746-3
62. Lunt D, Briffa T, Briffa NK, Ramsay J. Physical activity levels of adolescents with congenital heart disease. *Aust J Physiother*. 2003;49:43–50. PubMed ID: 12600253 doi:10.1016/S0004-9514(14)60187-2
63. Magnussen CG, Smith KJ, Juonala M. When to prevent cardiovascular disease? As early as possible: lessons from prospective cohorts beginning in childhood. *Curr Opin Cardiol*. 2013;28:561–68. PubMed ID: 23928921 doi:10.1097/HCO.0b013e32836428f4
64. Maher CA, Toohey M, Ferguson M. Physical activity predicts quality of life and happiness in children and adolescents with cerebral palsy. *Disabil Rehabil*. 2016;38:865–69. PubMed ID: 26218617 doi:10.3109/09638288.2015.1066450
65. Mattioni Maturana F, Peyrard A, Temesi J, Millet GY, Murias JM. Faster V̇O₂ kinetics after priming exercises of different duration but same fatigue. *J Sports Sci*. 2018;36:1095–102. PubMed ID: 28721747 doi:10.1080/02640414.2017.1356543
66. Mielgo-Ayuso J, Aparicio-Ugarriza R, Castillo A, et al. Physical activity patterns of the spanish population are mostly determined by sex and age: findings in the ANIBES study. *PLoS ONE*. 2016;11: e0149969. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0149969>.
67. Moser C, Tirakitsoontorn P, Nussbaum E, Newcomb R, Cooper DM. Muscle size and cardiorespiratory response to exercise in cystic fibrosis. *Am J Respir Crit Care Med*. 2000;162(5):1823–7. PubMed ID: 11069820 doi:10.1164/ajrcrm.162.5.2003057
68. Mutlu EK, Mutlu C, Taskiran H, Ozgen IT. Association of physical activity level with depression, anxiety, and quality of life in children with type 1 diabetes mellitus. *J Pediatr Endocrinol Metab*. 2015; 28:1273–8. PubMed ID: 26197465
69. Nagano Y, Baba R, Kuraishi K, et al. Ventilatory control during exercise in normal children. *Pediatr Res*. 1998;43:704–7. PubMed ID: 9585019 doi:10.1203/00006450-199805000-00021

70. National Health and Nutrition Examination Survey (NHANES). National Youth Fitness Survey (NYFS) Treadmill Examination Manual. 2012.
<https://www.cdc.gov/nchs/data/nyfs/treadmill.pdf>
71. Nemet D, Hong S, Mills PJ, Ziegler MG, Hill M, Cooper DM. Systemic vs. local cytokine and leukocyte responses to unilateral wrist flexion exercise. *J Appl Physiol*. 2002;93:546–54. PubMed ID: 12133863
72. Nemet D, Oh Y, Kim HS, Hill M, Cooper DM. Effect of intense exercise on inflammatory cytokines and growth mediators in adolescent boys. *Pediatrics*. 2002;110:681–89. PubMed ID: 12359780 doi:10.1542/peds.110.4.681
73. Nemet D, Pontello AM, Rose-Gottron C, Cooper DM. Cytokines and growth factors during and after a wrestling season in adolescent boys. *Med Sci Sports Exerc*. 2004;36:794–800. PubMed ID: 15126712
74. Nishi A, Milner DA, Giovannucci EL, et al. Integration of molecular pathology, epidemiology and social science for global precision medicine. *Expert Rev Mol Diagn*. 2016;16:11–23. PubMed ID: 26636627 doi:10.1586/14737159.2016.1115346
75. Ostrowski K, Rohde T, Asp S, Schjerling P, Pedersen BK. Pro- and anti-inflammatory cytokine balance in strenuous exercise in humans. *J Physiol*. 1999;515(pt 1):287–91. PubMed ID: 9925898 doi:10.1111/j.1469-7793.1999.287ad.x
76. Pacheco DR, Silva MJ, Alexandrino AM, Torres RM. Exercise-related quality of life in subjects with asthma: a systematic review. *J Asthma*. 2012;49:487–95. PubMed ID: 22554022 doi:10.3109/02770903.2012.680636
77. Paridon SM, Mitchell PD, Colan SD, et al. A cross-sectional study of exercise performance during the first 2 decades of life after the Fontan operation. *J Am Coll Cardiol*. 2008;52:99–107. PubMed ID: 18598887 doi:10.1016/j.jacc.2008.02.081
78. Pate RR, Wang CY, Dowda M, Farrell SW, O'Neill JR. Cardiorespiratory fitness levels among US youth 12 to 19 years of age: findings from the 1999-2002 National Health and Nutrition Examination Survey. *Arch Pediatr Adolesc Med*. 2006;160:1005–12. PubMed ID: 17018458 doi:10.1001/archpedi.160.10.1005
79. Patton GC, Azzopardi P, Kennedy E, Coffey C, Mokdad A. Global measures of health risks and disease burden in adolescents. In: Bundy DA, Silva ND, Horton S, Jamison DT, Patton GC, eds. *Child and Adolescent Health and Development*. 3rd ed. Washington, DC: The International Bank for Reconstruction and Development/the World Bank; 2017: 57–72. PubMed ID: 30212128
80. Pavey TG, Gilson ND, Gomersall SR, Clark B, Trost SG. Field evaluation of a random forest activity classifier for wrist-worn accelerometer data. *J Sci Med Sport*. 2017;20:75–80. PubMed ID: 27372275 doi:10.1016/j.jsams.2016.06.003
81. Reverri EJ, Morrissey BM, Cross CE, Steinberg FM. Inflammation, oxidative stress, and cardiovascular disease risk factors in adults with cystic fibrosis. *Free Radic Biol Med*. 2014;76:261–77. PubMed ID: 25172163 doi:10.1016/j.freeradbiomed.2014.08.005
82. Reybrouck T, Vangesselen S, Mertens L, Gewillig M. Increased interbreath variability of gas exchange during exercise in children with cardiomyopathy. *Heart*. 2007;93:377–8. PubMed ID: 17322519 doi:10.1136/hrt.2006.094508
83. Rörsch A. The progress of science—Past, present and future. *Humanities*. 2014;3:442–516. <http://www.mdpi.com/2076-0787/3/4/442>.
84. Ruck MD, Keating DP, Saewyc EM, Earls F, Ben-Arieh A. The United Nations convention on the rights of the child: its relevance for adolescents. *J Res Adolesc*. 2016;26:16–29. <http://doi.wiley.com/10.1111/jora.12172>.

85. Salvadego D, Lazzer S, Busti C, et al. Gas exchange kinetics in obese adolescents. Inferences on exercise tolerance and prescription. *Am J Physiol Regul Integr Comp Physiol*. 2010;299:R1298–305. PubMed ID: 20686169 doi:10.1152/ajpregu.00038.2010
86. Sata F. Developmental origins of health and disease (DOHaD) and epidemiology. *Nihon Eiseigaku Zasshi*. 2016;71:41–46. PubMed ID: 26832616 doi:10.1265/jjh.71.41
87. Scheffler RW. The power of exercise and the exercise of power: the Harvard Fatigue Laboratory, distance running, and the disappearance of work, 1919–1947. *J Hist Biol*. 2015;48:391–423. PubMed ID: 25287571 doi:10.1007/s10739-014-9392-1
88. Schneider M, Dunn A, Cooper D. Affect, exercise, and physical activity among healthy adolescents. *J Sport Exerc Psychol*. 2009;31:706–23. PubMed ID: 20384008 doi:10.1123/jsep.31.6.706
89. Schneiderman JE, Wilkes DL, Atenafu EG, et al. Longitudinal relationship between physical activity and lung health in patients with cystic fibrosis. *Eur Respir J*. 2014;43:817–23. PubMed ID: 24176992 doi:10.1183/09031936.00055513
90. Seals DR, Melov S. Translational geroscience: emphasizing function to achieve optimal longevity. *Aging*. 2014;6:718–30. PubMed ID: 25324468 doi:10.18632/aging.100694
91. Shay CM, Ning H, Daniels SR, Rooks CR, Gidding SS, Lloyd-Jones DM. Status of cardiovascular health in US adolescents: prevalence estimates from the National Health and Nutrition Examination Surveys (NHANES) 2005–2010. *Circulation*. 2013;127:1369–76. PubMed ID: 23547177 doi:10.1161/CIRCULATIONAHA.113.001559
92. Stary HC. Lipid and macrophage accumulations in arteries of children and the development of atherosclerosis. *Am J Clin Nutr*. 2000;72:1297S–306S. PubMed ID: 11063472 doi:10.1093/ajcn/72.5.1297s
93. Staudenmayer J, Pober D, Crouter S, Bassett D, Freedson P. An artificial neural network to estimate physical activity energy expenditure and identify physical activity type from an accelerometer. *J Appl Physiol*. 2009;107:1300–7. PubMed ID: 19644028 doi:10.1152/japplphysiol.00465.2009
94. Tipton CM. Contemporary exercise physiology: fifty years after the closure of Harvard Fatigue Laboratory. *Exerc Sport Sci Rev*. 1998;26:315–39. PubMed ID: 9696994 doi:10.1249/00003677-199800260-00014
95. Tirakitsoontorn P, Nussbaum E, Moser C, Hill M, Cooper DM. Fitness, acute exercise, and anabolic and catabolic mediators in cystic fibrosis. *Am J Respir Crit Care Med*. 2001;164:1432–7. PubMed ID: 11704591 doi:10.1164/ajrccm.164.8.2102045
96. Tremblay MS, Barnes JD, González SA, et al. Global Matrix 2.0: report card grades on the physical activity of children and youth comparing 38 countries. *J Phys Act Health*. 2016;13:S343–66. PubMed ID: 27848745 doi:10.1123/jpah.2016-0594
97. Trost SG, Wong WK, Pfeiffer KA, Zheng Y. Artificial neural networks to predict activity type and energy expenditure in youth. *Med Sci Sports Exerc*. 2012;44:1801–9. PubMed ID: 22525766 doi:10.1249/MSS.0b013e318258ac11
98. Trost SG, Zheng Y, Wong WK. Machine learning for activity recognition: hip versus wrist data. *Physiol Meas*. 2014;35:2183–9. PubMed ID: 25340887 doi:10.1088/0967-3334/35/11/2183
99. Troutman WB, Barstow TJ, Galindo AJ, Cooper DM. Abnormal dynamic cardiorespiratory responses to exercise in pediatric patients after Fontan procedure. *J Am Coll Cardiol*. 1998; 31(3):668–73. PubMed ID: 9502651 doi:10.1016/S0735-1097(97)00545-7
100. Usemann J, Xu B, Delgado-Eckert E, et al. Dynamics of respiratory symptoms during infancy and associations with wheezing at school age. *ERJ Open Res*. 2018;4:00037–2018. PubMed ID: 30474038 doi:10.1183/23120541.00037-2018

101. Verhellen E. Convention on the Rights of the Child : Background, Motivation, Strategies, Main Themes. Leuven, Belgium: Garant; 2000. <https://eric.ed.gov/?id=ED454987>.
102. Viner RM, Ozer EM, Denny S, et al. Adolescence and the social determinants of health. *Lancet*. 2012;379:1641–52. <https://www.sciencedirect.com/science/article/pii/S0140673612601494>.
103. Wallace PJ, Mckinlay BJ, Coletta NA, et al. Effects of motivational self-talk on endurance and cognitive performance in the heat. *Med Sci Sports Exerc*. 2017;49:191–9. PubMed ID: 27580154 doi:10.1249/MSS.0000000000001087
104. Wasserman K, Whipp BJ. Breath-by-breath analysis of pulmonary gas exchange and the hyperpnea of exercise under non-steady-state and steady-state conditions. *Chest*. 1972;61:46S–7S. PubMed ID: 5009867 doi:10.1016/S0012-3692(15)32698-2
105. Weippert M, Behrens M, Gonschorek R, Bruhn S, Behrens K. Muscular contraction mode differently affects autonomic control during heart rate matched exercise. *Front Physiol*. 2015;6:156. PubMed ID: 26042047 doi:10.3389/fphys.2015.00156
106. Wittekind SG, Edwards NM, Khoury PR, et al. Association of habitual physical activity with cardiovascular risk factors and target organ damage in adolescents and young adults. *J Phys Act Health*. 2018;15:176–82. PubMed ID: 29172989 doi:10.1123/jpah.2017-0276
107. Woltanowski P, Wincewicz A, Sulkowski S. Protection of children's human rights and health: a legacy of Julian Kramsztyk, Janusz Korczak, and Ludwik Rajchman. *Glob Pediatr Health*. 2018; 5:2333794X1775415. PubMed ID: 29383326 doi:10.1177/ 2333794X17754157
108. World Health Organization. About social determinants of health. 2017. http://www.who.int/social_determinants/sdh_definition/en/.