

# UC Davis

## UC Davis Previously Published Works

### Title

Complete genome sequence of Haloterrigena turkmenica type strain (4kT)

### Permalink

<https://escholarship.org/uc/item/9ft9n1s9>

### Journal

Environmental Microbiome, 2(1)

### ISSN

2524-6372

### Authors

Saunders, Elisabeth  
Tindall, Brian J  
Fähnrich, Regine  
et al.

### Publication Date

2010

### DOI

10.4056/sigs.681272

Peer reviewed

## Complete genome sequence of *Haloterrigena turkmenica* type strain (4k<sup>T</sup>)

Elisabeth Saunders<sup>1,2</sup>, Brian J. Tindall<sup>3</sup>, Regine Fährnich<sup>3</sup>, Alla Lapidus<sup>1</sup>, Alex Copeland<sup>1</sup>, Tijana Glavina Del Rio<sup>1</sup>, Susan Lucas<sup>1</sup>, Feng Chen<sup>1</sup>, Hope Tice<sup>1</sup>, Jan-Fang Cheng<sup>1</sup>, Cliff Han<sup>1,2</sup>, John C. Detter<sup>1,2</sup>, David Bruce<sup>1,2</sup>, Lynne Goodwin<sup>1,2</sup>, Patrick Chain<sup>1,2</sup>, Sam Pitluck<sup>1</sup>, Amrita Pati<sup>1</sup>, Natalia Ivanova<sup>1</sup>, Konstantinos Mavromatis<sup>1</sup>, Amy Chen<sup>4</sup>, Krishna Palaniappan<sup>4</sup>, Miriam Land<sup>1,5</sup>, Loren Hauser<sup>1,5</sup>, Yun-Juan Chang<sup>1,5</sup>, Cynthia D. Jeffries<sup>1,5</sup>, Thomas Brettin<sup>1,5</sup>, Manfred Rohde<sup>6</sup>, Markus Göker<sup>3</sup>, James Bristow<sup>1</sup>, Jonathan A. Eisen<sup>1,7</sup>, Victor Markowitz<sup>4</sup>, Philip Hugenholtz<sup>1</sup>, Hans-Peter Klenk<sup>3</sup>, and Nikos C. Kyrpides<sup>1\*</sup>

<sup>1</sup> DOE Joint Genome Institute, Walnut Creek, California, USA

<sup>2</sup> Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico, USA

<sup>3</sup> DSMZ – German Collection of Microorganisms and Cell Cultures GmbH, Braunschweig, Germany

<sup>4</sup> Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, California, USA

<sup>5</sup> Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

<sup>6</sup> HZI – Helmholtz Centre for Infection Research, Braunschweig, Germany

<sup>7</sup> University of California Davis Genome Center, Davis, California, USA

\*Corresponding author: Nikos C. Kyrpides

**Keywords:** extreme halophile, thermophile, free-living, aerobic, non-pathogenic, carotenoids-containing, *Halobacteriaceae*, GEBA

---

*Haloterrigena turkmenica* (Zvyagintseva and Tarasov 1987) Ventosa *et al.* 1999, comb. nov. is the type species of the genus *Haloterrigena* in the euryarchaeal family *Halobacteriaceae*. It is of phylogenetic interest because of the yet unclear position of the genera *Haloterrigena* and *Natri-nema* within the *Halobacteriaceae*, which created some taxonomic problems historically. *H. turkmenica*, was isolated from sulfate saline soil in Turkmenistan, is a relatively fast growing, chemoorganotrophic, carotenoid-containing, extreme halophile, requiring at least 2 M NaCl for growth. Here we describe the features of this organism, together with the complete genome sequence, and annotation. This is the first complete genome sequence of the genus *Haloterrigena*, but the eighth genome sequence from a member of the family *Halobacteriaceae*. The 5,440,782 bp genome (including six plasmids) with its 5,287 protein-coding and 63 RNA genes is part of the *Genomic Encyclopedia of Bacteria and Archaea* project.

---

### Introduction

Strain 4k<sup>T</sup> (= DSM 5511 = ATCC 51198 = VKM B-1734) is the type strain of the species *Haloterrigena turkmenica*, which is the type species of the genus *Haloterrigena* [1,2]. The strain was initially described in 1987 as *Halococcus turkmenicus* VKM B-1734 (basonym) by Zvyagintseva and Tarasov [3]. In 1999, Ventosa *et al.* proposed to transfer *H. turkmenicus* 4k as the type strain of the species *H. turkmenica* to the new genus *Haloterrigena* [1], whose name means salt, *halos*, (-requiring) and

born from the earth, *terrigena*. Inconsistent data published on sequence similarity and DNA-DNA hybridization for some *Haloterrigena* and *Natri-nema* strains created some confusion and taxonomic problems initially, but the problems were largely resolved in 2003 by Tindall [4], pointing to uncertainty about strain history. It has been suggested that the discrepancies may also be a result of 16S rDNA interoperon heterogeneity [5]. Published data appears to indicate that both strains


GSL-11 and JCM 9743 (formally included in the species *H. turkmenica* by Ventosa *et al.* [1]) may be members of the genus *Natrinema* [4,6]. Those strains will not be considered further here.

There are no reliable reports of other strains of *H. turkmenica* having been isolated. 16S rRNA sequence identity with the other seven type strains in the genus, which were mainly isolated from salt lakes, range from 98.0% for *H. salina* [7] to 94.4% for *H. longa* [6]. The sequence similarity to the *Natrinema* type strains is somewhere in-between, 95.2-96.4% [8], underlining the taxonomic problems [4]. The sequence similarity to phylotypes in environmental metagenomic libraries was not above 87%, indicating a rather poor representation of closely related strains in the habitats analyzed (status January 2010). Here we present a summary classification and a set of features for *H. turkmenica* strain 4k<sup>T</sup>, together with the descrip-

tion of the complete genome sequencing and annotation.

## Classification and features


Figure 1 shows the phylogenetic neighborhood of *H. turkmenica* strain 4k<sup>T</sup> in a 16S rRNA based tree. The three 16S rRNA gene sequences in the genome differ from each other by up to two nucleotides, and differ by up to six nucleotides from the previously published 16S rRNA sequence (AB004878) generated from DSM 5511. The difference between the genome data and the previously reported 16S rRNA gene sequences is most likely due to sequencing errors in the previously reported sequence data. As expected, *Haloterrigena* and *Natrinema* strains appear as intermixed in the tree, indicating a paraphyletic status of *Haloterrigena* (within which *Natronorubrum* and *Natrinema* branch off) and of *Natrinema* (within which *H. longa* is placed) [18].



**Figure 1.** Phylogenetic tree highlighting the position of *H. turkmenica* strain 4k relative to the other species within the genera *Haloterrigena* and *Natrinema* and the type strains of the other genera within the family *Halobacteriaceae*. The tree was inferred from 1,368 aligned characters [9,10] of the 16S rRNA sequence under the maximum likelihood criterion [11] and rooted with *Natronomonas pharaonis* [12]. The branches are scaled in terms of the expected number of substitutions per site. Numbers above branches are support values from 800 bootstrap replicates [13] if larger than 60%. Strains with a genome sequencing project registered in GOLD [14] are printed in blue; published genomes in bold, e.g. the recently published GEBA genomes from *Halogeometricum borinquense* [15], *Halorhabdus utahensis* [16], and *Halomicrobium mukohataei* [17].

*H. turkmenica* cells occur mostly as single cells, rarely in pairs or tetrads [1]. They are described as Gram-negative, ovoid to coccoid, 1.5-2  $\mu\text{m}$  in diameter [1], but can also be rod-shaped (Figure 2 and Table 1) [1]. Neither spores, nor flagella, nor lipid granules were reported. Colonies are pigmented red or light pink due of the presence of  $C_{50}$ -carotenoids [1]. Strain 4k<sup>T</sup> is chemoorganotrophic and aerobic, and requires at least 2 M NaCl [1]. Detailed physiological characteristics were

described by Zvyagintseva and Tarasov [3]. The G+C content of DNA was reported to be 59.2-60.2 mol % (Thermal denaturation method [1]), which is significantly less than the 64.3% found in the genome. At optimal growth temperatures, *H. turkmenica* is the fastest growing member of the *Halobacteriaceae*, with only 1.5 hours generation time [26]. Besides the chemical characterization of siderophores [29], there are no published reports on the molecular biology of *H. turkmenica*.



**Figure 2.** Scanning electron micrograph of *H. turkmenica* strain 4k<sup>T</sup>

Both diphytanyl moieties ( $C_{20}$ ,  $C_{20}$ ) and phytanyl-esterterpanyl moieties ( $C_{20}$ ,  $C_{25}$ ) are present in polar lipids [1]. The presence of both phytanyl and esterterpanyl side chains implies the presence of three different prenyl transferases involved in lipid biosynthesis, which are probably chain length specific as well as stereospecific for the incorporation of the isoprenoid side chains into the glycerol backbone [30]. The presence of significant levels of both the diphytanyl moieties ( $C_{20}$ ,  $C_{20}$ ) and phytanyl-esterterpanyl moieties ( $C_{20}$ ,  $C_{25}$ ) is characteristic of all members examined of this evolutio-

nary branch of the family *Halobacteriaceae*. Membrane polar lipids are glycerol-diether analogues of PG, PGP-Me and the disulfated diglycosyl diether lipid  $S_2$ -DGD (mannose-2,6 disulfate 1 $\rightarrow$ 2 glucose-glycerol diether) [31], the characteristic glycolipid of *Natrialba asiatica* [32]. The presence of respiratory lipoquinones have not been reported, but it may be predicted that MK-8 and MK-8 (VIII-H<sub>2</sub>) should be present, since this is a feature of all members of the family *Halobacteriaceae* examined to date.

**Table 1.** Classification and general features of *H. turkmenica* 4k<sup>T</sup> according to the MIGS recommendations [19]

| MIGS ID  | Property               | Term   | Evidence code |
|----------|------------------------|--|---------------|
|          |                        | Domain <i>Archaea</i>                              | TAS [20]      |
|          |                        | Phylum <i>Euryarchaeota</i>                        | TAS [21,22]   |
|          |                        | Class <i>Halobacteria</i>                          | TAS [23]      |
|          | Current classification | Order <i>Halobacteriales</i>                       | TAS [24]      |
|          |                        | Family <i>Halobacteriaceae</i>                     | TAS [25]      |
|          |                        | Genus <i>Haloterrigena</i>                         | TAS [1]       |
|          |                        | Species <i>Haloterrigena turkmenica</i>            | TAS [1]       |
|          |                        | Type strain 4k                                     | TAS [3]       |
|          | Gram stain             | negative   | TAS [1]       |
|          | Cell shape             | rods   | TAS [1]       |
|          | Motility               | nonmotile  | IDA           |
|          | Sporulation            | non-sporulating                                    | NAS           |
|          | Temperature range      | 29-57°C  | TAS [26]      |
|          | Optimum temperature    | 51°C   | TAS [26]      |
|          | Salinity               | extreme halophile, requires at least 2% (w/v) NaCl | TAS [1]       |
| MIGS-22  | Oxygen requirement     | aerobic  | TAS [1]       |
|          | Carbon source          | yeast extract                                      | NAS           |
|          | Energy source          | chemoorganotroph                                   | TAS [1]       |
| MIGS-6   | Habitat                | soil   | TAS [1]       |
| MIGS-15  | Biotic relationship    | free living  | NAS           |
| MIGS-14  | Pathogenicity          | none   | NAS           |
|          | Biosafety level        | 1  | TAS [27]      |
|          | Isolation              | sulfate saline soil                                | TAS [3]       |
| MIGS-4   | Geographic location    | Ashkhabad, Turkmenistan                            | TAS [3]       |
| MIGS-5   | Sample collection time | about or before 1987                               | TAS [3]       |
| MIGS-4.1 | Latitude,              | 37.950,  | NAS           |
| MIGS-4.2 | Longitude              | 58.380   |               |
| MIGS-4.3 | Depth                  | unknown  |               |
| MIGS-4.4 | Altitude               | unknown  |               |

Evidence codes - IDA: Inferred from Direct Assay (first time in publication); TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from of the Gene Ontology project [28]. If the evidence code is IDA, then the property was directly observed by one of the authors or an expert mentioned in the acknowledgements.

## Genome sequencing and annotation

### Genome project history

This organism was selected for sequencing on the basis of its phylogenetic position, and is part of the *Genomic Encyclopedia of Bacteria and Archaea* project [33]. The genome project is deposited in the Genomes OnLine Database [14] and the

complete genome sequence in GenBank. Sequencing, finishing and annotation were performed by the DOE Joint Genome Institute (JGI). A summary of the project information is shown in Table 2.

**Table 2.** Genome sequencing project information

| MIGS ID   | Property                   | Term  |
|-----------|----------------------------|---|
| MIGS-31   | Finishing quality          | Finished  |
| MIGS-28   | Libraries used             | Three genomic libraries: one Sanger 8 kb pMCL200, one 454 pyrosequence standard library and one Illumina standard library |
| MIGS-29   | Sequencing platforms       | ABI3730, 454 GS FLX, and Illumina GA  |
| MIGS-31.2 | Sequencing coverage        | 6.9× Sanger; 19.9× pyrosequence   |
| MIGS-30   | Assemblers                 | Newbler version 1.1.03.24, phrap  |
| MIGS-32   | Gene calling method        | Prodigal 1.4, GenePRIMP   |
|           | Genbank ID                 | CP001860 (chromosome)<br>CP001861-CP001866 (plasmids)   |
|           | Genbank Date of Release    | January 19, 2010  |
|           | GOLD ID                    | Gc01189   |
|           | NCBI project ID            | 30411   |
|           | Database: IMG-GEBA         | 2501939622  |
| MIGS-13   | Source material identifier | DSM 5511  |
|           | Project relevance          | Tree of Life, GEBA  |

### Growth conditions and DNA isolation

*H. turkmenica* 4k<sup>T</sup>, DSM 5511, was grown in DSMZ medium 372 (*Halobacteria* medium) [34] at 37°C. DNA was isolated from 1-1.5 g of cell paste using Qiagen Genomic 500 DNA Kit (Qiagen, Hilden, Germany) with lysis modification L according to Wu *et al.* [33].

### Genome sequencing and assembly

The genome was sequenced using a combination of Sanger and 454 sequencing platforms. All general aspects of library construction and sequencing performed at the JGI can be found at the [JGI website](#). 454 Pyrosequencing reads were assembled using the Newbler assembler version 1.1.03.24 (Roche). Large Newbler contigs were broken into 6,060 overlapping fragments of 1,000 bp and entered into assembly as pseudo-reads. The sequences were assigned quality scores based on Newbler consensus q-scores with modifications to account for overlap redundancy and adjust inflated q-scores. A hybrid 454/Sanger assembly was made using the parallel phrap assembler (High Performance Software, LLC). Possible misassemblies were corrected with Dupfinisher or transposon bombing of bridging clones [35]. A total of 1,183 Sanger finishing reads were produced to close gaps, to resolve repetitive regions, and to raise the quality of the finished sequence. Illumina reads were used to improve the final consensus quality using an in-house developed tool (the Polisher). The error rate of the completed genome sequence is less than 1 in 100,000.

Together, the combination of the Sanger and 454 sequencing platforms provided 26.8× coverage of the genome. The final assembly contains 33,433 Sanger reads and 394,632 pyrosequencing reads.

### Genome annotation


Genes were identified using Prodigal [36] as part of the Oak Ridge National Laboratory genome annotation pipeline, followed by a round of manual curation using the JGI GenePRIMP pipeline [37]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) nonredundant database, UniProt, TIGR-Fam, Pfam, PRIAM, KEGG, COG, and InterPro databases. Additional gene prediction analysis and functional annotation was performed within the [Integrated Microbial Genomes - Expert Review](#) platform [38].

### Genome properties


The genome is 5,440,782 bp long and comprises one main circular chromosome of 3,889,038 bp length and six circular plasmids of 15.8 to 698.5 kbp length, with an overall GC content of 64.3% (Table 3 and Figures 3 and 4). Of the 5,350 genes predicted, 5,287 were protein coding genes, and 63 RNAs; 174 pseudogenes were also identified. The majority of the protein-coding genes (60.1%) were assigned a putative function while those remaining were annotated as hypothetical proteins. The distribution of genes into COGs functional categories is presented in Table 4.

**Table 3.** Genome Statistics

| Attribute                        | Value     | % of Total |
|----------------------------------|-----------|------------|
| Genome size (bp)                 | 5,440,782 | 100.00%    |
| DNA coding region (bp)           | 4,524,412 | 83.16%     |
| DNA G+C content (bp)             | 3,496,479 | 64.26%     |
| Number of replicons              | 7         |            |
| Extrachromosomal elements        | 6         |            |
| Total genes                      | 5,350     | 100.00%    |
| RNA genes                        | 63        | 1.18%      |
| rRNA operons                     | 3         |            |
| Protein-coding genes             | 5,287     | 98.82%     |
| Pseudo genes                     | 174       | 3.25%      |
| Genes with function prediction   | 3,213     | 60.06%     |
| Genes in paralog clusters        | 1,706     | 31.89%     |
| Genes assigned to COGs           | 3,259     | 60.92%     |
| Genes assigned Pfam domains      | 3,208     | 59.96%     |
| Genes with signal peptides       | 625       | 11.68%     |
| Genes with transmembrane helices | 1,140     | 21.31%     |
| CRISPR repeats                   | 1         |            |



**Figure 3.** Graphical circular map of the chromosome. From outside to the center: Genes on forward strand (color by COG categories), Genes on reverse strand (color by COG categories), RNA genes (tRNAs green, rRNAs red, other RNAs black), GC content, GC skew.



**Figure 4.** Graphical circular map of the six plasmids: pHTUR01 (A), pHTUR02 (B), pHTUR03 (C), pHTUR04 (D), pHTUR05 (E), pHTUR06 (F). Plasmids not drawn to scale.

**Table 4.** Number of genes associated with the general COG functional categories

| Code | Value | %age | Description  |
|------|-------|------|--|
| J    | 178   | 3.4  | Translation, ribosomal structure and biogenesis              |
| A    | 1     | 0.0  | RNA processing and modification                              |
| K    | 190   | 3.6  | Transcription  |
| L    | 150   | 2.8  | Replication, recombination and repair                        |
| B    | 3     | 0.1  | Chromatin structure and dynamics                             |
| D    | 35    | 0.7  | Cell cycle control, mitosis and meiosis                      |
| Y    | 0     | 0.0  | Nuclear structure  |
| V    | 44    | 0.8  | Defense mechanisms   |
| T    | 161   | 3.0  | Signal transduction mechanisms                               |
| M    | 125   | 2.4  | Cell wall/membrane biogenesis                                |
| N    | 29    | 0.5  | Cell motility  |
| Z    | 0     | 0.0  | Cytoskeleton   |
| W    | 0     | 0.0  | Extracellular structures                                     |
| U    | 26    | 0.5  | Intracellular trafficking and secretion                      |
| O    | 141   | 2.7  | Posttranslational modification, protein turnover, chaperones |
| C    | 258   | 4.9  | Energy production and conversion                             |
| G    | 221   | 4.2  | Carbohydrate transport and metabolism                        |
| E    | 349   | 6.6  | Amino acid transport and metabolism                          |
| F    | 78    | 1.5  | Nucleotide transport and metabolism                          |
| H    | 189   | 3.6  | Coenzyme transport and metabolism                            |
| I    | 176   | 3.3  | Lipid transport and metabolism                               |
| P    | 224   | 4.2  | Inorganic ion transport and metabolism                       |
| Q    | 87    | 1.6  | Secondary metabolites biosynthesis, transport and catabolism |
| R    | 630   | 11.9 | General function prediction only                             |
| S    | 321   | 6.1  | Function unknown   |
| -    | 2,091 | 39.5 | Not in COGs  |



## Acknowledgements

We would like to gratefully acknowledge the help of Susanne Schneider (DSMZ) for DNA extraction and quality analysis. This work was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley

National Laboratory under contract No. DE-AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, and Los Alamos National Laboratory under contract No. DE-AC02-06NA25396, as well as German Research Foundation (DFG) INST 599/1-1.

## References

1. Ventosa A, Gutiérrez MC, Kamekura M, Dyll-Smith ML. Proposal to transfer *Halococcus turkmenicus*, *Halobacterium trapanicum* JCM 9743 and strain GSL-11 to *Haloterrigena turkmenica* gen. nov., comb. nov. *Int J Syst Bacteriol* 1999; **49**:131-136. [PubMed](#)
2. Oren A, Arahal DR, Ventosa A. Emended descriptions of the genera of the family *Halobacteriaceae*. *Int J Syst Evol Microbiol* 2009; **59**:637-642. [PubMed](#) [doi:10.1099/ijs.0.008904-0](#)
3. Zvyagintseva IS, Tarasov AL. Extreme halophilic bacteria from saline soils. *Mikrobiologiya* 1987; **56**:839-844.
4. Tindall BJ. Taxonomic problems arising in the genera *Haloterrigena* and *Natrinema*. *Int J Syst Evol Microbiol* 2003; **53**:1697-1698. [PubMed](#) [doi:10.1099/ijs.0.02529-0](#)
5. Walsh DA, Baptiste E, Kamekura M, Doolittle WF. Evolution of the RNA polymerase B subunit Gene (*rpoB*) in *Halobacteriales*: a complementary molecular marker to the SSU rRNA Gene. *Mol Biol Evol* 2004; **21**:2340-2351. [PubMed](#) [doi:10.1093/molbev/msh248](#)
6. Cui HL, Tohty D, Zhou PJ, Liu SJ. *Haloterrigena longa* sp. nov. and *Haloterrigena limicola* sp. nov., extremely halophilic archaea isolated from a salt lake. *Int J Syst Evol Microbiol* 2006; **56**:1837-1840. [PubMed](#) [doi:10.1099/ijs.0.64372-0](#)
7. Gutiérrez MC, Castillo AM, Kamekura M, Ventosa A. *Haloterrigena salina* sp. nov., an extremely halophilic archaeon isolated from a salt lake. *Int J Syst Evol Microbiol* 2008; **58**:2880-2884. [PubMed](#) [doi:10.1099/ijs.0.2008/001602-0](#)
8. Chun J, Lee JH, Jung Y, Kim M, Kim S, Kim BK, Lim YW. EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. *Int J Syst Evol Microbiol* 2007; **57**:2259-2261. [PubMed](#) [doi:10.1099/ijs.0.64915-0](#)
9. Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; **17**:540-552. [PubMed](#)
10. Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics* 2002; **18**:452-464. [PubMed](#) [doi:10.1093/bioinformatics/18.3.452](#)
11. Stamatakis A, Hoover P, Rougemont J. A Rapid Bootstrap Algorithm for the RAxML Web Servers. *Syst Biol* 2008; **57**:758-771. [PubMed](#) [doi:10.1080/10635150802429642](#)
12. Falb M, Pfeiffer F, Palm P, Rodewald K, Hickmann V, Tittor J, Oesterhelt D. Living with two extremes: conclusions from the genome sequence of *Natronomonas pharaonis*. *Genome Res* 2005; **15**:1336-1343. [PubMed](#) [doi:10.1101/gr.3952905](#)
13. Pattengale ND, Alipour M, Bininda-Emonds ORP, Moret BME, Stamatakis A. How many bootstrap replicates are necessary? *Lect Notes Comput Sci* 2009; **5541**:184-200. [doi:10.1007/978-3-642-02008-7\\_13](#)
14. Liolios K, Chen IM, Mavromatis K, Tavernarakis N, Hugenholtz P, Markowitz VM, Kyrpides NC. The Genomes On Line Database (GOLD) in 2009: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res* 2010; **38**:D346-D354. [PubMed](#) [doi:10.1093/nar/gkp848](#)
15. Malfatti S, Tindall BJ, Schneider S, Fährnich R, Lapidus A, LaButtii K, Copeland A, Glavina Del Rio T, Nolan M, Chen F, et al. Complete genome sequence of *Halogeometricum borinquense* type strain (PR3<sup>T</sup>). *Stand Genomic Sci* 2009; **1**:150-158. [doi:10.4056/sigs.23264](#)
16. Anderson I, Tindall BJ, Pomrenke H, Göker M, Lapidus A, Nolan M, Copeland A, Glavina Del Rio T, Chen F, Tice H, et al. Complete genome sequence of *Halorhabdus utahensis* type strain (AX-2<sup>T</sup>). *Stand Genomic Sci* 2009; **1**:218-225. [doi:10.4056/sigs.31864](#)
17. Tindall BJ, Schneider S, Lapidus A, Copeland A, Glavina Del Rio T, Nolan M, Lucas S, Chen F, Tice H, Cheng JF, et al. Complete genome

- sequence of *Halomicrobium mukohataei* type strain (arg-2<sup>T</sup>). *Stand Genomic Sci* 2009; **1**:270-277. [doi:10.4056/signs.42644](https://doi.org/10.4056/signs.42644)
18. Farris JS. Formal definitions of paraphyly and polyphyly. *Syst Zool* 1974; **23**:548-554. [doi:10.2307/2412474](https://doi.org/10.2307/2412474)
  19. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thompson N, Allen MJ, Angiuoli SV, et al. Towards a richer description of our complete collection of genomes and metagenomes: the "Minimum Information about a Genome Sequence" (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](https://pubmed.ncbi.nlm.nih.gov/16811111/) [doi:10.1038/nbt1360](https://doi.org/10.1038/nbt1360)
  20. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](https://pubmed.ncbi.nlm.nih.gov/18711111/) [doi:10.1073/pnas.87.12.4576](https://doi.org/10.1073/pnas.87.12.4576)
  21. Garrity GM, Holt JG. Phylum All. *Euryarchaeota* ph. nov. In: *Bergey's Manual of Systematic Bacteriology*, vol. 1. 2nd ed. Edited by: Garrity, GM, Boone, DR and Castenholz, RW. Springer, New York; 2001:211-355.
  22. List Editor. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. Validation List no. 85. *Int J Syst Evol Microbiol* 2002; **52**:685-690. [PubMed](https://pubmed.ncbi.nlm.nih.gov/12311111/) [doi:10.1099/ijs.0.02358-0](https://doi.org/10.1099/ijs.0.02358-0)
  23. Garrity GM, Lilburn TG, Cole JR, Harrison SH, Euzéby J, Tindall BJ. "Part 1- The *Archaea*, Phyla *Crenarchaeota* and *Euryarchaeota*" Taxonomic Outline of the *Bacteria* and *Archaea*. 2007. [www.taxonomicoutline.org](http://www.taxonomicoutline.org).
  24. Grant WD, Larsen H. Group III. Extremely halophilic archaeobacteria. Order *Halobacteriales* ord. nov. In: Staley JT, Bryant MP, Pfennig N & Holt JG(eds) *Bergey's Manual of Systematic Bacteriology*, First edition, Vol 3, The Williams & Watkins Co., Baltimore, 1989, pp 2216-2228.
  25. Gibbons NE. Family V. *Halobacteriaceae* fam. nov. In: Buchanan RE & Gibbons NE(eds) *Bergey's Manual of Determinative Bacteriology*, eighth edition, The Williams & Watkins Co., Baltimore, 1974, pp 279.
  26. Robinson JL, Pyzyna B, Atrasz RG, Handerson CA, Morrill KL, Burd AM, Desoucy E, Fogleman RE, III, Naylor JB, Steele SM, et al. Growth kinetics of extremely halophilic *Archaea* (family *Halobacteriaceae*) as revealed by Arrhenius plots. *J Bacteriol* 2005; **187**:923-929. [PubMed](https://pubmed.ncbi.nlm.nih.gov/15611111/) [doi:10.1128/JB.187.3.923-929.2005](https://doi.org/10.1128/JB.187.3.923-929.2005)
  27. Biological Agents. Technical rules for biological agents [www.baua.de TRBA 466](http://www.baua.de/TRBA/466).
  28. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene ontology: tool for the unification of biology. *Nat Genet* 2000; **25**:25-29. [PubMed](https://pubmed.ncbi.nlm.nih.gov/103875556/) [doi:10.1038/75556](https://doi.org/10.1038/75556)
  29. Dave BP, Anshuman K, Hajela P. Siderophores of halophilic *Archaea* and their chemical characterization. *Indian J Exp Biol* 2006; **44**:340-344. [PubMed](https://pubmed.ncbi.nlm.nih.gov/16811111/)
  30. Tachibana A. A novel prenyltransferase, farnesylgeranyl diphosphate synthase, from the haloalkaliphilic archaeon, *Natronobacterium pharaonis*. *FEBS Lett* 1994; **341**:291-294. [PubMed](https://pubmed.ncbi.nlm.nih.gov/10160014579394804753/) [doi:10.1016/0014-5793\(94\)80475-3](https://doi.org/10.1016/0014-5793(94)80475-3)
  31. Kamekura M, Dyll-Smith ML, Upasani V, Ventosa A, Kates M. Diversity of alkaliphilic halobacteria: proposals for the transfer of *Natronobacterium vacuolatum*, *Natronobacterium magadii*, and *Natronobacterium pharaonis* to the genus *Halorubrum*, *Natrialba*, and *Natronomonas* gen. nov., respectively, as *Halorubrum vacuolatum* comb. nov., *Natrialba magadii* comb. nov., and *Natronomonas pharaonis* comb. nov., respectively. *Int J Syst Bacteriol* 1997; **47**:853-857. [PubMed](https://pubmed.ncbi.nlm.nih.gov/91111111/)
  32. Kamekura M, Dyll-Smith ML. Taxonomy of the family *Halobacteriaceae* and the description of two new genera *Halorubrobacterium* and *Natrialba*. *J Gen Appl Microbiol* 1995; **41**:333-350. [doi:10.2323/jgam.41.333](https://doi.org/10.2323/jgam.41.333)
  33. Wu D, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova N, Kunin V, Goodwin L, Wu M, Tindall BJ. A phylogeny-driven genomic encyclopedia of *Bacteria* and *Archaea*. *Nature* 2009; **462**:1056-1060. [PubMed](https://pubmed.ncbi.nlm.nih.gov/1038nature08656/) [doi:10.1038/nature08656](https://doi.org/10.1038/nature08656)
  34. List of growth media used at DSMZ: [http://www.dsmz.de/microorganisms/media\\_list.php](http://www.dsmz.de/microorganisms/media_list.php).
  35. Sims D, Brettin T, Detter JC, Han C, Lapidus A, Copeland A, Glavina Del Rio T, Nolan M, Chen F, Lucas S, et al. Complete genome of *Kytococcus sedentarius* type strain (541<sup>T</sup>). *Stand Genomic Sci* 2009; **1**:12-20. [doi:10.4056/signs.761](https://doi.org/10.4056/signs.761)
  36. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene

- recognition and translation initiation site identification. *BMC Genomics* (In press).
37. Pati A, Ivanova N, Mikhailova N, Ovchinikova G, Hooper SD, Lykidis A, Kyrpides NC. GenePRIMP: A Gene Prediction Improvement Pipeline for microbial genomes. *Nat Methods* (In press); <http://geneprimp.jgi-psf.org/>.
38. Markowitz VM, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 2009; **25**:2271-2278. [PubMed](https://pubmed.ncbi.nlm.nih.gov/19311111/) [doi:10.1093/bioinformatics/btp393](https://doi.org/10.1093/bioinformatics/btp393)