UC Berkeley

UC Berkeley Previously Published Works

Title

Kinetic modeling of nitrous oxide decomposition on Fe-ZSM-5 in the presence of nitric oxide based on parameters obtained from first-principles calculations

Permalink

<https://escholarship.org/uc/item/9gk9520h>

Journal

Catalysis Science and Technology, 11(10)

ISSN

2044-4753

Authors

Aranifard, S
Bell, AT
Keil, FJ
et al.

Publication Date

2021-05-21

DOI

10.1039/d1cy00252j

Peer reviewed

[Log in / register](#)

SCHEDULED MAINTENANCE

Maintenance work is planned for Wednesday 18 August 2021 from 07:00 to 23:59 (BST).

Pages may load more slowly than usual and you may not be able to access some content or log in. If this happens then please refresh your web browser or try waiting two to three minutes before trying again. Sorry for any inconvenience and thank you for your patience.

[Issue 10, 2021](#)

[Previous](#)

[Next](#)

From the journal:

Catalysis Science & Technology

Kinetic modeling of nitrous oxide decomposition on Fe-ZSM-5 in the presence of nitric oxide based on parameters obtained from first-principles calculations†‡

[Sara Aranifard](#) ^{*a} [Alexis T. Bell](#) ^b [Frerich J. Keil](#) ^c and [Andreas Heyden](#) ^{*a}

[Author affiliations](#)

Abstract

A variety of experiments for the N₂O decomposition over Fe-ZSM-5 catalysts have been simulated in the presence and absence of small amounts of nitric oxide and water vapor. A comprehensive reaction mechanism over mononuclear iron sites was considered, and all elementary reaction rate constants were obtained from density functional theory and transition state theory. Various experimental observations,

such as the acceleration of the N_2O decomposition in the presence of nitric oxide at low temperatures, can be described with the studied reaction mechanism on mononuclear iron sites. No other iron species, such as binuclear iron, are necessary for explaining experimental observations. At high temperatures, $\text{Z}^-\text{[FeO]}^+$ sites are active for N_2O decomposition, forming either $\text{Z}^-\text{[OFeO]}^+$ and $\text{Z}^-\text{[FeO}_2]^+$ sites on which a second N_2O can decompose to form $\text{Z}^-\text{[OFeO}_2]^+$ sites from which O_2 can rapidly desorb. At low temperatures, nitric oxide activates water poisoned $\text{Z}^-\text{[Fe(OH)}_2]^+$ sites to form active $\text{Z}^-\text{[FeOH]}^+$ species. The catalytic cycle on $\text{Z}^-\text{[FeOH]}^+$ involves N_2O dissociation to form $\text{Z}^-\text{[OFeOH]}^+$ sites that are inactive for a second N_2O decomposition. Instead, NO adsorbs and NO_2 desorbs in order to regenerate the $\text{Z}^-\text{[FeOH]}^+$ sites. On all active iron sites, the first N_2O dissociation step is the most critical rate-controlling step. The concentration of nitric oxide and water vapor together determine at which temperature the switch between the most dominant active site (hydroxo- *versus* oxo-iron species) occurs. Overall, this study motivates the investigation of $\text{Z}^-\text{[OFeOH]}^+$ sites as potential α -oxygen species that can oxidize various hydrocarbons at low temperatures.

[About](#)
[Cited by](#)
[Related](#)

Buy this article

£42.50*

* Exclusive of taxes

This article contains 17 page(s)

Other ways to access this content

Log in

Using your institution credentials

Sign in

With your membership or subscriber account

Supplementary files

Supplementary information

PDF (320K)

Article information

<https://doi.org/10.1039/D1CY00252J>

Submitted

09 Feb 2021

Accepted

31 Mar 2021

First published

31 Mar 2021

Citation

Catal. Sci. Technol., 2021, **11**, 3539-3555

BibTex

Go

Article type

Paper

Permissions

[Request permissions](#)

Social activity

Tweet

Share

Search articles by author

- ☐ Sara Aranifard
- ☐ Alexis T. Bell
- ☐ Frerich J. Keil
- ☐ Andreas Heyden

Go

Spotlight

Advertisements

» Journals, books & databases

- Home
- About us
- Membership & professional community
- Campaigning & outreach
- Journals, books & databases
- Teaching & learning
- News & events
- Locations & contacts
- Careers
- Awards & funding
- Advertise
- Help & legal
- Privacy policy
- Terms & conditions

© Royal Society of Chemistry 2021

Registered charity number: 207890