

UCSF

UC San Francisco Previously Published Works

Title

A candidate gene study of the type I interferon pathway implicates IKBKE and IL8 as risk loci for SLE

Permalink

<https://escholarship.org/uc/item/9gt3v3mz>

Journal

European Journal of Human Genetics, 19(4)

ISSN

1018-4813

Authors

Sandling, Johanna K
Garnier, Sophie
Sigurdsson, Snaevar
et al.

Publication Date

2011-04-01

DOI

10.1038/ejhg.2010.197

Peer reviewed

ARTICLE

A candidate gene study of the type I interferon pathway implicates *IKBKE* and *IL8* as risk loci for SLE

Johanna K Sandling¹, Sophie Garnier^{1,12}, Snaevor Sigurdsson^{1,13}, Chuan Wang¹, Gunnel Nordmark², Iva Gunnarsson³, Elisabet Svenungsson³, Leonid Padyukov³, Gunnar Sturfelt⁴, Andreas Jönsen⁴, Anders A Bengtsson⁴, Lennart Truedsson⁵, Catharina Eriksson⁶, Solbritt Rantapää-Dahlqvist⁷, Anders Målarstig⁸, Rona J Strawbridge⁸, Anders Hamsten⁸, Lindsey A Criswell⁹, Robert R Graham¹⁰, Timothy W Behrens¹⁰, Maija-Leena Eloranta², Gunnar Alm¹¹, Lars Rönnblom² and Ann-Christine Syvänen^{*1}

Systemic Lupus Erythematosus (SLE) is a systemic autoimmune disease in which the type I interferon pathway has a crucial role. We have previously shown that three genes in this pathway, *IRF5*, *TYK2* and *STAT4*, are strongly associated with risk for SLE. Here, we investigated 78 genes involved in the type I interferon pathway to identify additional SLE susceptibility loci. First, we genotyped 896 single-nucleotide polymorphisms in these 78 genes and 14 other candidate genes in 482 Swedish SLE patients and 536 controls. Genes with $P < 0.01$ in the initial screen were then followed up in 344 additional Swedish patients and 1299 controls. SNPs in the *IKBKE*, *TANK*, *STAT1*, *IL8* and *TRAF6* genes gave nominal signals of association with SLE in this extended Swedish cohort. To replicate these findings we extracted data from a genomewide association study on SLE performed in a US cohort. Combined analysis of the Swedish and US data, comprising a total of 2136 cases and 9694 controls, implicates *IKBKE* and *IL8* as SLE susceptibility loci ($P_{\text{meta}} = 0.00010$ and $P_{\text{meta}} = 0.00040$, respectively). *STAT1* was also associated with SLE in this cohort ($P_{\text{meta}} = 3.3 \times 10^{-5}$), but this association signal appears to be dependent of that previously reported for the neighbouring *STAT4* gene. Our study suggests additional genes from the type I interferon system in SLE, and highlights genes in this pathway for further functional analysis.

European Journal of Human Genetics (2011) 19, 479–484; doi:10.1038/ejhg.2010.197; published online 22 December 2010

Keywords: systemic lupus erythematosus; type I interferon system; candidate gene study; single nucleotide polymorphism; *IKBKE*; *IL8*

INTRODUCTION

Systemic Lupus Erythematosus (SLE, OMIM 152700) is an inflammatory autoimmune disease that primarily affects women during their childbearing years. Production of autoantibodies, tissue deposits of immune complexes and inflammation in kidneys, skin, joints and central nervous system are hallmarks of SLE. Despite a strong heritability of the disease, linkage studies have failed to identify genes outside the major histocompatibility complex (MHC) region as risk factors for SLE. Association studies have since proven a more fruitful approach. The confirmed findings from genomewide association studies (GWAS) include genes that were originally discovered in candidate gene studies, such as the interferon regulatory factor 5 (*IRF5*)¹ and the signal transducer and activator of transcription 4 (*STAT4*)² genes from the type I IFN system. IFNs are cytokines with antiviral activity that are produced in response to viral infections, of which the type I IFNs bind the IFN- α receptor (IFNAR). Today there

are more than 20 confirmed SLE susceptibility loci,^{3,4} of which several are in the type I IFN system.

The type I IFN system is activated in SLE patients,^{5,6} and an important role of the type I IFN system in the disease process was confirmed by studies showing an increased expression of type I IFN-inducible genes in SLE patients (an 'IFN-signature').⁷ A direct causative role of the type I IFN system in the etiopathogenesis of SLE was suggested by the observation that individuals treated with IFN- α can develop an SLE syndrome indistinguishable from the naturally occurring disease.⁸ Moreover, a phase I clinical trial with a monoclonal antibody against IFN- α , reports reduction of disease activity as well as neutralization of the IFN signature in SLE patients.^{9,10} Encouraged by the compelling evidence for the involvement of the type I IFN system in SLE, we performed an association study to identify additional genes from the type I IFN pathway that confer risk for SLE.

¹Molecular Medicine, Department of Medical Sciences, Uppsala University, Uppsala, Sweden; ²Section of Rheumatology, Department of Medical Sciences, Uppsala University, Uppsala, Sweden; ³Rheumatology Unit, Department of Medicine, Karolinska Institutet/Karolinska University Hospital, Stockholm, Sweden; ⁴Section of Rheumatology, Department of Clinical Sciences, Lund University, Lund, Sweden; ⁵Department of Laboratory Medicine, section of M.I.G., Lund University, Lund, Sweden; ⁶Department of Clinical Immunology, Umeå University Hospital, Umeå, Sweden; ⁷Department of Rheumatology, Umeå University Hospital, Umeå, Sweden; ⁸Atherosclerosis Research Unit, Department of Medicine, Solna Karolinska Institutet Stockholm, Stockholm, Sweden; ⁹Rosalind Russell Medical Research Center for Arthritis, Department of Medicine, University of California, San Francisco, CA, USA; ¹⁰Genentech Inc., South San Francisco, CA, USA; ¹¹Department of Biomedical Sciences and Veterinary Public Health, Swedish University of Agricultural Sciences, Uppsala, Sweden

*Correspondence: Professor A-C Syvänen, Department of Medical Sciences, Uppsala University, Molecular Medicine, Entrance 70, Research Department 2, 3rd floor, Uppsala University Hospital, 75185 Uppsala, Sweden. Tel: +46 18 6112959; Fax: +46 18 55 36 01; E-mail: Ann-Christine.Syvanen@medsci.uu.se

¹²Current address: INSERM UMR S937, Génétique Épidémiologique et Moléculaire des Pathologies Cardiovasculaires, Université Paris VI, Paris, France.

¹³Current address: Broad Institute of Harvard and MIT, Boston, MA, USA.

Received 5 May 2010; revised 9 August 2010; accepted 8 October 2010; published online 22 December 2010

MATERIALS AND METHODS

Subjects and genotyping

Our study included 826 Swedish and 1310 US SLE patients, fulfilling at least four of the classification criteria for SLE as defined by the American College of Rheumatology (ACR),¹¹ and 9694 healthy control individuals from the same geographic areas as the patients (Supplementary Table S1). DNA was extracted from blood samples of the patients and controls using standard procedures. The study was approved by the regional ethical boards and all subjects gave their informed consent to participate. The study was performed in three stages:

Discovery phase. First, a panel of SNPs in 82 genes with key functions related to the type I IFN signalling system and 14 additional genes, with suggested association with SLE, were selected for genotyping in 490 Swedish SLE patients and 543 controls. Patients were from the rheumatology clinics at the Lund, Karolinska (Solna) and Uppsala University Hospitals in Sweden. SNPs were selected on the basis of their average spacing of around 1 kb and LD information from the HapMap project ($r^2 < 0.8$ HapMap CEU release 16c), excluding SNPs with an Illumina quality score < 0.6 . Genotyping of 1258 SNPs in the 96 genes was performed using the Golden Gate Assay (Illumina Inc., San Diego, CA, USA). Samples and SNPs with $> 10\%$ missing data, SNPs with Hardy–Weinberg equilibrium test P -values < 0.001 and SNPs with MAF < 0.01 were excluded from further analysis. Four parent–offspring trios were included in the genotyping for inheritance checks, and no Mendelian inheritance errors were observed. After exclusion of genetic outliers, duplicate or related samples 482 cases, 536 controls and 896 SNPs in 92 genes were available for analysis.

Confirmatory set. In the second phase, SNPs in the eight most promising genes were followed up in 393 patients and 1645 controls from Sweden. Patients were from the rheumatology clinics at the Umeå, Uppsala and Karolinska (Solna) University Hospitals. For the patients and 972 of the controls, genotyping of 25 SNPs was performed using the 12-plex and 48-plex SNPstream systems¹² (Beckman Coulter Inc., Brea, CA, USA). Primer sequences are provided in Supplementary Table S2. The same quality control filters as for the discovery cohort were applied, and samples overlapping with the discovery phase were excluded. Additional Swedish population-based controls from the Stockholm area ($n=673$), previously genotyped using the Infinium II assay on human 1M v1 bead arrays¹³ (Illumina Inc.), were also included after applying the following quality filters: MAF $> 1\%$, HWE $P > 1 \times 10^{-6}$, SNP and sample call rates $> 95\%$. After exclusion of genetic outliers, duplicate or related samples, a total of 344 cases, 1299 controls and 21 SNPs were available for analysis.

GWAS. We sought replication of our results by using data from a GWAS on SLE in US Caucasians.¹⁴ In brief, 1435 North-American SLE cases of European descent and 3583 controls had been genotyped on HumanHap550 bead arrays (Illumina Inc.). An additional 4564 controls were also included as previously described.⁴ After strict quality control¹⁴ 1310 cases and 7859 controls remained.

Additional quality control and imputation

For the 673 Swedish controls genotyped on the 1M bead arrays, genotypes for 13 confirmatory phase SNPs that were not directly genotyped were imputed using the software IMPUTE and phasing data from the HapMap project.¹⁵ In the US GWAS dataset genotypes were imputed for 15 SNPs using the IMPUTE software. SNPs had imputation confidence scores ≥ 0.90 with one exception (rs4694178, confidence 0.85 in the US data). For all the Swedish SLE patients, and for more than half of the Swedish controls genotyped using the GoldenGate or SNPstream methods, data from 6060 uncorrelated ancestry informative markers (AIMs) became available during the course of our study.⁴ Using this data, genetic outliers were identified with principal component analysis by the EIGENSTRAT software¹⁶ and excluded from the study (the ten first principal components were inferred and a cut-off of $\sigma > 6$ was used to identify outliers). These samples were also checked for cryptic relatedness by investigation of identity-by-state (IBS) status in PLINK¹⁷ (<http://pngu.mgh.harvard.edu/purcell/plink/>) using a set of 12k previously genotyped SNPs. For the Swedish controls, genotyped on the 1M arrays, all markers were used to identify cryptically related samples and 6035 successfully genotyped AIMs were used to identify genetic outliers.

Association analysis and power calculation

The association analysis of the directly genotyped SNPs was performed by comparing allele frequencies in cases and controls with Fisher's exact or Chi2 tests using PLINK. A null distribution for the quantile–quantile (Q–Q) plot was generated with PLINK and plotted using R (<http://www.r-project.org/>). Analyses including imputed genotypes were performed using SNPTEST,¹⁵ which takes imputation uncertainty into account. SNPs, which were not captured by imputation were only analysed in the directly genotyped samples. Conditional logistic regression analysis assuming an additive model was performed using PLINK to test for independence of association signals observed in neighbouring genes, or between associated SNPs within the same gene. Tests for pairwise SNP interactions were performed using the epistasis command in PLINK. The combined analysis of the Swedish and US case-control data was performed using the software Metal (<http://www.sph.umich.edu/csg/abecasis/Metal/index.html>). Pooled odds ratios were calculated using the Mantel–Haenszel method under a fixed effects model, and tests for heterogeneity of odds ratios between studies were calculated using the MedCalc software (<http://www.medcalc.be/>). Power calculations were performed using the software QUANTO (<http://hydra.usc.edu/GxE/>) assuming a log-additive model, and a prevalence for SLE in Sweden of 0.05%.

RESULTS

We selected a panel of SNPs in 78 genes with key functions related to the type I IFN signalling system to study their association with SLE in a Swedish case-control cohort. The selected genes encode Toll-like receptors (TLRs) and intracellular sensor molecules for nucleic acids (ie RIG-I-like receptors: RLR) and members of their signalling pathways, including several transcription factors that are active in the IFN producing pDCs and membrane proteins of the pDCs. Genes encoding the members of the type I IFN family and other genes regulated by the TLRs that are involved in the response to the type I and type III IFNs, and genes for which the expression is directly regulated by type I IFNs, were also included in the panel. Although the type I and type III IFNs bind two different receptors, the IFNAR and IFN- λ receptor, they share downstream signalling and IFN- λ -induced genes are also induced by type I IFNs. A total of 14 additional genes that are not directly involved in the type I IFN system, but have been suggested to be associated with SLE were also included (Supplementary Table S3). This analysis identified 21 SNPs in seven genes that yielded unadjusted P -values < 0.01 : *IKBKE*, *TANK*, *STAT1*, *IL8*, *NRP1*, *TRAF6* and *PIAS4* (Figure 1 and Supplementary Table S3). At this significance level only nine associated SNPs would be expected to yield association signals by chance, which indicates the presence of true association signals in our data (Supplementary Figure S1). The power for the discovery phase, which included 482 SLE cases and 536 controls, was 65% to detect an OR of 1.5 at 0.01 significance for a 10% frequency allele. However, our power to detect genes with an OR of 1.2 was considerably lower (10%), and thus we cannot exclude that genes, which remained undetected in our study may contribute to the risk for SLE.

To increase power, we designed a panel of SNPs in the *IKBKE*, *TANK*, *STAT1*, *IL8*, *NRP1*, *TRAF6* and *PIAS4* genes for genotyping in an independent collection of Swedish SLE patients and controls. The *STAT5B* gene was also included in the follow-up study, as initially one SNP in the gene showed a P -value < 0.01 . This was, however, before an additional quality control step, by which related samples and population outliers were excluded. For genes, where multiple SNPs showed P -values < 0.05 in the discovery phase, candidate SNPs for follow-up were tested for independence of their association signal in relation to the most strongly associated SNP in each gene (Supplementary Table S4). A partly redundant set of 21 SNPs that accounted for the association signals from the discovery phase was then analysed in the second Swedish case-control cohort. Also in this




Figure 1 Association with SLE for the 92 genes tested in the discovery phase (482 Swedish cases and 536 controls). The negative logarithm of the *P*-value from a χ^2 -test is plotted against chromosomal location. The grey line represents $P=0.05$, and the black line $P=0.01$.

Table 1 Combined analysis of association with SLE in the Swedish discovery and confirmation cohorts^a

Gene	CHR	SNP ^b	Basepair	Minor/major alleles	MAF cases	MAF controls	P ^c	Risk allele	OR (95% CI)
<i>IKBKE</i>	1	rs1539243	204 714 410	T/C	0.16	0.18	0.031	C	1.19 (1.02–1.39)
<i>IKBKE</i>	1	rs17433930	204 719 362	G/A	0.07	0.09	0.013	A	1.33 (1.06–1.67)
<i>TANK</i>	2	rs3754974	161 721 201	G/A	0.05	0.04	0.0087	G	1.44 (1.09–1.89)
<i>TANK</i>	2	rs1267075	161 768 063	G/A	0.19	0.18	0.75	G	1.02 (0.88–1.19)
<i>STAT1</i>	2	rs2030171	191 577 408	A/G	0.35	0.29	1.2E–05	A	1.31 (1.16–1.49)
<i>STAT1</i>	2	rs16833172	191 584 314	A/G	0.04	0.03	0.015	A	1.51 (1.09–2.09)
<i>IL8</i>	4	rs4694178	74 831 552	C/A	0.52	0.46	5.2E–05	C	1.26 (1.12–1.42)
<i>NRP1</i>	10	rs734187	33 524 702	A/G	0.24	0.24	0.67	A	1.03 (0.90–1.18)
<i>NRP1</i>	10	rs1331314	33 564 687	G/C	0.11	0.10	0.32	G	1.11 (0.92–1.34)
<i>NRP1</i>	10	rs2073320	33 593 263	A/G	0.38	0.37	0.42	A	1.05 (0.93–1.18)
<i>TRAF6</i>	11	rs5030482	36 466 602	C/T	0.12	0.14	0.0094	T	1.25 (1.05–1.49)
<i>TRAF6</i>	11	rs5030472	36 470 362	T/C	0.09	0.12	0.013	C	1.28 (1.05–1.55)
<i>STAT5B</i>	17	rs6503691	37 647 616	T/C	0.09	0.10	0.44	C	1.08 (0.88–1.32)
<i>PIAS4</i>	19	rs2289863	3 979 783	C/T	0.27	0.29	0.19	T	1.09 (0.96–1.25)

Abbreviations: CI, confidence Interval; MAF, minor Allele Frequency; OR, odds ratio; SNP, single-nucleotide polymorphism.

^a826 Swedish cases and 1835 controls genotyped in the discovery and replication phases including controls genotyped on the 1M chip.

^bGenotypes for rs17433930, rs3754974, rs2030171, rs16833172, rs4694178, rs734187 and rs1331314 were imputed in the Swedish controls genotyped on the 1M chip.

^c*P*-values calculated using SNPtest.

P-values <0.05 are indicated in bold.

confirmatory cohort genetic outliers were excluded. By combining the data from the Swedish discovery and confirmatory cohorts, totalling 826 cases and 1835 controls, we observed strong association signals with $P < 5 \times 10^{-4}$ for the SNPs rs2030171 in *STAT1* and rs4694178 in *IL8* (Table 1 and Supplementary Table S5). Multiple linked SNPs in the TRAF family member-associated NF κ B activator (*TANK*) and TNF receptor-associated factor 6 (*TRAF6*) genes also gave suggestive signals of association with SLE in the combined Swedish cohort. In addition, two weakly correlated SNPs in the *IKBKE* gene (rs1539243 and rs17433930, LD $r^2=0.34$) displayed *P*-values <0.05. Conditional regression analysis did not provide evidence for more than one allele contributing to risk for SLE in genes with multiple associated SNPs (Supplementary Table S5), nor was there any evidence for epistatic interactions between SNPs at different loci.

We sought independent replication of our results by using data from a GWAS on SLE in European Americans.¹⁴ The SNPs, rs1539243 and rs17433930, in the *IKBKE* gene showed signals of association with SLE also in the US data ($P=0.0028$ and $P=0.0021$, respectively) (Table 2 and Supplementary Table S6). The SNP rs1539243 had been directly genotyped in the US cohort, and for rs17433930 imputed genotypes were analysed. Combined analysis of the data from the

Swedish and US cohorts revealed convincing association signals with SLE for these SNPs ($P_{\text{meta}}=0.00026$, OR=1.19, and $P_{\text{meta}}=0.00010$, OR=1.33, for rs1539243 and rs17433930, respectively), a result which was significant also after Bonferroni correction ($P_{\text{meta_corr}} < 0.01$). The two *IKBKE* SNPs rs17433930 and rs1539243, which we found to be associated with SLE are located in the tenth intron and fourth exon of the gene, respectively, where rs1539243 is a synonymous SNP in amino-acid residue 67 (Ile) of the IKK ϵ kinase.

Although the SNP rs4694178, located 3.3 kb downstream of the *IL8* gene, had only a trend-wise significant *P*-value in the US cohort compared with the more convincing association with SLE in the Swedish cohort ($P_{\text{US}}=0.064$, $P_{\text{SWE}}=5 \times 10^{-5}$), its association signal remained significant in the combined analysis after Bonferroni correction ($P_{\text{meta}}=0.00040$, OR=1.17, $P_{\text{meta_corr}} < 0.01$) (Table 2). This SNP was imputed in the US data (confidence=0.85), however, the directly genotyped SNP rs9999446, which is strongly correlated with the SNP rs4694178 ($r^2=0.86$) yielded a similar result ($P=0.085$) for association with SLE. For the SNP rs10199181 in the *STAT1* gene, we observed combined *P*-values $< 1 \times 10^{-3}$. The *STAT1* gene is located close to the *STAT4* gene in a region of high LD on chromosome 2q32.2. *STAT4* contains two linked SNPs, rs10181656 and rs7582694, which are strongly associated with SLE,¹⁸ and

Table 2 Meta-analysis of association with SLE for the Swedish and US cohorts

Gene	SNP	P ^d	Sweden ^a		US ^b		Risk allele	P	Meta-analysis ^c	
			OR (95% CI)	P ^d	OR (95% CI) ^e	Corrected P ^f			OR (95% CI) ^e	
<i>IKBKE</i>	rs1539243	0.031	1.19 (1.02–1.39)	0.0028	1.20 (1.07–1.35)	C	0.00026	0.0054	1.19 (1.09–1.31)	
<i>IKBKE</i>	rs17433930	0.013	1.33 (1.06–1.67)	0.0021	1.33 (1.11–1.58)	A	0.00010	0.0022	1.33 (1.16–1.53)	
<i>TANK</i>	rs3754974	0.0087	1.44 (1.09–1.89)	0.98	0.99 (0.80–1.22)	NA	0.22	1.0	1.13 (0.96–1.34)	
<i>TANK</i>	rs1267075	0.75	1.02 (0.88–1.19)	0.87	0.99 (0.89–1.11)	NA	0.99	1.0	1.00 (0.92–1.10)	
<i>STAT1</i>	rs2030171	1.2E–05	1.31 (1.16–1.49)	0.018	1.11 (1.02–1.21)	A	3.3E–05	0.00069	1.17 (1.09–1.26)	
<i>STAT1</i>	rs16833172	0.015	1.51 (1.09–2.09)	0.16	1.23 (0.95–1.59)	A	0.016	0.34	1.33 (1.09–1.63)	
<i>IL8</i>	rs4694178	5.2E–05	1.26 (1.12–1.42)	0.064	1.13 (1.04–1.23)	C	0.00040	0.0084	1.17 (1.10–1.26)	
<i>NRP1</i>	rs734187	0.67	1.03 (0.90–1.18)	0.66	1.01 (0.92–1.12)	A	0.56	1.0	1.02 (0.94–1.10)	
<i>NRP1</i>	rs1331314	0.32	1.11 (0.92–1.34)	0.11	1.11 (0.97–1.26)	G	0.061	1.0	1.11 (1.00–1.23)	
<i>NRP1</i>	rs2073320	0.42	1.05 (0.93–1.18)	0.61	0.98 (0.90–1.06)	NA	0.94	1.0	1.00 (0.94–1.07)	
<i>TRAF6</i>	rs5030482	0.0094	1.25 (1.05–1.49)	0.12	1.10 (0.97–1.24)	T	0.0097	0.20	1.14 (1.03–1.26)	
<i>TRAF6</i>	rs5030472	0.013	1.28 (1.05–1.55)	0.19	1.09 (0.96–1.25)	C	0.020	0.42	1.15 (1.03–1.28)	
<i>STAT5B</i>	rs6503691	0.44	1.08 (0.88–1.32)	0.78	0.98 (0.86–1.12)	NA	0.91	1.0	1.01 (0.90–1.13)	
<i>PIAS4</i>	rs2289863	0.19	1.09 (0.96–1.25)	0.72	0.98 (0.89–1.08)	NA	0.76	1.0	1.02 (0.95–1.10)	

Abbreviations: CI, confidence interval; MAF, minor Allele Frequency; NA, not Available; OR, odds ratio; SNP, single-nucleotide polymorphism.

^a826 Swedish cases and 1835 controls genotyped in the discovery and replication phases including controls genotyped on the 1M chip.

^b1310 US SLE cases and 7859 controls genotyped in a GWAS. Genotypes for rs17433930, rs3754974, rs2030171, rs16833172, rs4694178, rs734187, rs1331314 and rs5030482 were imputed in this dataset as described in Gateva *et al*.

^cMeta-analysis including 2136 SLE cases and 9694 controls.

^dP-values calculated using SNPtest.

^eOdds ratios are calculated relative to the risk allele defined in the analysis of the combined Swedish cohort.

^fP-value corrected for the 21 tested SNPs.

P-values <0.05 are indicated in bold.

conditional regression analysis of the data from the combined Swedish cohort indicates that the association signals from the *STAT1* and *STAT4* SNPs are not independent of each other (data not shown, remaining $P_{\text{conditional}} > 0.3$).

DISCUSSION

Our association study of genes from the type I IFN pathway and additional candidate genes for SLE highlighted two genes, *IKBKE* and *IL8*, as potential risk factors for SLE. In addition, the genes *TRAF6* and *TANK* showed significant association with SLE in the Swedish cohorts. Furthermore, *IRF5*, *TYK2*, *STAT4*, *IFIH1*, *IRAK1*, *IRF8* and the *PHRF1/IRF7* region have been reported by us elsewhere to be associated with SLE.^{1,4,18} Thus, polymorphisms in multiple genes connected to the type I IFN signalling system are important for SLE disease susceptibility.

IKBKE, the inhibitor of nuclear factor kappa-B kinase subunit epsilon gene, encodes IKK ϵ , a kinase that together with the TANK-binding kinase (TBK1) has a role in the innate antiviral response. IKK ϵ and TBK1 are activated when two intracellular RLR helicases, encoded by the *IFIH1* and *DDX58* genes, recognise viral RNA in virus-infected cells (Figure 2). These kinases are also activated upon stimulation of endosomal TLR3 by double stranded DNA, or cell membrane TLR4 by bacterial lipopolysaccharide (LPS). Together with TBK1, IKK ϵ mediates phosphorylation of the transcription factors IRF3 and IRF7, which leads to their activation and subsequent transcription of type I IFN and other inflammatory cytokines, but also activation of NF κ B has been reported.^{19,20} The activation of IRF3 and IRF7 can be inhibited by an ubiquitin-editing enzyme (A20), encoded by the *TNFAIP3* gene. As also variation in *TNFAIP3* and *IFIH1* are associated with SLE,^{4,21,22} this further supports an important role for the RLR pathway in the disease process. There is also evidence that IKK ϵ can phosphorylate STAT1, and thus contribute to the type I IFN signalling through the IFNAR.²³

The *IKBKE* gene has recently been implicated in risk for rheumatoid arthritis (RA).²⁴ The two polymorphisms with the most significant association with RA were tested in the discovery phase of our

study (rs2151222 and rs3748022 with $P=0.084$ and $P=0.49$ for association to SLE, respectively), however, these variants appear to be independent from the *IKBKE* variants rs1539243 and rs17433930 that we identified as risk alleles for SLE ($r^2 < 0.1$ discovery phase). The conjecture that *IKBKE* has a role in arthritis is supported by data from an animal model. IKK ϵ knockout mice have been shown to be less sensitive to induction of arthritis and exhibit less joint destruction than control mice.²⁵ In a published GWAS on women with SLE²⁶ the *IKBKE* SNP rs1539243 was tested, but no association was observed (dbGaP, <http://www.ncbi.nlm.nih.gov/gap>). The power of that study was, however, only 23% at the $P=0.05$ level to detect the association we observe.

The chemokine IL-8 has a wide range of pro-inflammatory effects, and its production can be triggered by immune complexes that also have the capacity to induce type I IFN production.²⁷ Recent data also suggests that IL-8 production in virus-infected cells is IFN dependent.²⁸ SLE patients with renal²⁹ or CNS involvement³⁰ have elevated IL-8 levels in their serum and cerebrospinal fluid, respectively, and serum IL-8 levels and disease activity correlate in SLE patients.³¹ As SLE disease flares are associated with increased IFN- α production,⁶ these observations provide a link between the SLE disease process, IL-8 and the type I IFN system.

Although we observed a significant association of *IL8* with SLE in this study, there are previous conflicting reports on the association between *IL8* and SLE,^{26,32–34} only two of which, a Spanish case-control study and a GWAS in women with SLE, having similar power to our discovery phase. The power of these studies was around 50% at the 0.05 significance level to detect the effect for *IL8* that we observe. The LD between the SNP rs2227306 tested in the Spanish study and the SNP rs4694178 in our study is very high ($r^2=0.97$ in the HapMap CEU population), but they do not observe any association with SLE for this SNP.³⁴ In our study, the association signal is mainly contributed by the Swedish cohort, which has a higher frequency of the risk allele than both the US and Spanish cohorts (control frequency of rs4694178 C: Sweden=0.46, USA=0.41 and the linked allele rs2227306 T: Spain=0.41). In the women GWAS,²⁶ a SNP




Figure 2 A schematic illustration of signalling within the type I interferon (IFN) system. Circled gene names have confirmed association to SLE: *STAT4*, *IRF5*, *IRF7*, *IRF8*, *TNFAIP3* (A20), *TNFSF4* (OX40L), *FCGR2A*, *TYK2*, *IFIH1* (MDA5), *IRAK1*, as well as *IKBKE* and *IL8*, identified in this study. Genes with dashed circles are *TANK* and *TRAF6*, which we find associated with SLE in the Swedish but not US cohorts, and *STAT1* that has an association signal, which cannot be distinguished from that of *STAT4*. *Left*: Induction of inflammatory cytokines and type I IFN genes in response to viral RNA/DNA by the helicases IFIH1 (MDA5), DDX58 (RIG-I) and the DNA sensor ZBP1 (DAI), or by bacterial LPS through toll-like receptor 4. These pathways signal using TANK that interacts with IKBKE or TBK1, which together can mediate phosphorylation of the interferon regulatory factors IRF3 and 7, leading to the transcription of type I IFN genes. Activation of this pathway can be inhibited by TNFAIP3 (A20). *Middle*: Induction of IFN production in plasmacytoid dendritic cells (pDCs) by endogenous immune complexes. Immune complexes are endocytosed through FCGRIIA, which leads to activation of TLR7/9. After a signalling cascade, which includes IRAK1 and TRAF6, the interferon regulatory factors IRF5 and IRF7 are activated leading to the transcription of type I IFN genes. Also IRF8 can contribute to this activation. *Right*: IFN signalling through the type I IFN receptor IFNAR. Association of type I IFN to the receptor activates the kinases TYK2 and JAK1, which signal through STAT1 and STAT2. Also other STATs, including STAT4, are activated in this process. The STAT1/STAT2 complex associates with IRF9, which in turn binds to interferon-stimulated response elements (ISREs) and induces expression of IFN-induced genes, such as IRF5, IRF7 and indirectly IL8. *Far right*: Membrane proteins expressed on pDCs. TNFSF4 (OX40L), together with endogenous type I IFNs provide a means for the pDCs to control the T-cell response, specifically to profile it towards a T_H1 response. Activation of the NFκB pathway, and thus production of inflammatory cytokines, also occurs through TLR7/9 (not shown). Proteins are denoted by their corresponding gene names.

perfectly correlated with *IL8* rs4694178 (rs4694636, $r^2=1$ HapMap CEU) was tested, and showed an odds ratio suggestive of association with SLE (OR CI: 1.01–1.49, dbGaP). The modest power to detect an effect of this size, or genetic heterogeneity between populations could explain these conflicting results.

Although the *TANK* and *TRAF6* genes, which gave clear association signals ($P < 0.001$) in the Swedish cohort, did not replicate in the US GWAS data, these genes remain interesting candidates. *TANK* signals immediately upstream of IKKε in the TLR4- and IFIH1/DDX58-mediated activation of type I IFNs and inflammatory cytokines in response to bacterial and viral stimuli, respectively (Figure 2). Association of *TANK* with SLE would thus further support an important role for the RLR pathway in SLE. *TRAF6* is a ubiquitin ligase that mediates signal transduction from, for example, members of the TLR family leading to activation of NFκB and IRFs. Polyubiquitination of IRF5 after its interaction with IRAK1 is mediated by *TRAF6*, which enables IRF5 to translocate to the nucleus and exert its effect on gene expression. On the other hand, *TNFAIP3* can inhibit TLR-induced activity of NFκB by de-ubiquitination of

TRAF6.³⁵ Interestingly, also *IRAK1* has been associated with risk for SLE,³⁶ and because SNPs in the *TRAF6* region and in *TNFAIP3* have also been associated with risk for RA,^{37–39} it seems as variants of all these genes have the potential to contribute to loss of tolerance and autoimmune reactions. Further studies will be needed to determine whether the *TANK* and *TRAF6* genes have an effect on SLE, and whether this effect is specific for Scandinavian populations.

Our study confirms the important role of the type I IFN system in SLE, and suggests multiple genes from this pathway as candidates for functional studies and as interesting therapeutic targets (Figure 2). These results also point more specifically to the importance of genes in the RLR pathway, which is activated in response to viral infections because of the ability of IFIH1/DDX58 to recognise cytoplasmic viral RNA. This pathway is active in cells other than the pDCs, including monocyte derived dendritic cells. In addition to IFIH1, factors such as *TANK*, *IKBKE* and *TNFAIP3* contribute to signalling in the RLR pathway (reviewed in⁴⁰). However, there is also evidence for involvement in SLE of the MYD88-dependent pathway activated by endosomal TLR7/9 by RNA/DNA from dying cells, immune complexes

(IC) or by viral RNA/DNA, because *IRF8*, *IRAK1*, *FCGR2A* and potentially *TRAF6*, in this pathway are associated with SLE. Thus, at least two pathways seem important in SLE, both leading to production of type I IFN and inflammatory cytokines through activation of *IRF3*, *IRF5* and *IRF7*, and additional transcription factors, especially *NFκB*. Association to SLE has further been demonstrated for the type I IFN signalling through the *IFNAR*, specifically for *TYK2* and *STAT4*, as well as the IFN-regulated genes *IFIH1*, *IRF5*, *IRF7* and *IL8*. Consequently, a large number of genes, located in different functional parts of the type I IFN system, are associated with SLE indicating a general role for the type I IFN system genes in autoimmunity.

CONFLICT OF INTEREST

Robert R Graham and Timothy W Behrens are employees of Genentech Corp. The other authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This study was supported by a Target Identification in Lupus (TIL) grant from the Alliance for Lupus Research, US, the Swedish Research Council for Medicine, the Knut and Alice Wallenberg Foundation, the Swedish Rheumatism Foundation, the King Gustaf V 80-year Foundation, COMBINE, the European Community 6th Framework Program (LSHM-CT-2007-037273), the Swedish Heart-Lung Foundation, the Torsten and Ragnar Söderberg Foundation, the 'Visare Norr' Fund for Northern County Councils of Sweden, The Åke Wiberg Foundation, ALF funding from Stockholm County Council and Karolinska Institutet, the National Institutes of Health (R01 AR44804, K24 AR02175), the National Center for Research Resources (5 M01 RR00079), and a Kirkland Scholar Award to LAC. We thank Rezvan Kiani Dehkordi at the Rheumatology Clinic, Uppsala University Hospital, for assistance with DNA sample collection. Professor Göran Hallmans, Head of the Medical Biobank of Northern Sweden for providing blood samples. Ann-Christin Wiman, Caisa Pöntinen, Molecular Medicine, Department of Medical Sciences, Uppsala University, for assistance with genotyping. Genotyping was performed using equipment available at the SNP Technology Platform in Uppsala, Sweden (<http://www.genotyping.se>).

- Sigurðsson S, Nordmark G, Goring HH *et al*: Polymorphisms in the tyrosine kinase 2 and interferon regulatory factor 5 genes are associated with systemic lupus erythematosus. *Am J Hum Genet* 2005; **76**: 528–537.
- Remmers EF, Plenge RM, Lee AT *et al*: *STAT4* and the risk of rheumatoid arthritis and systemic lupus erythematosus. *N Engl J Med* 2007; **357**: 977–986.
- Graham RR, Hom G, Ortmann W, Behrens TW: Review of recent genome-wide association scans in lupus. *J Intern Med* 2009; **265**: 680–688.
- Gateva V, Sandling JK, Hom G *et al*: A large-scale replication study identifies *TNIP1*, *PRDM1*, *JAZF1*, *UHRF1BP1* and *IL10* as risk loci for systemic lupus erythematosus. *Nat Genet* 2009; **41**: 1228–1233.
- Hooks JJ, Moutsopoulos HM, Geis SA, Stahl NI, Decker JL, Notkins AL: Immune interferon in the circulation of patients with autoimmune disease. *N Engl J Med* 1979; **301**: 5–8.
- Bengtsson AA, Sturfelt G, Truedsson L *et al*: Activation of type I interferon system in systemic lupus erythematosus correlates with disease activity but not with antiretroviral antibodies. *Lupus* 2000; **9**: 664–671.
- Baechler EC, Batliwalla FM, Karypis G *et al*: Interferon-inducible gene expression signature in peripheral blood cells of patients with severe lupus. *Proc Natl Acad Sci USA* 2003; **100**: 2610–2615.
- Ronnblom LE, Alm GV, Oberg KE: Possible induction of systemic lupus erythematosus by interferon-alpha treatment in a patient with a malignant carcinoid tumour. *J Intern Med* 1990; **227**: 207–210.
- Wallace DJ, Petri M, Olsen N *et al*: MEDI-545, an anti-interferon alpha monoclonal antibody, shows evidence of clinical activity in systemic lupus erythematosus. *Arthritis Rheum* 2007; **56**(Suppl 9): S526–S527.
- Yao Y, Richman L, Higgs BW *et al*: Neutralization of interferon-alpha/beta-inducible genes and downstream effect in a phase I trial of an anti-interferon-alpha monoclonal antibody in systemic lupus erythematosus. *Arthritis Rheum* 2009; **60**: 1785–1796.
- Tan EM, Cohen AS, Fries JF *et al*: The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982; **25**: 1271–1277.
- Bell PA, Chaturvedi S, Gelfand CA *et al*: SNPstream UHT: ultra-high throughput SNP genotyping for pharmacogenomics and drug discovery. *Biotechniques* 2002; (Suppl 74), 70–72, 76–77.

- Broadbent HM, Peden JF, Lorkowski S *et al*: Susceptibility to coronary artery disease and diabetes is encoded by distinct, tightly linked SNPs in the ANRIL locus on chromosome 9p. *Hum Mol Genet* 2008; **17**: 806–814.
- Hom G, Graham RR, Modrek B *et al*: Association of systemic lupus erythematosus with C8orf13-BLK and ITGAM-ITGAX. *N Engl J Med* 2008; **358**: 900–909.
- Marchini J, Howie B, Myers S, McVean G, Donnelly P: A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet* 2007; **39**: 906–913.
- Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D: Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* 2006; **38**: 904–909.
- Purcell S, Neale B, Todd-Brown K *et al*: PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007; **81**: 559–575.
- Sigurðsson S, Nordmark G, Garnier S *et al*: A risk haplotype of *STAT4* for systemic lupus erythematosus is over-expressed, correlates with anti-dsDNA and shows additive effects with two risk alleles of *IRF5*. *Hum Mol Genet* 2008; **17**: 2868–2876.
- Sharma S, tenOever BR, Grandvaux N, Zhou GP, Lin R, Hiscott J: Triggering the interferon antiviral response through an IKK-related pathway. *Science* 2003; **300**: 1148–1151.
- Sankar S, Chan H, Romanow WJ, Li J, Bates RJ: IKK- α signals through *IRF3* and *NFκB* to mediate the production of inflammatory cytokines. *Cell Signal* 2006; **18**: 982–993.
- Graham RR, Cotsapas C, Davies L *et al*: Genetic variants near *TNFAIP3* on 6q23 are associated with systemic lupus erythematosus. *Nat Genet* 2008; **40**: 1059–1061.
- Musone SL, Taylor KE, Lu TT *et al*: Multiple polymorphisms in the *TNFAIP3* region are independently associated with systemic lupus erythematosus. *Nat Genet* 2008; **40**: 1062–1064.
- tenOever BR, Ng SL, Chua MA, McWhirter SM, Garcia-Sastre A, Maniatis T: Multiple functions of the IKK-related kinase IKKepsilon in interferon-mediated antiviral immunity. *Science* 2007; **315**: 1274–1278.
- Dieguez-Gonzalez R, Akar S, Calaza M *et al*: Genetic variation in the nuclear factor kappaB pathway in relation to susceptibility to rheumatoid arthritis. *Ann Rheum Dis* 2009; **68**: 579–583.
- Corr M, Boyle DL, Ronacher L, Flores N, Firestein GS: Synergistic benefit in inflammatory arthritis by targeting I kappaB kinase epsilon and interferon beta. *Ann Rheum Dis* 2009; **68**: 257–263.
- Harley JB, Alarcon-Riquelme ME, Criswell LA *et al*: Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in *ITGAM*, *PXK*, *KIAA1542* and other loci. *Nat Genet* 2008; **40**: 204–210.
- Santer DM, Yoshio T, Minota S, Moller T, Elkon KB: Potent induction of IFN- α and chemokines by autoantibodies in the cerebrospinal fluid of patients with neuropsychiatric lupus. *J Immunol* 2009; **182**: 1192–1201.
- Manuse MJ, Parks GD: TLR3-dependent upregulation of RIG-I leads to enhanced cytokine production from cells infected with the parainfluenza virus SV5. *Virology* 2010; **397**: 231–241.
- Holcombe RF, Baethge BA, Wolf RE *et al*: Correlation of serum interleukin-8 and cell surface lysosome-associated membrane protein expression with clinical disease activity in systemic lupus erythematosus. *Lupus* 1994; **3**: 97–102.
- Trysberg E, Carlsten H, Tarkowski A: Intrathecal cytokines in systemic lupus erythematosus with central nervous system involvement. *Lupus* 2000; **9**: 498–503.
- Lit LC, Wong CK, Tam LS, Li EK, Lam CW: Raised plasma concentration and *ex vivo* production of inflammatory chemokines in patients with systemic lupus erythematosus. *Ann Rheum Dis* 2006; **65**: 209–215.
- Rovin BH, Lu L, Zhang X: A novel interleukin-8 polymorphism is associated with severe systemic lupus erythematosus nephritis. *Kidney Int* 2002; **62**: 261–265.
- Huang CM, Huo AP, Tsai CH, Chen CL, Tsai FJ: Lack of association of interleukin-6 and interleukin-8 gene polymorphisms in Chinese patients with systemic lupus erythematosus. *J Clin Lab Anal* 2006; **20**: 255–259.
- Sanchez E, Sabio JM, Callejas JL *et al*: Association study of genetic variants of pro-inflammatory chemokine and cytokine genes in systemic lupus erythematosus. *BMC Med Genet* 2006; **7**: 48.
- Boone DL, Turer EE, Lee EG *et al*: The ubiquitin-modifying enzyme A20 is required for termination of Toll-like receptor responses. *Nat Immunol* 2004; **5**: 1052–1060.
- Jacob CO, Zhu J, Armstrong DL *et al*: Identification of *IRAK1* as a risk gene with critical role in the pathogenesis of systemic lupus erythematosus. *Proc Natl Acad Sci USA* 2009; **106**: 6256–6261.
- Raychaudhuri S, Thomson BP, Remmers EF *et al*: Genetic variants at *CD28*, *PRDM1* and *CD2/CD58* are associated with rheumatoid arthritis risk. *Nat Genet* 2009; **41**: 1313–1318.
- Thomson W, Barton A, Ke X *et al*: Rheumatoid arthritis association at 6q23. *Nat Genet* 2007; **39**: 1431–1433.
- Plenge RM, Cotsapas C, Davies L *et al*: Two independent alleles at 6q23 associated with risk of rheumatoid arthritis. *Nat Genet* 2007; **39**: 1477–1482.
- Takeuchi O, Akira S: MDA5/RIG-I and virus recognition. *Curr Opin Immunol* 2008; **20**: 17–22.



This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported Licence. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)