

UC Berkeley

UC Berkeley Previously Published Works

Title

Early-life exposure to p,p'-DDT and p,p'-DDE in South African children participating in the VHEMBE study: An assessment using repeated serum measurements and pharmacokinetic modeling

Permalink

<https://escholarship.org/uc/item/9hf4375c>

Authors

Verner, Marc-André
Chevrier, Jonathan
Ngueta, Gérard
et al.

Publication Date

2018-10-01

DOI

10.1016/j.envint.2018.07.010

Peer reviewed



Published in final edited form as:

Environ Int. 2018 October ; 119: 478–484. doi:10.1016/j.envint.2018.07.010.

Early-life exposure to p,p' -DDT and p,p' -DDE in South African children participating in the VHEMBE study: An assessment using repeated serum measurements and pharmacokinetic modeling

Marc-André Verner^{a,b,*}, Jonathan Chevrier^c, Gérard Ngueta^{a,b}, Stephen Rauch^d, Riana Bornman^{e,f}, and Brenda Eskenazi^d

^aUniversité de Montréal Public Health Research Institute (IRSPUM), Montreal, QC, Canada

^bDepartment of Occupational and Environmental Health, School of Public Health, Université de Montréal, Montreal, QC, Canada

^cDepartment of Epidemiology, Biostatistics and Occupational Health, Faculty of Medicine, McGill University, Montreal, QC, Canada

^dCenter for Environmental Research and Children's Health (CERCH), School of Public Health, University of California, Berkeley, CA, USA

^eSchool of Health Systems and Public Health, University of Pretoria, Pretoria, South Africa

^fUniversity of Pretoria Institute for Sustainable Malaria Control, University of Pretoria, Pretoria, South Africa

Abstract

Background: The World Health Organization recommends indoor residual spraying (IRS) of insecticides (including dichlorodiphenyltrichloroethane [DDT]) to fight malaria vectors in endemic countries. There is limited information on children's exposure to DDT in sprayed areas, and tools to estimate early-life exposure have not been thoroughly evaluated in this context.

Objectives: To document serum p,p' -DDT/E levels in 47 mothers and children participating in the Venda Health Examination of Mothers, Babies and their Environment (VHEMBE), a study conducted in an area where IRS insecticides are used annually, and to evaluate the precision and accuracy of a published pharmacokinetic model for the estimation of children's p,p' -DDT/E levels.

Methods: p,p' -DDT/E levels were measured in maternal serum at delivery, and in children's serum at 12 and 24 months of age. A pharmacokinetic model of gestational and lactational exposure was used to estimate children's p,p' -DDT/E levels during pregnancy and the first two years of life, and estimated levels were compared to measured levels.

*Corresponding author at: Department of Occupational and Environmental Health, School of Public Health, Université de Montréal, 2375 ch. de la Côte-Sainte-Catherine, Suite 4105, Montreal, QC H3T 1A8, Canada. marc-andre.verner.1@umontreal.ca (M.-A. Verner).

Results: The geometric means of children's serum p,p' -DDT/E levels at 12 and 24 months were higher than those of maternal serum levels. Regression models of measured children's p,p' -DDT/E levels vs. levels estimated with the pharmacokinetic model (which only accounted for children's exposure through placental transfer and breastfeeding) had coefficients of determination (R^2) ranging from 0.75 to 0.82. Estimated p,p' -DDE levels were not significantly different from measured levels, whereas p,p' -DDT levels were overestimated by 36% at 12 months, and 51% at 24 months.

Conclusion: Results indicate that children living in a sprayed area have serum p,p' -DDT/E levels exceeding their mothers' during the first two years of life. The pharmacokinetic model may be useful to estimate children's levels in the VHEMBE population.

1. Introduction

The World Health Organization (WHO) recommends indoor residual spraying (IRS) to control malaria vectors in endemic countries and includes dichlorodiphenyltrichloroethane (DDT) in their list of recommended insecticides (WHOPES, 2015). In many countries, including South Africa, DDT has been used for decades (Mabaso et al., 2004), resulting in elevated body burdens of DDT and its breakdown product dichlorodiphenyldichloroethylene (DDE) in individuals living in sprayed homes (Gaspar et al., 2017). Concerns have been raised regarding the potential health effects related to exposure to these chemicals, especially in developing children (Bouwman and Kylin, 2009; Chevrier et al., in press; Coker et al., 2018; Eskenazi et al., 2009; Huang et al., in press).

Because of their biological half-lives in the order of years (Ritter et al., 2009; Smith, 1999; Wolff et al., 2000), p,p' -DDT/E accumulate in human tissues. p,p' -DDT/E have been shown to transfer to children during pregnancy (Aylward et al., 2014) and breastfeeding (LaKind et al., 2009), resulting in exposures during critical windows of development. Multiple studies have provided evidence of the substantial impact of breastfeeding on children's p,p' -DDE levels. Whereas neonates' serum p,p' -DDT/E levels are similar or slightly lower than maternal levels at birth on a lipid basis (Aylward et al., 2014), studies with longitudinal measurements have shown that children's serum levels can increase substantially during breastfeeding (Lackmann, 2006). Studies have reported that breastfed children have higher p,p' -DDE levels than children who had been fed formula, from as early as 6 weeks of age (Lackmann et al., 2004) to as late as 14 years (Gascon et al., 2015). Breastfeeding duration has also been associated with higher children p,p' -DDE levels during the first years of life (Caspersen et al., 2016; Ribas-Fito et al., 2005). It has been estimated that the dose of lipophilic persistent organic pollutants like p,p' -DDT/E received by the child through breastfeeding can be substantially higher than the maternal daily dose (Haddad et al., 2015), primarily due to bioaccumulation in mothers and partitioning into breast milk lipids.

The scientific literature on children's exposure to p,p' -DDT/E in sprayed areas is scarce. In addition to placental transfer and breast-feeding, children living in houses sprayed with p,p' -DDT may be exposed through direct contact with contaminated dust and consumption of local chicken and eggs. In a previous study of undisturbed dust from houses in the Limpopo province of South Africa, the majority of dust samples from sprayed houses (82%) had

quantifiable levels of p,p' -DDT (Gaspar et al., 2016). To our knowledge, no study has obtained repeated p,p' -DDT/E levels during early childhood in the context of IRS. While tools have been developed to estimate children's early-life exposure to p,p' -DDT/E (Gyalpo et al., 2012; Verner et al., 2009; Verner et al., 2013), their ability to estimate children's body burden has not been thoroughly evaluated in this context.

In this study, we aimed to i) document serum p,p' -DDT/E levels in a subset of mothers and children ($n = 47$) from the Limpopo province participating in the Venda Health Examination of Mothers, Babies and their Environment (VHEMBE); and to ii) evaluate the precision and accuracy of a published pharmacokinetic model for the estimation of children's p,p' -DDT/E levels during the first two years of life.

2. Methods

2.1. Study participants

The Venda Health Examination of Mothers, Babies and their Environment (VHEMBE) study population is detailed in Eskenazi et al. (2018). Briefly, women who presented at Tshilidzini Hospital (Thohoyandou, Limpopo, South Africa) for delivery between August 2012 and December 2013 were eligible if they were ≥ 18 years old, spoke Tshivenda at home, lived within 20 km of the hospital, planned to remain in the area, had not been diagnosed with malaria during pregnancy, had contractions > 5 min apart, and gave birth to a viable singleton. A total of 920 out of the 1649 women who were approached were eligible. Out of the eligible women, 152 refused to participate, 14 did not have a completed baseline questionnaire, and blood volume was insufficient for p,p' -DDT/E analysis in 3 women, resulting in a total of 751 participants. Blood was sampled prior to delivery ($n = 595$) or immediately after delivery ($n = 156$) (range: 31 days prior to delivery to 2 days after delivery). Breastfeeding mothers provided breast milk samples by expressing directly in sterile collection bottles during a home visit one week postpartum ($n = 706$) and at our field office visits when children were aged 12 months ($n = 532$). In addition, study nurses collected venous blood from children via venipuncture during visits to the field office at 12 months ($n = 642$) and 24 months ($n = 662$). Blood and breast milk samples were processed immediately after collection and stored at -80°C until shipment to analytical laboratories on dry ice. Due to budgetary restrictions, children's serum p,p' -DDT/E levels could only be measured in a subset of the cohort, and breast milk p,p' -DDT/E levels could only be measured at one collection time; we selected the 12-month samples because milk is fully mature at this point (versus transition milk at one week postpartum), and it could potentially reflect maternal exposure after delivery (whereas early breast milk levels likely reflect exposure prior to delivery). Serum and breast milk samples were selected from a subset of 47 mother-child dyads based on i) availability of p,p' -DDT or p,p' -DDE levels in maternal serum (previous analyses), ii) sufficient children's serum volume for p,p' -DDT/E analyses, and iii) availability of a breast milk sample at 12 months postpartum for p,p' -DDT/E analyses. All mothers or legal guardian provided informed consent prior to data and sample collection. Protocols were reviewed and approved by institutional review boards at the Université de Montréal, University of California, Berkeley, McGill University, the

University of Pretoria, the Limpopo Department of Health and Social Development, and Tshilidzini Hospital.

2.2. Maternal and child characteristics

Questionnaires were administered before hospital discharge, at the one-week home visit, and during the 12-month and 24-month visits at the field office. Collected information included demographics (e.g., child sex, age at delivery, duration of breastfeeding) and IRS during pregnancy and the period preceding each time point. The total duration of breastfeeding was approximated based on the question “How many months old was [child’s name] when you stopped breastfeeding completely?” The duration of exclusive breastfeeding was approximated based on the question “How many months did you exclusively breast-feed, that is, for how many months did [child’s name] only receive breast milk without any additional food or water?” Mothers were weighed at study enrollment and at the 12-month and 24-month visits using a Tanita HD-351 scale (Tokyo, Japan). Birth weight was measured using a Tanita BD-815U neonatal scale (Tokyo, Japan) in the delivery suite shortly after delivery, and children were weighed at 12-month and 24-month visits using a Tanita BD-590 pediatric scale (Tokyo, Japan). All weight measurements were taken in triplicate and averaged.

2.3. Measurement of serum p,p' -DDT/E levels

Maternal serum p,p' -DDT/E levels were measured at the Environmental Health Laboratory at Emory University’s Rollins School of Public Health using gas chromatography-tandem mass spectrometry (GC–MS) with isotope dilution quantification (Barr et al., 2003). Limits of detection were 0.01 µg/L for p,p' -DDT and 0.03 µg/L for p,p' -DDE. Quality control samples (i.e., sealed blanks, field blanks, and spiked samples) were also analyzed. Triglycerides and total cholesterol were measured using standard enzymatic methods (Roche Chemicals, Indianapolis, USA). Breast milk and children’s serum concentrations of p,p' -DDT/E were measured by the Centre de Toxicologie du Québec (CTQ) of the Institut national de santé publique du Québec (INSPQ) by gas chromatography mass spectrometry (GC–MS) as described in Fisher et al. (2016). The limits of detection for both compounds were between 0.01 and 0.05 µg/L for child serum and 0.2 µg/L for breast milk. Lipids in children’s serum samples levels were measured by enzymatic methods combined with colorimetry at the laboratory of Centre Hospitalier de l’Université Laval (Quebec, Canada). Breast milk lipid content was assessed by extraction, evaporation and gravimetric measurement. Maternal and child serum p,p' -DDT/E levels were expressed on a total lipid basis estimated based on triglyceride and total cholesterol levels (Phillips et al., 1989). Breast milk levels were also expressed on a lipid basis based on total lipids.

2.4. Pharmacokinetic modeling

We used a published and validated pharmacokinetic model to estimate children’s p,p' -DDT/E levels. The model is presented in detail in Verner et al. (2013). Briefly, the model included two compartments representing the maternal and child lipids in which p,p' -DDT/E are assumed to distribute primarily given their high lipophilicity. The compartments were connected through placental transfer and breast-feeding. During pregnancy, maternal and fetal concentrations were assumed to be equal on a lipid basis. Transfer through

breastfeeding was based on breast milk consumption, breast milk lipid content and maternal p,p' -DDT/E levels, assuming p,p' -DDT/E levels in breast milk lipids were equal to levels in maternal lipids. Elimination in the mother and child was based on published biological half-lives. The pharmacokinetic model allows simulating maternal p,p' -DDT/E levels from maternal birth onwards, and child's levels during gestation, breast-feeding, and after breastfeeding has ceased.

To simulate p,p' -DDT/E levels for each participating mother-child dyad, the pharmacokinetic model incorporated individual-specific information on the mother (i.e., age at delivery, weight at admission for delivery, and weight at 12 months and 24 months postpartum), the child (i.e., sex, birth weight, and weight at 12 months and 24 months of age), the duration of exclusive and total breastfeeding, and the maternal serum p,p' -DDT/E levels. 16 out of the 47 mothers were only seen post-delivery; for these participants, body weight at delivery was estimated using a linear regression model of pre- and post-delivery body weight based on data from participants with both measurements (model $R^2 = 0.96$). Prepregnancy weight and gestational weight gain were not available. We assumed a 16 kg weight gain during pregnancy for all participants based on a previous study of South African women (Wrottesley et al., 2017). During non-exclusive breastfeeding, we assumed milk intake to be half the intake during exclusive breastfeeding. For each mother-child dyad and compound, the model was run iteratively to estimate a maternal daily dose (absorbed) yielding a simulated maternal level matching the measured level. Then, the model was run using the estimated maternal daily dose to generate a time-course of maternal, child and breast milk levels from maternal birth until two years after delivery. These simulations provided complete profiles of p,p' -DDT/E for each mother-child dyad (see example in Fig. 1).

Simulations were performed for dyads with maternal serum levels above the limit of detection for p,p' -DDT ($n = 46$) or p,p' -DDE ($n = 47$). We set out to perform a first set of simulations where children were exposed solely through placental transfer and breastfeeding, with the possibility to perform a second set of simulations including environmental exposure (e.g., dust ingestion) in the child if estimated levels were below measured levels in the first set of simulations.

2.5. Statistical analyses

To compare maternal serum p,p' -DDT/E levels at delivery to children's levels at 12 months and 24 months, we first calculated child/mother ratios for each mother-child dyad with levels above the limit of detection (one dyad with maternal serum p,p' -DDT level below the limit of detection was excluded from p,p' -DDT analyses), and subsequently calculated the geometric mean of child/mother ratios at 12 months and 24 months. We also computed Pearson correlation analyses between maternal, child and breast milk levels of p,p' -DDT/E (levels were \log_{10} -transformed prior to correlation analyses).

To evaluate the pharmacokinetic model precision and accuracy, we performed linear regression analyses of \log_{10} -transformed measured levels vs. \log_{10} -transformed simulated levels (for children's serum levels and breast milk levels). Precision was assessed by calculating coefficients of determination (R^2); a precise model would yield a R^2 close to 1.

Accuracy was assessed in terms of β coefficients and intercepts; an accurate model would yield a β coefficient with 95% confidence intervals including 1, and an intercept with a 95% confidence interval including 0. We also calculated mean relative errors to complement these measurements of model accuracy. Relative errors were calculated as follows (Eq. (1)):

$$\text{Relative error}_i = 100 \times \frac{\text{Estimated level}_i - \text{Measured level}_i}{\text{Measured level}_i} \quad (1)$$

3. Results

Children participating in this study were born to mothers who were between 18 and 39 years of age at delivery (Table 1). The geometric mean total duration of breastfeeding was 18.7 months, whereas the geometric mean duration of exclusive breastfeeding was < 2 months. One mother (2%) reported that her home was sprayed with DDT during pregnancy, six mothers (13%) reported that their homes were sprayed with DDT during the first year after birth, and six mothers (13%) reported that their homes were sprayed with DDT during the second year after birth. *p,p'*-DDT/E were detected in all serum and milk samples, except for one maternal serum *p,p'*-DDT level that was below the limit of detection. The geometric means of *p,p'*-DDT/E levels were higher in children's serum than maternal serum at 12 months and 24 months (Table 1). A high proportion (95%) of child/mother serum *p,p'*-DDT/E level ratios were above 1, indicating that children's *p,p'*-DDT/E levels at 12 months and 24 months were mostly higher than maternal levels at delivery. The geometric means of child/mother level ratios were higher for *p,p'*-DDE (3.6 at 12 months, and 4.1 at 24 months) than for *p,p'*-DDT (2.5 at 12 months, and 2.4 at 24 months) (Table 2). The maximum child/mother level ratios were 42 for *p,p'*-DDT and 32 for *p,p'*-DDE.

Maternal serum, children's serum and breast milk levels were highly correlated, with Pearson correlation coefficients ranging from 0.866 to 0.984 (Table 3). *p,p'*-DDT and *p,p'*-DDE levels were highly correlated within each matrix, with correlation coefficients ranging from 0.895 to 0.923.

Regression models of measured children's *p,p'*-DDT/E levels vs. levels estimated with the pharmacokinetic model had coefficients of determination (R^2) ranging between 0.75 and 0.82 at 12 months and 24 months (Fig. 2 and Table 4). R^2 s for children's levels at 24 months of age were only slightly lower than at 12 months of age. Linear regression models of measured vs. estimated children's *p,p'*-DDT/E levels had β coefficients ranging from 0.898 to 0.968, which were not statistically different from 1. The only intercept that was significantly > 0 was for *p,p'*-DDE levels at 24 months (Table 4). Mean relative errors were not significantly different from 0 for *p,p'*-DDE. However, they were significantly > 0 for *p,p'*-DDT levels, indicating that children's *p,p'*-DDT levels were overestimated by the pharmacokinetic model. Because estimated *p,p'*-DDT/E levels were either accurate or overestimated in simulations only accounting for placental transfer and breastfeeding, we did not include additional environmental exposure (e.g., dust ingestion, dermal contact, food consumption) in our model simulations. Relative errors were not significantly different in males and females (results not shown).

Regression models of measured breast milk p,p' -DDT and p,p' -DDE levels vs. levels estimated with the pharmacokinetic model had R^2 s of 0.78 and 0.83 at 12 months postpartum, respectively (Fig. 2 and Table 4). Although β values for regression models of measured vs. estimated breast milk levels were not significantly different from one, the intercept for p,p' -DDT was > 0 . Mean relative errors indicated that estimated breast milk levels were underestimated by ~30%.

The maximum children's concentrations estimated by the pharmacokinetic model had geometric means of 253 ng/g lipids (range: 16 to 6888) for p,p' -DDT and 952 ng/g lipids (range: 85 to 28,217) for p,p' -DDE. Estimated daily lactational exposure (ng/kg/day) at ~one month postpartum (time when the daily dose on a body weight basis is the highest) was compared to estimated maternal daily dose (ng/kg/day); the geometric mean lactational/maternal dose ratio was 77 for p,p' -DDT and p,p' -DDE. In other words, nursing infants were exposed to a daily dose that was 77 times greater than the maternal daily dose on a body weight basis.

4. Discussion

In this study of South African children's exposure to p,p' -DDT/E, we found that children's serum levels measured at 12 months and 24 months of age were substantially higher than their mothers' at delivery. These findings are consistent with previous reports of increasing serum p,p' -DDE levels in breastfed infants from six weeks to six months of life (Lackmann, 2006), a period during which all children included in our study were breastfed. In addition to lactational exposure, direct environmental exposure (e.g., food intake, dust/soil ingestion, dermal contact) could contribute to children's p,p' -DDT/E levels. However, simulations using the pharmacokinetic model, which only considered placental transfer and breastfeeding, accounted for most if not all of the measured children's p,p' -DDT/E levels at 12 months and 24 months of age (see mean relative errors in Table 4). Estimated daily doses through lactation (ng/kg/day) were 77 times higher than estimated maternal daily doses expressed in the same units. Similar conclusions with regards to the contribution of breast milk to children's body burden were drawn by Bouwman et al. (1992) based on children's blood and breast milk levels from the Kwazulu (now Kwazulu-Natal) area of South Africa.

R^2 s obtained for the regression models of measured vs. estimated p,p' -DDT levels (0.77 at 12 months, and 0.75 at 24 months [$n = 46$]) were higher than those obtained in a previous validation exercise in Inuit children (0.47 at 6 months [$n = 103$]) and Slovak children (0.58 at 6 months [$n = 216$], and 0.45 at 16 months [$n = 717$]). R^2 s obtained for p,p' -DDE levels (0.82 at 12 months, and 0.79 at 24 months [$n = 47$]) were similar to that in Inuits (0.83 at 6 months [$n = 156$]), but higher than those obtained in Slovaks (0.49 at 6 months [$n = 216$], and 0.44 at 16 months [$n = 728$]). Mean relative errors indicated that children's p,p' -DDT levels were overestimated by 36% at 12 months, and 51% at 24 months (Table 4). Similarly, children's p,p' -DDT levels were overestimated in the previous validation effort in Inuits (73% at 6 months) and Slovaks (127% at 6 months, and 68% at 16 months) (data not published). This overestimation may be due to certain pharmacokinetic parameters, including the assumption that 100% of p,p' -DDT lactational dose is absorbed by the child. On the other hand, β coefficients for linear regressions of measured vs. simulated p,p' -DDT

levels were not significantly different from 1, and intercepts were not significantly different from 0. Of note, confidence intervals around measures of accuracy may be inflated given the small sample size. Therefore, caution should be exerted when interpreting results based on statistical tests in this study.

Our study has a number of limitations. Although the assumption of 100% absorption in the gastro-intestinal tract is supported by studies of other lipophilic persistent organic pollutants like dioxins and polychlorinated biphenyls (McLachlan, 1993), it is possible that absorption is lower for p,p' -DDT/E. The assumption that p,p' -DDT/E levels are equal in maternal and breast milk lipids may also be an over-simplification, as suggested in a study of paired serum and milk samples (LaKind et al., 2009). Many parameters including diffusion across the membranes and lipid composition in serum and breast milk could influence p,p' -DDT/E partitioning and explain the observed underestimation of breast milk levels. However, we decided not to calibrate the pharmacokinetic model parameters like absorption and milk:serum partition coefficients based on measured children's levels to avoid overfitting, which would have reduced model generalizability. Another limitation of our study is that we did not include individual-specific data on enzyme polymorphism and folate status, which may influence half-life (Denk and Milutinovic, 2018). Finally, it is important to note that R^2 s for measured and estimated children's levels may also account for other environmental and lifestyle factors that are shared between mothers and children. For example, consumption of chicken foraging around the dwelling (Van Dyk et al., 2010) and their eggs (Bouwman et al., 2015) may be related among family members and contribute similarly to maternal and child intake.

We believe the pharmacokinetic model used herein could be employed to estimate serum levels in children participating to the VHEMBE study whose serum was not analyzed. Not only can the pharmacokinetic model estimate levels at 12 months and 24 months, it can generate complete time courses of children's levels across the first two years of life. Multiple exposure metrics can be derived from these time courses like monthly levels, maximum levels, and area under the curve (cumulative exposure) during specific periods (see Fig. 1). Such exposure metrics will allow evaluating associations between exposures during prenatal and postnatal windows of development and health outcomes, an approach that has previously been used to identify windows of vulnerability to toxic insults (Iszatt et al., 2015; Jusko et al., 2016; Rosenquist et al., 2017; Verner et al., 2010). Estimated p,p' -DDT levels could be mathematically corrected for the systematic overestimation (lack of model accuracy) based on calculated mean relative errors (see Table 4). For example, multiplying estimated children's p,p' -DDT levels by 0.69 would correct for the average 44% overestimation by the pharmacokinetic model, while maintaining model precision.

In conclusion, our study showed that children from a sprayed area in South Africa had serum p,p' -DDT/E at 12 months and 24 months of age exceeding their mothers' at delivery. We also demonstrated that a published pharmacokinetic model of placental and lactational transfer allows estimating children's serum p,p' -DDT/E levels with a precision ranging from 75 to 82%. However, indicators of accuracy suggested that estimated children's serum p,p' -DDT were overestimated. Because the pharmacokinetic model was developed a priori, and because results suggested that children's body burden during the first two years is mostly

attributable to mother-child transfer, the pharmacokinetic model could be considered to be transportable to comparable populations with similar cultural practices and where IRS with DDT is still used to fight malaria.

Acknowledgments

We would like to thank study participants. This study was funded by the National Institute of Environmental Health Sciences, Award ID: 1R01ES020360-01. Marc-André Verner is the recipient of a Research Scholar J1 Award from the Fonds de recherche du Québec – Santé (FRQS). Jonathan Chevrier holds a Canada Research Chair in Environmental Health Sciences.

References

- Aylward LL, Hays SM, Kirman CR, Marchitti SA, Kenneke JF, English C, et al., 2014 Relationships of chemical concentrations in maternal and cord blood: a review of available data. *J. Toxicol. Environ. Health B Crit. Rev* 17 (3), 175–203. [PubMed: 24749481]
- Barr JR, Maggio VL, Barr DB, Turner WE, Sjodin A, Sandau CD, et al., 2003 New high-resolution mass spectrometric approach for the measurement of polychlorinated biphenyls and organochlorine pesticides in human serum. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci* 794 (1), 137–148.
- Bouwman H, Kylin H, 2009 Malaria control insecticide residues in breast milk: the need to consider infant health risks. *Environ. Health Perspect.* 117 (10), 1477–1480. [PubMed: 20019894]
- Bouwman H, Becker PJ, Cooppan RM, Reinecke AJ, 1992 Transfer of DDT used in malaria control to infants via breast milk. *Bull. World Health Organ* 70 (2), 241–250. [PubMed: 1632883]
- Bouwman H, Bornman R, van Dyk C, Barnhoorn I, 2015 First report of the concentrations and implications of DDT residues in chicken eggs from a malaria-controlled area. *Chemosphere* 137, 174–177. [PubMed: 26197434]
- Caspersen IH, Kvale HE, Haugen M, Brantsaeter AL, Meltzer HM, Alexander J, et al., 2016 Determinants of plasma PCB, brominated flame retardants, and organochlorine pesticides in pregnant women and 3 year old children in The Norwegian Mother and Child Cohort Study. *Environ. Res* 146, 136–144. [PubMed: 26749444]
- Chevrier J, Rauch S, Crause M, Obida M, Gaspar F, Bornman R, et al., 2018 Maternal exposure to DDT and pyrethroids and birth outcomes among residents of an area sprayed for malaria control participating in the VHEMBE birth cohort study. *Am. J. Epidemiol* 10.1093/aje/kwy143. (in press).
- Coker E, Chevrier J, Rauch S, Bradman A, Obida M, Crause M, et al., 2018 Association between prenatal exposure to multiple insecticides and child body weight and body composition in the VHEMBE South African birth cohort. *Environ. Int* 113, 122–132. [PubMed: 29421401]
- Denk MK, Milutinovic NS, 2018 Reductive dehalogenation of DDT with folate models: formation of the DDT metabolite spectrum under biomimetic conditions. *Chemosphere* 191, 408–411. [PubMed: 29054081]
- Eskenazi B, Chevrier J, Rosas LG, Anderson HA, Bornman MS, Bouwman H, et al., 2009 The Pine River statement: human health consequences of DDT use. *Environ. Health Perspect* 117 (9), 1359–1367. [PubMed: 19750098]
- Eskenazi B, An S, Rauch SA, Coker ES, Maphula A, Obida M, et al., 2018 Prenatal exposure to DDT and pyrethroids for malaria control and child neurodevelopment: the VHEMBE cohort, South Africa. *Environ. Health Perspect* 126 (4), 047004. [PubMed: 29648420]
- Fisher M, Arbuckle TE, Liang CL, Leblanc A, Gaudreau E, Foster WG, et al., 2016 Concentrations of persistent organic pollutants in maternal and cord blood from the maternal-infant research on environmental chemicals (MIREC) cohort study. *Environ. Health* 15 (1), 59. [PubMed: 27142700]
- Gascon M, Vrijheid M, Gari M, Fort M, Grimalt JO, Martinez D, et al., 2015 Temporal trends in concentrations and total serum burdens of organochlorine compounds from birth until adolescence and the role of breastfeeding. *Environ. Int* 74, 144–151. [PubMed: 25454231]
- Gaspar FW, Chevrier J, Bornman R, Crause M, Obida M, Barr DB, et al., 2016 Corrigendum to “Undisturbed dust as a metric of long-term indoor insecticide exposure: residential DDT contamination from indoor residual spraying and its association with serum levels in the

- VHEMBE cohort” [Environ. Int. 85C (2015) 163–167]. Environ. Int 94, 778–783. [PubMed: 27236616]
- Gaspar FW, Chevrier J, Quiros-Alcala L, Lipsitt JM, Barr DB, Holland N, et al., 2017 Levels and determinants of DDT and DDE exposure in the VHEMBE cohort. Environ. Health Perspect 125 (7), 077006. [PubMed: 28696207]
- Gyalpo T, Fritsche L, Bouwman H, Bornman R, Scheringer M, Hungerbuhler K, 2012 Estimation of human body concentrations of DDT from indoor residual spraying for malaria control. Environ. Pollut 169, 235–241. [PubMed: 22682611]
- Haddad S, Ayotte P, Verner MA, 2015 Derivation of exposure factors for infant lactational exposure to persistent organic pollutants (POPs). Regul. Toxicol. Pharmacol 71 (2), 135–140. [PubMed: 25510832]
- Huang J, Eskenazi B, Bornman R, Rauch S, Chevrier J, 2018 Maternal peripartum serum DDT/E and urinary pyrethroid metabolite concentrations and child infections at age 2 years in the VHEMBE birth cohort: associations and modifiers. Environ. Health Perspect. 126 (6), 067006. [PubMed: 29906263]
- Izatt N, Stigum H, Verner MA, White RA, Govarts E, Murinova LP, et al., 2015 Prenatal and postnatal exposure to persistent organic pollutants and infant growth: a pooled analysis of seven European birth cohorts. Environ. Health Perspect 123 (7), 730–736. [PubMed: 25742056]
- Jusko TA, De Roos AJ, Lee SY, Thevenet-Morrison K, Schwartz SM, Verner MA, et al., 2016 A birth cohort study of maternal and infant serum PCB-153 and DDE concentrations and responses to infant tuberculosis vaccination. Environ. Health Perspect 124 (6), 813–821. [PubMed: 26649893]
- Lackmann GM, 2006 Human milk, environmental toxins and pollution of our infants: disturbing findings during the first six months of life. Int. J. Biomed. Sci 2 (2), 178–183. [PubMed: 23674980]
- Lackmann GM, Schaller KH, Angerer J, 2004 Organochlorine compounds in breast-fed vs. bottle-fed infants: preliminary results at six weeks of age. Sci. Total Environ 329 (1–3), 289–293. [PubMed: 15262173]
- LaKind JS, Berlin CM, Jr., Sjodin A, Turner W, Wang RY, Needham LL, et al., 2009 Do human milk concentrations of persistent organic chemicals really decline during lactation? Chemical concentrations during lactation and milk/serum partitioning. Environ. Health Perspect 117 (10), 1625–1631. [PubMed: 20019916]
- Mabaso ML, Sharp B, Lengeler C, 2004 Historical review of malarial control in southern African with emphasis on the use of indoor residual house-spraying. Tropical Med. Int. Health 9 (8), 846–856.
- McLachlan MS, 1993 Digestive tract absorption of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls in a nursing infant. Toxicol. Appl. Pharmacol 123 (1), 68–72. [PubMed: 8236263]
- Phillips DL, Pirkle JL, Burse VW, Bernert JT, Jr., Henderson LO, Needham LL, 1989 Chlorinated hydrocarbon levels in human serum: effects of fasting and feeding. Arch. Environ. Contam. Toxicol 18 (4), 495–500. [PubMed: 2505694]
- Ribas-Fito N, Grimalt JO, Marco E, Sala M, Mazon C, Sunyer J, 2005 Breastfeeding and concentrations of HCB and p,p'-DDE at the age of 1 year. Environ. Res 98 (1), 8–13. [PubMed: 15721878]
- Ritter R, Scheringer M, MacLeod M, Schenker U, Hungerbuhler K, 2009 A multi-individual pharmacokinetic model framework for interpreting time trends of persistent chemicals in human populations: application to a postban situation. Environ. Health Perspect 117 (8), 1280–1286. [PubMed: 19672409]
- Rosenquist AH, Hoyer BB, Julvez J, Sunyer J, Pedersen HS, Lenters V, et al., 2017 Prenatal and postnatal PCB-153 and p,p'-DDE exposures and behavior scores at 5–9 years of age among children in Greenland and Ukraine. Environ. Health Perspect 125 (10), 107002. [PubMed: 28974479]
- Smith D, 1999 Worldwide trends in DDT levels in human breast milk. Int. J. Epidemiol 28 (2), 179–188. [PubMed: 10342677]
- Van Dyk JC, Bouwman H, Barnhoorn IE, Bornman MS, 2010 DDT contamination from indoor residual spraying for malaria control. Sci. Total Environ 408 (13), 2745–2752. [PubMed: 20381127]

- Verner MA, Ayotte P, Muckle G, Charbonneau M, Haddad S, 2009 A physiologically based pharmacokinetic model for the assessment of infant exposure to persistent organic pollutants in epidemiologic studies. *Environ. Health Perspect* 117 (3), 481–487. [PubMed: 19337526]
- Verner MA, Plusquellec P, Muckle G, Ayotte P, Dewailly E, Jacobson SW, et al., 2010 Alteration of infant attention and activity by polychlorinated biphenyls: unravelling critical windows of susceptibility using physiologically based pharmacokinetic modeling. *Neurotoxicology* 31 (5), 424–431. [PubMed: 20609431]
- Verner MA, Sonneborn D, Lancz K, Muckle G, Ayotte P, Dewailly E, et al., 2013 Toxicokinetic modeling of persistent organic pollutant levels in blood from birth to 45 months of age in longitudinal birth cohort studies. *Environ. Health Perspect* 121 (1), 131–137. [PubMed: 23086694]
- WHOPES, 2015 WHO recommended insecticides for indoor residual spraying against malaria vectors. Available: http://www.who.int/whopes/Insecticides_IRS_2_March_2015.pdf?ua=1.
- Wolff MS, Zeleniuch-Jacquotte A, Dubin N, Toniolo P, 2000 Risk of breast cancer and organochlorine exposure. *Cancer Epidemiol. Biomark. Prev* 9 (3), 271–277.
- Wrottesley SV, Pisa PT, Norris SA, 2017 The influence of maternal dietary patterns on body mass index and gestational weight gain in urban black South African women. *Nutrients* 9 (7).




Fig. 1. Example of a pharmacokinetic profile for a mother-child dyad where the mother was 25 years of age at delivery, had a serum *p,p'*-DDT concentration of 385 ng/g lipids at delivery and breastfed her child exclusively for 6 months.




Fig. 2. Measured p,p' -DDT/E levels vs. levels estimated using the pharmacokinetic model. Note: axes are log₁₀-transformed. Dotted line represents expected concentrations (where measured levels = simulated levels) based on an intercept of 0 and a slope of 1.

Table 1

Characteristics of study participants, and DDT/E levels in serum and breast milk.

Characteristic	N (%)	GM	GSD	Range
Maternal age at delivery (years)		26	1.2	18–39
Maternal body weight at delivery (kg)		74	1.2	55–119
Maternal body weight (kg) 12 months after delivery		67	1.3	46–118
Maternal body weight (kg) 24 months after delivery		69	1.2	44–113
Sex of child				
Female	20 (43)			
Male	27 (57)			
Birth weight (kg)		3.1	1.2	1.7–4.1
Child weight at 12months (kg)		9.1	1.2	6.6–12.1
Child weight at 24 months (kg)		11.2	1.1	9.1–13.5
Duration of exclusive breastfeeding (months)		1.7	3.3	0.1–8.0
Duration of total breastfeeding (months)		18.7	1.2	11.9–25.3
Maternal serum levels (ng/g lipids)				
<i>p,p'</i> -DDT		90	4.6	< LOD-1816
<i>p,p'</i> -DDE		338	3.9	31–7440
Children's serum levels at 12months (ng/g lipids)				
<i>p,p'</i> -DDT		229	5.5	11–11,893
<i>p,p'</i> -DDE		1215	4.4	123–19,772
Children's serum levels at 24 months (ng/g lipids)				
<i>p,p'</i> -DDT		221	5.2	16–15,299
<i>p,p'</i> -DDE		1377	4.1	145–33,605
Breast milk levels at 12 months (ng/g lipids)				
<i>p,p'</i> -DDT		114	5.4	8–5301
<i>p,p'</i> -DDE		409	4.6	28–5984

GM: geometric mean; GSD: geometric standard deviation; LOD: limit of detection.

Table 2

Ratio between children's serum levels at 12 months and 24 months, and maternal levels at delivery.

Compound	N	Ratio child (12 mo)/mother GM (min-max)	Ratio child (24 mo)/mother GM (min-max)
<i>p,p'</i> -DDT	46	2.5 (0.7–42)	2.4 (0.3–34)
<i>p,p'</i> -DDE	47	3.6 (0.8–32)	4.1 (0.9–25)

GM: geometric mean.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3Pearson correlations between log₁₀-transformed serum and breast milk *p,p'*-DDT/E levels.

		<i>p,p'</i> -DDT				<i>p,p'</i> -DDE			
		Maternal serum	Breast milk	Child serum (12 mo)	Child serum (24 mo)	Maternal serum	Breast milk	Child serum (12 mo)	Child serum (24 mo)
<i>p,p'</i> -DDT	Maternal serum	-							
	Breast milk	0.890	-						
	Child, serum (12 mo)	0.878	0.984	-					
	Child, serum (24 mo)	0.866	0.967	0.984	-				
<i>p,p'</i> -DDE	Maternal serum	0.895	0.840	0.880	0.788	-			
	Breast milk	0.818	0.917	0.799	0.850	0.916	-		
	Child serum (12 mo)	0.827	0.917	0.907	0.877	0.901	0.983	-	
	Child serum (24 mo)	0.829	0.929	0.924	0.923	0.889	0.963	0.981	-

Table 4

Linear regression of measured vs. estimated p,p' -DDT/E levels. Levels were \log_{10} -transformed prior to analyses.

	Intercept (95% CI)	β (95% CI)	R^2	Mean relative error (%) (95% CI)
<i>p,p'</i> -DDT				
Children's serum levels (12 months)	0.051 (-0.349,0.451)	0.968 (0.806, 1.130)	0.77	36 (11, 61)
Children's serum levels (24 months)	0.096 (-0.317, 0.510)	0.938 (0.772, 1.105)	0.75	51 (13, 89)
Breast milk levels (12 months)	0.320(0.020, 0.619)	0.971 (0.814, 1.128)	0.78	-31 (-44, -18)
<i>p,p'</i> -DDE				
Children's serum levels (12 months)	0.320 (-0.079, 0.719)	0.936 (0.804, 1.069)	0.82	-11(-29, 6)
Children's serum levels (24 months)	0.475 (0.056, 0.894)	0.898 (0.760, 1.037)	0.79	-17 (-35, 2)
Breast milk levels (12 months)	0.311 (-0.011,0.634)	0.975 (0.842, 1.107)	0.83	-32 (-47, -17)