

UC Office of the President

Recent Work

Title

Hand Nicotine and Cotinine In Children Exposed to Cigars: A Pilot Study

Permalink

<https://escholarship.org/uc/item/9hh9k3dg>

Journal

Tobacco Regulatory Science, 7(3)

ISSN

2333-9748

Authors

Mahabee-Gittens, E Melinda
Matt, Georg E
Jandarov, Roman J
[et al.](#)

Publication Date

2021-05-01

DOI

10.18001/trs.7.3.2

Peer reviewed

Hand Nicotine and Cotinine In Children Exposed to Cigars: A Pilot Study

E. Melinda Mahabee-Gittens, MD, MS
Georg E. Matt, PhD
Roman J. Jandarov, PhD
Ashley L. Merianos, PhD

Objectives: Past research has not examined secondhand and thirdhand smoke (THS) exposure in children of cigar smokers. We examined hand nicotine and cotinine levels in children of cigar smokers to explore the contribution of cigar smoke to tobacco smoke exposure (TSE). **Methods:** Participants were children (N = 24; mean (SD) age = 6.5 (3.6) years) whose parents smoked cigars only or poly-used cigars and/or cigarettes. Primary outcomes were hand nicotine and urinary cotinine levels. **Results:** All children had detectable hand nicotine (range: 7.6-312.5ng/wipe) and cotinine (range: 0.3-100.3ng/ml). Positive correlations were found between hand nicotine and cotinine ($r = 0.693$, $p = .001$), hand nicotine and parents who also smoked cigarettes ($r = 0.407$, $p = .048$), and hand nicotine and number of smokers around the child ($r = 0.436$, $p = .03$). Hand nicotine ($r = -0.464$, $p = .02$), but not cotinine ($r = -0.266$, $p = .26$), was negatively correlated with child age. Multiple regression results indicated a positive association between hand nicotine and cotinine ($p = .002$; semi-partial $r^2 = 0.415$), irrespective of child age. **Conclusions:** The significant association of hand nicotine with urinary cotinine suggests that THS pollution should be assessed in evaluating children's overall TSE to cigars and other tobacco products, and hand nicotine may be a proxy for overall TSE. Younger children may have increased THS pollutant uptake.

Key words: adolescent/youth; cigar/cigarillo; secondhand smoke; thirdhand smoke; tobacco exposure; toxicant
Tob Regul Sci.™ 2021;7(3):170-176
DOI: doi.org/10.18001/TRS.7.3.2

Although cigar use rates remain relatively high at 3.9% among adults in the United States (US),¹ past studies have not yet examined tobacco smoke exposure (TSE) in children exposed to cigars. Cigar use rates are highest in males, younger adults, and persons of low socioeconomic status.¹⁻³ Racial/ethnic disparities exist in cigar and cigarillo use with highest levels among black (4.9%) compared to white adults (4.1%).¹ The estimated annual healthcare services and expenditures attributable to cigar smoking are high amounting to \$1.8 billion, of which \$284 million is attributable to cigar only use and \$1.5 billion due to poly-use of cigars and other tobacco products.⁴

Given the disparities associated with cigar use, it is likely that males and non-Hispanic Blacks have the highest cigar-related healthcare expenditures. TSE rates are highest in children who are non-Hispanic black, low-income, or who live in rented homes.⁵ Cigar smoke contains higher levels of harmful pollutants compared to cigarettes,⁴ which may result in higher adverse effects on children exposed to cigar smoke compared to cigarette smoke. Moreover, availability, advertising, and marketing for little cigars and cigarillos is more widespread in neighborhoods where black or low-income populations reside, potentially leading to increased cigar use and exposure.⁶

E. Melinda Mahabee-Gittens, Professor, Division of Emergency Medicine, Cincinnati Children's Hospital Medical Center, University of Cincinnati College of Medicine, Cincinnati, OH, United States. Georg E. Matt, Professor, Department of Psychology, San Diego State University, San Diego, CA, United States. Roman J. Jandarov, Assistant Professor, Department of Environmental Health, Division of Biostatistics and Bioinformatics, University of Cincinnati College of Medicine, Cincinnati, OH, United States. Ashley L. Merianos, Associate Professor, School of Human Services, University of Cincinnati, Cincinnati, OH, United States.
Correspondence Dr Mahabee-Gittens; melinda.mahabee-gittens@cchmc.org

After tobacco has been smoked, chemical residue from secondhand smoke (SHS) remains on surfaces, clothes and skin, becomes embedded in materials, and accumulates in dust. This aged SHS, or thirdhand smoke (THS), contains nicotine and other tobacco smoke toxicants.⁷ Nicotine residue is picked up by children on their hands even when active smoking is not occurring.⁸ The association between hand nicotine and cotinine (ie, nicotine metabolite),⁹ suggests that hand nicotine is a marker of overall TSE (SHS+THS) in children's environments.⁸ Research indicates that all children of cigarette smokers, even those whose parents enforce smoking bans, have detectable hand nicotine and cotinine with higher levels in younger children and those exposed to more cigarettes.^{5,8,10-13}

Whereas the nicotine concentration in cigars compared to cigarettes is similar, the total tobacco content in cigars is higher than cigarettes, and the smoke emissions and TSE patterns from cigars and cigarettes may vary.¹⁴ TSE biomarker levels differ by cigar type and use patterns and the use of other tobacco products (eg, dual- or poly-use with cigars, cigarettes, and/or marijuana).^{15,16} Thus, it is likely that biomarker levels of SHS and THS exposure differ in children who are exposed to cigars and other tobacco and non-tobacco products; an understudied area. To help fill this gap, we conducted a pilot study to investigate hand nicotine and urinary cotinine biomarker levels in children whose parents smoke cigars only or dual- or poly-used cigar products and cigarettes, and their relationship with child characteristics, parent-reported smoking, and TSE patterns.

METHODS

Child and parent dyads in this study were part of a 2-group randomized controlled trial, "Healthy Families" (clinicaltrials.gov:NCT02531594),¹⁷ that was conducted in one of 2 pediatric emergency departments (PED) or urgent cares (UC) at a large children's hospital. This hospital IRB-approved study enrolled the study subset, all of whom had public insurance or were self-pay and presented to the PED/UC with a potential TSE-related complaint (eg, cough). Parents completed electronic assessments to determine tobacco use type and amount smoked around the child by parent and

household members (Black and Milds (B&Ms), other cigars, cigarettes, electronic cigarettes) and whether home smoking bans were in place. Included children were ages 0-14 years, nonsmokers, had parents who reported smoking B&Ms and/or other cigars (N = 19), and dual- or poly-use with combustible cigarettes (N= 5). No parents currently used electronic cigarettes. There were 88 children with complete data on hand nicotine and parent tobacco use. Of those, 24 (27.3%) had parents who smoked B&Ms and/or cigars and were included in the present study. No differences in participant characteristics were found between those included (N = 24) and excluded (N = 64) in the study. Although 19 parents were B&M and/or other cigar smokers, 17 (89.5%) of these children were around 1-4 cigarette smokers (25th-50th-75th: 1-2-3) in the past week.

The palm and volar aspect of children's dominant hands were wiped by research coordinators and analyzed for nicotine with a level of quantification (LOQ) = 0.1ng/wipe. Field blank hand wipes were collected to adjust for potential sample contamination¹⁸ (range: < LOQ-6.7ng/wipe). A sub-sample provided a urine sample (N = 20) that was analyzed for cotinine (LOQ = 0.05ng/ml) using liquid chromatography-tandem mass spectrometry.¹⁹ Urine samples were not collected from the full sample due to difficulty obtaining samples from young children. We assessed sociodemographics and parent-reported smoking and TSE patterns.

Data Analysis

Logarithmic transformations were performed of hand nicotine and cotinine to control for skew and unequal variances. We report geometric means (GeoM), confidence intervals (CI), medians (Mdn), and interquartile ranges (IQR). Simple linear regression and correlation analyses were conducted to examine associations between child characteristics, parent-reported smoking, TSE patterns, hand nicotine, and cotinine. We examined a multiple regression model to assess the association of hand nicotine and child age with cotinine as our outcome. All statistical tests were 2-tailed with the level of significance set at .05.

RESULTS

Table 1 shows participant characteristics. All children had detectable hand nicotine rang-

Table 1
PED/UC Patient Sociodemographic and Self-reported
Smoking and TSE Patterns by Hand Nicotine and Urinary Cotinine Levels

Variable	N (%) ^b	Hand Nicotine Concentration (ng/wipe; N = 24) ^a			Urinary Cotinine Concentration (ng/ml; N = 20)		
		GeoM (95%CI)	Mdn (IQR)	P-value ^c	GeoM (95%CI)	Mdn (IQR)	P-value ^c
<i>Sociodemographic Characteristics</i>							
Child Age, M (SD)	6.5 (3.6)	-	-	.02	-	-	.26
25th-50th-75th	4-6.5-8.75						
Child Sex				.89			.64
Male	10 (41.7)	45.2 (23.1-87.9)	80.3 (20.8-90.5)		5.6 (1.4-17.0)	3.3 (1.2-13.9)	
Female	14 (58.3)	47.8 (26.9-84.3)	44.3 (23.7-108.5)		7.7 (2.6-19.8)	7.5 (1.8-10.3)	
Child Race							
White, non-Hispanic	3 (12.5)	56.9 (15.2-206.1)	61.8 (-)	Ref	2.1 (0.7-35.5)	2.1 (-)	Ref
Black, non-Hispanic	20 (83.3)	49.0 (31.3-76.6)	56.4 (24.0-97.9)	.80	7.5 (3.4-15.3)	7.3 (1.7-13.9)	.31
Other, non-Hispanic	1 (4.2)	9.3 (-)	-	-	-	-	-
Parent Education Level				.02			.16
≤ High school graduate	11 (45.8)	74.8 (44.1-126.2)	90.2 (31.6-102.0)		11.0 (3.8-29.3)	7.1 (3.7-32.3)	
≥ Some college	13 (54.2)	31.3 (17.9-54.1)	32.5 (12.7-66.6)		4.3 (1.3-11.5)	1.8 (0.9-10.3)	
Income Level				.66			.48
< \$15,000	20 (83.3)	44.9 (28.6-70.2)	44.3 (24.0-91.2)		5.9 (2.5-12.5)	5.0 (1.6-12.9)	
≥ \$15,000	4 (16.7)	56.8 (11.3-268.6)	73.1 (20.0-122.7)		10.7 (-0.1-122.8)	8.4 (2.3-50.5)	
Housing Type				.81			.40
Single-family	5 (20.8)	51.3 (16.1-159.1)	91.5 (19.1-101.3)		4.0 (1.2-10.6)	3.3 (2.0-7.9)	
Multi-Unit/Apartment	19 (79.2)	45.6 (28.4-72.8)	53.4 (23.8-89.9)		7.9 (3.0-18.4)	7.5 (1.6-14.0)	

(continued on next page)

ing from 7.6-312.5ng/wipe (GeoM = 46.7, 95% CI:[31.3;69.5]; Mdn = 56.4, IQR = 24.0-91.2) and urinary cotinine ranging from 0.3-100.3ng/ml (GeoM = 6.7, 95% CI:[3.2;13.0]; Mdn = 6.4, IQR = 1.6-12.9).

Children whose parents completed ≤ high school (GeoM = 74.8, p = .02) had higher mean hand nicotine than those whose parents completed ≥ some college (GeoM = 31.3). Children who lived with parents who smoked B&Ms and/or cigars with cigarettes had higher mean hand nicotine (GeoM = 96.9, p = .048) than those who lived with parents who smoked B&Ms and/or cigars (GeoM = 38.5). Children without home smoking bans (GeoM = 11.4, 95% CI:[5.3;23.3], p = .006)

had higher mean cotinine than children living in homes with smoking bans (GeoM = 1.5ng/ml, 95% CI:[0.2;4.4]).

Hand nicotine (r = -0.464, p = .02), but not cotinine (r = -0.266, p = .26), was negatively correlated with child age. Of note, a one-year-old had the highest hand nicotine level. Positive correlations were found between hand nicotine level and parents who smoked cigars and cigarettes (vs cigars only) (r = 0.407, p = .048), and hand nicotine and cumulative number of tobacco smokers (ie, cigarillo, cigar, cigarette smokers) around the child in the past week (r = 0.436, p = .03). Hand nicotine was positively correlated with cotinine (r = 0.693, p = .001). A negative correlation was found

Table 1 (continued)
PED/UC Patient Sociodemographic and Self-reported
Smoking and TSE Patterns by Hand Nicotine and Urinary Cotinine Levels

Variable	N (%) ^b	Hand Nicotine Concentration (ng/wipe; N = 24) ^a			Urinary Cotinine Concentration (ng/ml; N = 20)		
		GeoM (95%CI)	Mdn (IQR)	P-value ^c	GeoM (95%CI)	Mdn (IQR)	P-value ^c
<i>Self-reported Smoking and TSE Patterns</i>							
Parent Specific Tobacco Product Type Use Patterns							
Exclusive B&M Use	17 (70.8)	36.1 (22.4-57.8)	31.6 (17.1-90.0)	-	5.8 (2.6-11.9)	6.4 (1.7-11.1)	-
Exclusive Cigar Use	1 (4.2)	136.0 (-)	-	-	85.0 (-)	-	-
B&M and Cigar Use	1 (4.2)	32.5 (-)	-	-	1.6 (-)	-	-
B&M and Cigarette Use	4 (16.6)	72.2 (46.9-110.8)	74.5 (55.4-91.0)	-	3.2 (-)	-	-
B&M, Cigar, and Cigarette Use	1 (4.2)	312.5 (-)	-	-	73.5 (-)	-	-
Parent Smokes Cigarettes				.048			.85
No	19 (79.2)	38.5 (24.6-59.9)	32.5 (21.1-90.2)		6.5 (2.8-13.5)	6.4 (1.6-12.9)	
Yes	5 (20.8)	96.9 (40.5-229.7)	89.8 (57.4-169.3)		7.6 (-0.3-106.9)	5.1 (1.3-44.1)	
Home Smoking Ban				.16			.006
No	17 (70.8)	55.8 (34.3-90.3)	71.9 (25.4-101.3)		11.4 (5.3-23.3)	8.7 (4.1-21.4)	
Yes	7 (29.2)	30.2 (13.3-67.3)	32.5 (13.8-61.8)		1.5 (0.2-4.4)	1.2 (0.6-2.6)	
No. Tobacco Smokers Around Child, 25th-50th-75th^{d,e}	2.25-3.5-4			.03			.10
No. B&M Smokers Around Child, 25th-50th-75th^e	1-1-1	-	-	.86	-	-	.82
No. Cigar Smokers Around Child, 25th-50th-75th^e	0-0-0	-	-	.12	-	-	.19
No. Cigarette Smokers Around Child, 25th-50th-75th^e	1-2-3	-	-	.06	-	-	.18
No. Cigarettes Smoked Around Child, 25th-50th-75th^e	0-2.5-9.75	-	-	.18	-	-	.14

Note.

Abbreviations: GeoM, geomean; CI, confidence interval; Mdn, median; IQR, interquartile range; ref, reference category; B&Ms, Black and Milds.

N = 24 and missing values excluded.

^a Hand nicotine concentrations have been corrected for field blank contamination.

^b N and percent unless noted otherwise.

^c All p-values refer to linear regression model results.

^d Tobacco smokers includes all smokers who use cigarillos, cigars, and/or cigarettes.

^e Measures include all smokers (including the parent) the child was around in any location (home, car, outside).

between cotinine and having a home smoking ban ($r = -0.588$, $p = 0.006$), but no other associations were found between cotinine and the other covariates (ie, child age, number of cigarettes smoked).

A multiple linear regression model with cotinine as the outcome ($R^2 = 0.49$, $F(2,17) = 8.04$, $p = .003$) revealed that hand nicotine (semi-partial r^2

$= 0.415$; $p = .002$) had a significant positive linear association with cotinine independent of child age. That is, independent of age, higher hand nicotine levels were associated with higher cotinine levels. There was no association between child age (semi-partial $r^2 = 0.006$; $p = .66$) and cotinine independent of hand nicotine.

DISCUSSION

All nonsmoking children of cigar smokers had detectable hand nicotine (GeoM = 46.7 ng/wipe, 95% CI:[31.3;69.5]). These levels are in stark contrast to hand nicotine levels of children in THS-free homes of nonsmokers (GeoM: 0.7; 95% CI:[0.2;1]) and similar to those in homes of cigarette smokers with indoor smoking bans (GeoM:17.2; 95% CI:[7.1;39.9])²⁰ and without indoor smoking bans (GeoM:86.4; 95% CI:[61.0;122]).⁸ Consistent with hand nicotine levels, urinary cotinine of 6.7ng/ml was comparable to children of cigarette smokers from our previous work.¹² Cotinine levels contrast with much lower levels in children of nonsmokers in THS-free homes (GeoM=0.08ng/ml, 95% CI:[0.0;0.17]).²⁰

There was a strong positive correlation between cotinine and hand nicotine supporting the hypothesis that hand nicotine represents a measure of children's overall TSE via inhalation (eg, SHS) or via dermal absorption or ingestion (eg, from THS in dust or on surfaces).⁸ Hand nicotine and urinary cotinine, however, are not redundant measures of TSE as indicated by the negative association of age with hand nicotine and the lack of association with cotinine. The negative correlation observed with child age and hand nicotine has not been reported previously. However, we have reported higher hand nicotine levels in 2-4-year-olds compared to younger and older children,¹⁰ which is suspected to be the result of increased exploratory behaviors and contact with THS-polluted surfaces in this age group.²¹ The finding that children who lived with parents who smoked cigars and cigarettes had higher hand nicotine levels may be due to increased overall THS levels in their homes. However, future studies should include larger numbers of parents who dual-used cigarettes and cigars and parents who smoke cigarettes only so that hand nicotine levels in these children can be compared. Finally, the finding that children in homes without smoking bans had higher mean cotinine is consistent with prior research on the effects of indoor smoking on SHS exposure.²² Whereas it is known that homes with indoor smoking bans also have lower nicotine levels on surface levels,²³ we did not observe lower hand nicotine levels in children who lived in homes with smoking bans compared to those without bans. This could be due to higher

levels of THS pollution from previous smokers in homes with smoking bans,^{23,24} or differences in protective TSE practices in younger children's homes.²⁵ These findings need to be examined further in larger samples of varying sociodemographics and parental tobacco product use.

Study limitations include the small sample size which did not allow comparisons between children whose parents used B&Ms, other cigars, cigarettes, or who were dual- or poly-users. There was no control group of children of nonsmokers; thus, we were unable to differentiate between THS and SHS exposure. However, because nicotine can only be derived from tobacco products, nicotine on children's hands can be attributed only to SHS and THS in their environment, representing overall TSE. Further, given the varying ranges of nicotine content, additives and flavors in cigars,¹⁴ future TSE biomarker studies should employ specific, detailed assessments about cigar use, including brands, cigar types (eg, traditional or little cigars, cigarillos), smoking topography (eg, puffs, duration), and locations of cigars smoked, all of which affect children's TSE. Although co-use of marijuana is common in tobacco product users,^{16,26} marijuana use was not assessed, which could have affected our results. Furthermore, with the exception of B&Ms, we did not specifically assess types of cigars used which could have affected our study findings. Participants may have underreported cigar use or mistaken B&Ms for cigarettes due to their similar physical characteristics, but this was not assessed.^{27,28} Additionally, recency and frequency of handwashing likely affects the amount of hand nicotine,²⁹ and future studies are advised to collect such information to adjust measured hand nicotine levels. Finally, there are other exposure-relevant variables (eg, amount of indoor time, distance to exposure source, hand-to-mouth behaviors, THS pollutant levels)³⁰⁻³² that future studies should incorporate to improve models of SHS and THS exposure to cigar smoke.

In conclusion, this study highlights the role of cigar use in children's overall TSE as it contributes to SHS and THS pollution of home environments. This study suggests that cigar use contributes to overall TSE through SHS and THS exposure. To study the unique and combined effects on exposure, future studies should measure different types of tobacco product use (eg, cigarettes, little and

traditional cigars, dual- and poly-use) and measure exposure through cotinine and hand nicotine to assess differential contributions of SHS and THS.

IMPLICATIONS FOR TOBACCO REGULATION

Cigar use in children's homes plays an important role in child TSE. The statistically significant association of hand nicotine with cotinine levels suggests that THS pollution at home should be assessed in evaluating children's overall TSE to toxicants in cigars and other tobacco products, and hand nicotine may be a useful proxy for overall TSE. Moreover, the statistically significant association between hand nicotine levels and child age suggests that younger children may have increased uptake of THS pollutants from their environments. Future studies should include biomarkers of SHS and THS exposure in children of cigar smokers with detailed assessments to differentiate biomarker patterns by type, frequency, and exposure patterns of cigar and other tobacco products. Given the lower regulation of cigar products compared to cigarettes,³³ the tobacco industry's push to place little cigars and cigarillos that have similar features to cigarettes (ie, shape, size, packaging) but are sold at a lower cost on the market, lower taxation, and the availability of flavors, there is a need for policies to regulate the sale of cigars that are equivalent to regulation of cigarettes.^{6,28,34} Without these regulations, there is a risk of increased harm to vulnerable child populations that already have high TSE. Thus, policies are needed to decrease the availability and use of cigar products to protect children from toxicants due to cigar smoke exposure.

Human Subjects Approval Statement

This study was approved by the Institutional Review Board of Cincinnati Children's Hospital Medical Center.

Conflict of Interest Disclosure Statement

The authors have no conflicts of interest to disclose.

Acknowledgements

The authors thank Dr. Hoh, Dr. Quintana, and the laboratory staff at the San Diego State University

Environmental Health Laboratory for their role in the handwipe analyses. This research was funded by the National Institute of Environmental Health Sciences (NIH Grant Numbers R01ES027815, R01ES030743 and R21ES032161) and the National Institute on Drug Abuse (NIH Grant Number K01DA044313). The National Institutes of Health had no role in the design and conduct of the study; the collection, management, analysis, or interpretation of the data; the preparation, review, or approval of the manuscript; or the decision to submit the manuscript for publication.

References

1. Creamer MR, Wang TW, Babb S, et al. Tobacco product use and cessation indicators among adults - United States, 2018. *MMWR Morb Mortal Wkly Rep.* 2019;68(45):1013-1019. doi:10.15585/mmwr.mm6845a2
2. Kasza KA, Ambrose BK, Conway KP, et al. Tobacco-product use by adults and youths in the United States in 2013 and 2014. *N Engl J Med.* 2017;376(4):342-353. doi:10.1056/NEJMsa1607538
3. Kasza KA, Edwards KC, Tang Z, et al. Correlates of tobacco product initiation among youth and adults in the USA: findings from the PATH Study Waves 1-3 (2013-2016). *Tob Control.* 2020;29(Suppl 3):S191-S202. doi:10.1136/tobaccocontrol-2020-055671
4. Wang Y, Sung HY, Yao T, et al. Health care utilization and expenditures attributable to cigar smoking among US adults, 2000-2015. *Public Health Rep.* 2018;133(3):329-337. doi:10.1177/0033354918769873
5. Merianos AL, Jandarov RA, Choi K, Mahabee-Gittens EM. Tobacco smoke exposure disparities persist in U.S. children: NHANES 1999-2014. *Prev Med.* 2019;123:138-142. doi:10.1016/j.ypmed.2019.03.028
6. Kong AY, Queen TL, Golden SD, Ribisl KM. Neighborhood disparities in the availability, advertising, promotion, and youth appeal of little cigars and cigarillos, United States, 2015. *Nicotine Tob Res.* 2020;22(12):2170-2177. doi:10.1093/ntr/ntaa005
7. Jacob P, 3rd, Benowitz NL, Destaillats H, et al. Thirdhand smoke: new evidence, challenges, and future Directions. *Chem Res Toxicol.* 2017;30(1):270-294. doi:10.1021/acs.chemrestox.6b00343
8. Mahabee-Gittens EM, Merianos AL, Matt GE. Preliminary evidence that high levels of nicotine on children's hands may contribute to overall tobacco smoke exposure. *Tob Control.* 2018;27(2):217-219. doi:10.1136/tobaccocontrol-2016-053602
9. Benowitz NL. Cotinine as a biomarker of environmental tobacco smoke exposure. *Epidemiol Rev.* 1996;18(2):188-204. doi:10.1093/oxfordjournals.epirev.a017925
10. Mahabee-Gittens EM, Merianos AL, Hoh E, et al. Nicotine on children's hands: limited protection of smoking bans and initial clinical findings. *Tob Use Insights.* 2019;12:1179173X18823493. doi:10.1177/1179173X18823493
11. Mahabee-Gittens EM, Mazzella MJ, Doucette JT, et al.

- comparison of liquid chromatography mass spectrometry and enzyme-linked immunosorbent assay methods to measure salivary cotinine levels in ill children. *Int J Environ Res Public Health*. 2020;17(4):1157. doi:10.3390/ijerph17041157
12. Mahabee-Gittens EM, Merianos AL, Stone L, et al. Tobacco use behaviors and perceptions of parental smokers in the emergency department setting. *Tob Use Insights*. 2019;12:1179173X19841392. doi:10.1177/1179173X19841392
 13. Mills LM, Semple SE, Wilson IS, et al. Factors influencing exposure to secondhand smoke in preschool children living with smoking mothers. *Nicotine Tob Res*. 2012;14(12):1435-1444. doi:10.1093/ntr/nts074
 14. Koszowski B, Thanner MH, Pickworth WB, et al. Nicotine content and physical properties of large cigars and cigarillos in the United States. *Nicotine Tob Res*. 2018;20(3):393-398. doi:10.1093/ntr/ntx054
 15. Chang CM, Rostron BL, Chang JT, et al. Biomarkers of exposure among U.S. adult cigar smokers: Population Assessment of Tobacco and Health (PATH) Study Wave 1 (2013-2014). *Cancer Epidemiol Biomarkers Prev*. 2019;28(5):943-953. doi:10.1158/1055-9965.EPI-18-0539
 16. Smith DM, O'Connor R J, Wei B, et al. Nicotine and toxicant exposure among concurrent users (co-users) of tobacco and cannabis. *Nicotine Tob Res*. 2020;22(8):1354-1363. doi:10.1093/ntr/ntz122
 17. Mahabee-Gittens EM, Ammerman RT, Khoury JC, et al. Healthy families: study protocol for a randomized controlled trial of a screening, brief intervention, and referral to treatment intervention for caregivers to reduce secondhand smoke exposure among pediatric emergency patients. *BMC Public Health*. 2017;17(1):374. doi:10.1186/s12889-017-4278-8
 18. Quintana PJ, Matt GE, Chatfield D, et al. Wipe sampling for nicotine as a marker of thirdhand tobacco smoke contamination on surfaces in homes, cars, and hotels. *Nicotine Tob Res*. 2013;15(9):1555-1563. doi:10.1093/ntr/ntt014
 19. Wang L, Bernert JT, Benowitz NL, et al. Collaborative method performance study of the measurement of nicotine, its metabolites, and total nicotine equivalents in human urine. *Cancer Epidemiol Biomarkers Prev*. 2018;27(9):1083-1090. doi:10.1158/1055-9965.EPI-17-1127.
 20. Kelley ST, Liu W, Quintana PJE, et al. Altered microbiomes in thirdhand smoke-exposed children and their home environments. *Pediatr Res*. 2021 Mar 2. doi:10.1038/s41390-021-01400-1 Epub ahead of print. PMID: 33654287.
 21. Xue J, Zartarian V, Moya J, et al. A meta-analysis of children's hand-to-mouth frequency data for estimating non-dietary ingestion exposure. *Risk Anal*. 2007;27(2):411-420. doi:10.1111/j.1539-6924.2007.00893.x
 22. Collins BN, Nair US, DiSantis KI, et al. Long-term results from the FRESH RCT: sustained reduction of children's tobacco smoke exposure. *Am J Prev Med*. 2020;58(1):21-30. doi:10.1016/j.amepre.2019.08.021
 23. Matt GE, Quintana PJE, Hoh E, et al. Persistent tobacco smoke residue in multiunit housing: legacy of permissive indoor smoking policies and challenges in the implementation of smoking bans. *Prev Med Rep*. 2020;18:101088. doi:10.1016/j.pmedr.2020.101088
 24. Hood NE, Ferketich AK, Klein EG, et al. Associations between self-reported in-home smoking behaviours and surface nicotine concentrations in multiunit subsidised housing. *Tob Control*. 2014;23(1):27-32. doi:10.1136/tobaccocontrol-2012-050666
 25. Zhang X, Martinez-Donate A, Rhoads N. Parental practices and attitudes related to smoke-free rules in homes, cars, and outdoor playgrounds in US households with underage children and smokers, 2010-2011. *Prev Chronic Dis*. 2015;12:E96. doi:10.5888/pcd12.140553
 26. Kong G, Creamer MR, Simon P, et al. Systematic review of cigars, cigarillos, and little cigars among adolescents: setting research agenda to inform tobacco control policy. *Addict Behav*. 2019;96:192-197. doi:10.1016/j.addbeh.2019.04.032
 27. Agaku I, Odani S, Vardavas C, Neff L. Self-identified tobacco use and harm perceptions among US youth. *Pediatrics*. 2018;141(4). doi:10.1542/peds.2017-3523
 28. Delnevo CD, Hrywna M, Giovenco DP, et al. Close, but no cigar: certain cigars are pseudo-cigarettes designed to evade regulation. *Tob Control*. 2017;26(3):349-354. doi:10.1136/tobaccocontrol-2016-052935
 29. Curwin BD, Hein MJ, Sanderson WT, et al. Nicotine exposure and decontamination on tobacco harvesters' hands. *Ann Occup Hyg*. 2005;49(5):407-413. doi:10.1093/annhyg/meh112
 30. Sheu R, Stonner C, Ditto JC, et al. Human transport of thirdhand tobacco smoke: a prominent source of hazardous air pollutants into indoor nonsmoking environments. *Sci Adv*. 2020;6(10):eaay4109. doi:10.1126/sciadv.aay4109
 31. Invernizzi G, Ruprecht A, De Marco C, et al. Residual tobacco smoke: measurement of its washout time in the lung and of its contribution to environmental tobacco smoke. *Tob Control*. 2007;16(1):29-33. doi:10.1136/tc.2006.017020
 32. Matt GE, Bernert JT, Hovell MF. Measuring secondhand smoke exposure in children: an ecological measurement approach. *J Pediatr Psychol*. 2008;33(2):156-175. doi:10.1093/jpepsy/jsm123
 33. US Food and Drug Administration. Retailers: Chart of Required Warning Statements on Tobacco Product Packaging and Advertising. <https://www.fda.gov/tobacco-products/retail-sales-tobacco-products/retailers-chart-required-warning-statements-tobacco-product-packaging-and-advertising>. Published August 13, 2018. Accessed November 24, 2020.
 34. King JL, Shan L, Azagba S. Association between purchasing behaviors and cigar use: a longitudinal analysis of Waves 1-3 of the Population Assessment of Tobacco and Health (PATH) Study. *PLoS One*. 2020;15(6):e0235496. doi:10.1371/journal.pone.0235496

Copyright of Tobacco Regulatory Science is the property of Tobacco Regulatory Science Group and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.