

# UC San Diego

## UC San Diego Previously Published Works

### Title

Adipose tissue from subjects with type 2 diabetes exhibits impaired capillary formation in response to GRO $\alpha$ : involvement of MMPs-2 and -9

### Permalink

<https://escholarship.org/uc/item/9hj1z37r>

### Journal

Adipocyte, 11(1)

### ISSN

2162-3945

### Authors

Levy, Yifat Amir  
Ciaraldi, Theodore P  
Mudaliar, Sunder R  
[et al.](#)

### Publication Date

2022-12-31

### DOI

10.1080/21623945.2022.2070949

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed

## Adipose tissue from subjects with type 2 diabetes exhibits impaired capillary formation in response to GRO $\alpha$ : involvement of MMPs-2 and -9

Yifat Amir Levy<sup>a,b</sup>, Theodore P Ciaraldi<sup>a,b</sup>, Sunder R. Mudaliar<sup>a,b</sup>, Susan A. Phillips<sup>a,c</sup>, and Robert R. Henry<sup>a,b,\*</sup>

<sup>a</sup>Center for Metabolic Research, Veterans Affairs San Diego Healthcare System, San Diego, CA, USA; <sup>b</sup>Departments of Medicine, University of California, La Jolla, CA, USA; <sup>c</sup>Departments of Pediatrics, University of California, La Jolla, CA, USA

### ABSTRACT

Type 2 Diabetes (T2D) is associated with impaired vascularization of adipose tissue (AT). IL8, GRO $\alpha$  and IL15 are pro-angiogenic myokines, secreted at elevated levels by T2D myotubes. We explored the direct impact of these myokines on AT vascularization. AT explants from subjects with T2D and without diabetes (non-diabetic, ND) were treated with rIL8, rGRO $\alpha$  and rIL15 in concentrations equal to those in conditioned media (CM) from T2D and ND myotubes, and sprout formation evaluated. Endothelial cells (EC) were isolated from T2D and ND-AT, treated with rGRO $\alpha$  and tube formation evaluated. Finally, we investigated the involvement of MMP-2 and -9 in vascularization. ND and T2D concentrations of IL8 or IL15 caused similar stimulation of sprout formation in ND- and T2D-AT. GRO $\alpha$  exerted a similar effect in ND-AT. When T2D-AT explants were exposed to GRO $\alpha$ , sprout formation in response to T2D concentrations was reduced compared to ND. Exposure of EC from T2D-AT to GRO $\alpha$  at T2D concentrations resulted in reduced tube formation. Reduced responses to GRO $\alpha$  in T2D-AT and EC were also seen for secretion of MMP-2 and -9. The data indicate that skeletal muscle can potentially regulate AT vascularization, with T2D-AT having impairments in sensitivity to GRO $\alpha$ , while responding normally to IL8 and IL15.

### ARTICLE HISTORY

Received 24 December 2021  
Revised 19 April 2022  
Accepted 22 April 2022

### KEYWORDS

Type 2 diabetes;  
angiogenesis; adipose tissue;  
myokines; MMPs; groa

## Introduction



Obesity, pre-diabetes, and Type 2 Diabetes (T2D) lie along a continuum of metabolic disease. A common feature is insulin resistance in peripheral tissues, primarily skeletal muscle, adipose tissue, and liver [1].


Capillaries play a crucial role mediating the supply of oxygen, glucose and other substrates, hormones, and inflammatory cells, from the circulation to adipose tissue (AT) [2]. Inappropriate capillarization of AT may limit the delivery of the above factors, lead to relative hypoxia and a subsequent inflammatory response that exacerbates metabolic dysfunction [3,4]. The importance of altered vascularization in insulin-resistant states is supported by the observations that capillary density (CD) is reduced in AT from obese individuals [3,5]. Two major factors determine the vascular phenotypes of AT. One is the intrinsic character of local endothelial cells (EC) [6,7], and another is the microenvironment of pro- and anti-angiogenic factors within each tissue [4,8,9].

Over the past two decades, evidence has accumulated that skeletal muscle (SkM) is a secretory tissue that releases

a variety of myokines which can act in autocrine, paracrine and/or endocrine fashions to modulate multiple processes in other tissues (reviewed in [10]). While some of the secreted factors are unique to SkM, many are broadly produced and secreted cyto- and chemokines and growth factors. Among them are IL8, GRO $\alpha$  and IL15, which while acting as modulators of inflammation [11], are known also to regulate angiogenesis [11–13]. Our laboratory and others have reported that myotubes from human subjects with T2D secrete higher levels of IL8, GRO and IL15 compared to ND myotubes [14,15]. In addition, we reported that the elevated levels of IL8 secreted from T2D myotubes impaired angiogenesis in both human umbilical vein endothelial cells (HUVEC) and human skeletal muscle explants when compared to levels from ND myotubes [16].

In the current report, we investigate the potential effects of the elevated levels of IL8, GRO $\alpha$  and IL15 secreted from T2D SkM on capillarization of AT from ND and T2D subjects. As a surrogate for angiogenesis in AT, we employed adipose tissue explants, which can form endothelial sprouts, and EC isolated from AT. Understanding the roles of IL8, GRO $\alpha$ , and IL15 in regulating angiogenesis and the signalling pathways

**CONTACT** Theodore P Ciaraldi  [tciaraldi@ucsd.edu](mailto:tciaraldi@ucsd.edu)  VA San Diego Healthcare System, 3350 La Jolla Village Drive (111G), San Diego, CA 92161, \*Deceased

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/21623945.2022.2070949>.

This work was authored as part of the Contributor's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

This is an Open Access article that has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights (<https://creativecommons.org/publicdomain/mark/1.0/>). You can copy, modify, distribute and perform the work, even for commercial purposes, all without asking permission.

involved could shed light on their ability to modulate vascularization and the metabolic function of AT in T2D.

## Results

### Effects of exogenous myokines on capillary length in adipose tissue explants


To explore whether higher levels of pro-angiogenic factors IL8, IL15 and GRO $\alpha$  could be involved in altered capillary sprout formation in AT, rIL8, rGRO $\alpha$  and rIL15, either alone or in combination, were added to ND-AT and T2D-AT explants in concentrations equal to the averages of those found in CM from T2D and ND myotubes ([T2D-IL8] = 3280 and [ND-IL8] = 2071 pg/mL; [GRO $\alpha$ ] = 4620 and 1859 pg/mL; and [IL15] = 1.07 and 0.6 pg/mL, respectively) [14]. Capillary outgrowth was measured after 4–6 days. Capillary outgrowth was stimulated to a similar extent in ND-AT and in T2D-AT by the rIL8 + rGRO $\alpha$  + rIL15 combination at ND and T2D concentrations (Supplementary Material, Figure 1a, b). The same was true after exposure to rIL15 alone; no difference between groups or concentrations (SM, Figure 1c, d). Regarding rIL8, after 5 days of treatment, capillary

outgrowth in ND-AT was modestly, but significantly ( $p < 0.05$ ), lower in response to the T2D concentration of IL8 compared to ND-IL8. This difference was lost by day 6 of treatment. In T2D-AT, the response was the same to both concentrations of IL8 (SM, Figure 1e, f).

Finally, when exposed to rGRO $\alpha$  alone, ND-AT responded the same to both levels of rGRO $\alpha$  (Figure 1a). However, T2D-AT displayed a differential sensitivity to rGRO $\alpha$ , with treatment by higher (T2D) concentrations suppressing capillary outgrowth compared to treatment at the lower ND-CM level (Figure 1b) ( $p < 0.005$  at each day 4–6). Figure 1c presents representative images of capillary outgrowth from T2D-AT explants after exposure to ND- (Figure 1c-a) and T2D-CM (Figure 1c-b) concentrations of rGRO $\alpha$ . These results suggest that higher levels of GRO $\alpha$  such as those secreted from T2D myotubes might contribute to the lower capillarization observed in T2D-AT [5].

### Effects of exogenous GRO $\alpha$ on tube formation by endothelial cells isolated from ND- and T2D-AT

Adipose tissue explants represent a mixed system, where multiple cell types may contribute pro- and anti-angiogenic factors to the microenvironment in addition to those provided by myotubes. To understand more about the direct effects of myotube-produced levels of



**Figure 1.** Effects of ND- and T2D [GRO $\alpha$ ] on capillary outgrowth from ND- and T2D-AT.

A & B. Quantification of capillary length from AT explants from ND (A) or T2D (B) subjects after incubation with rGRO $\alpha$  in concentrations equivalent to ND- (solid bars) or T2D (open bars) myotube CM for 4d – 6d. Average + SD, ND-AT,  $n = 14$  subjects and T2D-AT  $n = 15$ . Shown is the average capillary length/ $500\mu\text{m}^2$  for three fields of at least three different explants/subject. \*\*\* $p < 0.005$  T2D-rGRO $\alpha$  vs. ND- rGRO $\alpha$ . C. Representative images of adipose tissue explants from T2D-AT after incubation with rGRO $\alpha$  in concentrations equivalent to ND- (a) or T2D-CM (b).


GRO $\alpha$  on AT angiogenesis, endothelial cells were isolated from AT of ND (ND-AT-EC) and T2D (T2D-AT-EC) subjects. These AT-EC display characteristic endothelial markers Von Willebrand Factor and CD31 (SM, Figure 2b & 2d) and can form tube-like structures when stimulated, like established endothelial cell lines (SM, Figure 2e) [17].

Like the procedure with AT explants, AT-EC were exposed to concentrations of rGRO $\alpha$  equal to those present in media conditioned by ND- and T2D-CM myotubes. Since the readout in this assay is the formation of tube-like structures by existing cells, rather than capillary outgrowth, treatment times could be shortened (to 8 hr rather than days). When treated with rGRO $\alpha$ , the response of ND-AT-EC was very similar to that of ND-AT, with equal stimulation of the angiogenic response at both concentrations (Fig. 2Aa & b and B top panel). However, T2D-AT-EC behaved in the same manner as T2D-AT, where the higher concentration of GRO $\alpha$  suppressed tube formation, compared to the ND level (Figure 2 a c) & (d and b bottom panel) ( $p < 0.005$ ).

### **MMP-2 & MMP-9 and GRO $\alpha$ in regulation of angiogenesis in T2D-AT and T2D-AT-EC**

A critical step in the angiogenic process is the release of matrix metalloproteinases (MMPs) from endothelial cells, with MMP-2 and -9 playing important roles in supporting and promoting angiogenesis [18]. To identify possible molecular mechanisms for the dose-dependency of the T2D-AT angiogenic response to GRO $\alpha$ , T2D-AT explants were treated with [ND-GRO $\alpha$ ] and [T2D-GRO $\alpha$ ] for 4d and the levels of MMP-2 and MMP-9 released into the media measured. Exposure to the higher concentration of GRO $\alpha$ , [T2D-GRO $\alpha$ ], resulted in significantly less MMP-2 and MMP-9 in the media (Figure 3 a and b) compared to the response to [ND-GRO $\alpha$ ] ( $p < 0.01$ ).

Similar responses were detected when T2D-AT-EC were treated with [ND-GRO $\alpha$ ] and [T2D-GRO $\alpha$ ] for 8 h and the levels of MMP-2 and MMP-9 released into the media measured. Exposure to the higher concentration of GRO $\alpha$  ([T2D-GRO $\alpha$ ]) resulted in significantly less MMP-2 and MMP-9 released into the media (Figure 3 a and c) compared to the response to [ND-GRO $\alpha$ ] ( $p < 0.01$ ).



**Figure 2.** Effects of ND- and T2D-[rGRO $\alpha$ ] on capillary tube formation by endothelial cells isolated from ND- and T2D-AT. A. Representative images of tube formation of endothelial cells isolated from adipose tissue explants from ND-AT (a, b) or T2D-AT (c, d) after incubation with rGRO $\alpha$  in concentrations equivalent to ND-CM (a, c) or T2D-CM [rGRO $\alpha$ ] (b, d).

B. Quantification of tube formation of endothelial cells isolated from adipose tissue explants from ND (top) or T2D (bottom) subjects after incubation with rGRO $\alpha$  in concentrations equivalent to ND- rGRO $\alpha$  (solid bars) or T2D- rGRO $\alpha$  (open bars). Average + SD, ND-AT-EC,  $n = 4$  subjects and T2D-AT-EC  $n = 6$ . Shown is the average capillary number/500 $\mu\text{m}^2$  for 4 fields of at least two independent experiments for each subject. \*\*\* $p < 0.005$  T2D-rGRO $\alpha$  vs. ND-rGRO $\alpha$ .


Addition of the MMP-2/MMP-9 inhibitor BiPS (25  $\mu$ M) reduced tube formation in response to both [ND- GRO $\alpha$ ] ( $p < 0.0001$ ) and [T2D- GRO $\alpha$ ] ( $p < 0.05$ ) in T2D-AT-EC. However, we found that the inhibition of tube formation by BiPS was greater in T2D-AT-EC treated with [ND-GRO $\alpha$ ] compared to [T2D- GRO $\alpha$ ] ( $p < 0.01$ ) (Figure 4).

## Discussion


Adipose tissue is perhaps the most plastic tissue in the body, adjusting its mass in response to changing patterns of substrate availability, environmental stresses, neuro-hormonal stimuli, or pathological conditions. A key feature of adipose tissue remodelling is the development of an appropriate vascular network. Indeed, evidence suggests that the development of the vascular supply is a rate-limiting step in AT expansion [19–21]. Failure to adequately accomplish this goal can lead to several consequences, including reduced exposure to hormones and substrates, hypoxia, increased tissue inflammation, and, ultimately, insulin resistance [3,4]. The importance of adequate AT blood flow (ATBF) is emphasized by the observation that AT CD is reduced in obesity and insulin resistance states [3,5] and

impairments in ATBF may be apparent before insulin resistance [4,22]. In adult AT, expansion of the vascular network occurs through the multistep process of angiogenesis [23]. Major determinants of the efficiency of AT angiogenesis are the presence of pro- and anti-angiogenic factors in the AT microenvironment [4,8,9,24] and the performance of resident endothelial cells [6,7].

In addition to the crucial mechanical and metabolic functions of skeletal muscle, it has become apparent over the past 20 years that muscle can also serve as a secretory tissue, releasing protein factors from myotubes termed myokines that can act in autocrine, paracrine, and endocrine fashions to regulate multiple processes (reviewed in [12,25]). A number of these myokines, such as VEGF, IL8, IL15 and GRO $\alpha$ , are known to exert pro-angiogenic activities. Multiple investigators have shown that the secretion of selected myokines can be regulated by the presence of T2D (reviewed in [12,25]). We reported that myotubes from T2D subjects, compared to ND myotubes, release elevated amounts of multiple myokines, including IL8, IL15 and GRO, while VEGF secretion was similar between groups [11,14]. Indeed, IL8 and GRO are two of the most abundantly secreted myokines [11,14].



**Figure 3.** Effects of rGRO $\alpha$  on secretion of MMP-2 and MMP-9 from T2D-AT explants and endothelial cells isolated from T2D-AT. rGRO $\alpha$  in concentrations equivalent to ND- or T2D- GRO $\alpha$  was added to T2D-AT and T2D-AT-EC. Culture media was collected from T2D-AT after 4d and from T2D-AT-EC after 8 h. A. Representative western-blot of MMP-2 and MMP-9 from T2D-AT and T2D-AT-EC. B & C. Quantification of blots for MMP-9 and MMP-2 content in the media of T2D-AT (b) and T2D-AT-EC (c); ND-rGRO $\alpha$  (solid bars) and T2D-rGRO $\alpha$  (open bars). Results are the average + SD of 2 independent experiments for each subject: T2D-AT  $n = 6$ , and T2D-AT-EC  $n = 6$ . \* $p < 0.05$ , T2D-rGRO $\alpha$  vs. ND-rGRO $\alpha$ .



**Figure 4.** Effects of MMP-2/MMP-9 inhibition with BiPS on capillary tube formation of T2D-AT-EC induced by rGRO $\alpha$ . A. Representative images of tube formation of T2D-AT-EC after incubation with rGRO $\alpha$  in concentrations equivalent to ND- rGRO $\alpha$  (a, b) or T2D- rGRO $\alpha$  (c, d), with 0 (a, c) or 25  $\mu$ M BiPS (b, d), for 8 h. B. Quantification of tube formation of T2D-AT-EC after incubation with rGRO $\alpha$  in concentrations equivalent to ND- rGRO $\alpha$  (top) or T2D- rGRO $\alpha$  (bottom) with 0 (solid bars) or 25  $\mu$ M BiPS (open bars). Average  $\pm$  SD, T2D-AT-EC  $n = 3$  subjects. Shown is the average capillary number/500 $\mu$ m<sup>2</sup> for 3 fields of at least 2 independent experiments for each subject. ++++  $p < 0.005$  ND-rGRO $\alpha$  with 25  $\mu$ M BiPS vs. ND-rGRO $\alpha$  without BiPS, +  $p < 0.05$  T2D-rGRO $\alpha$  with 25  $\mu$ M BiPS vs. T2D-rGRO $\alpha$  without BiPS.

Thus, it is possible that these specific myokines could, by acting in a paracrine manner, contribute to the regulation of angiogenesis in skeletal muscle. Yet, while T2D myotubes release elevated levels of these pro-angiogenic factors, skeletal muscle tissue in T2D is characterized by a reduced CD [26], like that observed in AT [21]. Previously we tested possible explanations for this paradox, using the HUVEC line as a surrogate for muscle endothelial cells. Exposure of HUVEC to media conditioned by ND (ND-CM) and T2D (T2D-CM) myotubes, resulted in stimulation of capillary tube formation. However, the response to T2D-CM was considerably less than that to ND-CM [17]. As IL8, IL15 and GRO were all present at higher levels in T2D-CM, we examined the effects of these factors, applied individually or in combination at concentrations matching the average values in ND- and T2D-CM. Multiple lines of evidence revealed that it was IL8 that was responsible for the biphasic dose-dependent effect on capillary tube formation by HUVEC [16]. Thus, it is possible that the lower CD seen in T2D SkM could be due, at least in part, to the

higher levels of IL8 present in the microenvironment of diabetic muscle.

Given the high levels of IL8 and GRO $\alpha$  secreted by myotubes and the mass of skeletal muscle, it is possible that delivery of these myokines into the circulation could contribute in an endocrine manner to regulation of angiogenesis in AT as well. This postulate was tested in the current work. Important differences from our previous work include the fact that the present studies include an additional variable, the identity of the response system, AT from ND and T2D subjects, and not a homogenous system such as HUVEC. Unlike the situation with HUVEC, there was not a dose-dependent angiogenic response by AT, in this instance capillary outgrowth, to the IL8+ IL15+ GRO $\alpha$  combination. Both HUVEC and AT responded similarly to ND- and T2D equivalent levels of IL15 alone [16]. The most striking differences between HUVEC and skeletal muscle explants and AT involved responses to IL8 and GRO $\alpha$ . While T2D related levels of IL8 reduced angiogenesis in HUVEC and skeletal muscle explants, it had very modest transient effects in AT. Distinctly in AT it

was the T2D-related higher level of GRO $\alpha$  that blunted angiogenesis. Furthermore, it is only T2D-AT and T2D-AT-EC that displayed a differential sensitivity to GRO $\alpha$ .

Mixed effects of GRO $\alpha$  on cell migration and angiogenesis have been observed in multiple systems. In HUVEC, GRO $\alpha$  primarily stimulates cytoskeletal reorganization [27] and cell migration and tube formation [28] in a dose-dependent manner, with a tendency for responses to be blunted at higher concentrations. However, in mast cells [29] and human airway smooth muscle cells [30] GRO $\alpha$  inhibits cell migration. Little is known about possible roles of GRO $\alpha$  in T2D [31]. Circulating levels are reported to be elevated in obese [32] and T2D subjects [33] by some, but not all [34], investigators. GRO $\alpha$  is synthesized and secreted equally by the adipocyte and stromal-vascular fractions of omental adipose tissue (OAT) [35]. Such secretion is higher in tissue from obese individuals [35], yet the absolute amount of GRO $\alpha$  secreted from OAT appears to much less than that produced by myotubes [14,15,36,37]. Considering the relative masses of OAT and SkM in humans, it is possible that myotube-derived GRO $\alpha$  contributes a major portion of the GRO $\alpha$  present in the circulation and within tissues.

GRO $\alpha$  and IL8 share several characteristics, belonging to the same structural class of chemokines (CXC with a Glu-Leu-Arg sequence in the N-terminus) and interacting with the same receptors [25,27]. This similarity gives rise to the question of why in T2D-AT and T2D-AT-EC there is a dose-dependency for angiogenesis in response to GRO $\alpha$ , but not IL8, while in HUVEC the opposite is true [16]. Cell type specificity in the preference between GRO $\alpha$  and IL8 is not without precedent, as in human lung microvascular endothelial cells, IL8 is considerably more potent than GRO $\alpha$  in stimulating actin reorganization [27]. Further support for a 'functional distinction' [38] between GRO $\alpha$  and IL8 is provided by studies in rabbit knee joints, where IL8 is more potent for neutrophil recruitment, while it is GRO $\alpha$  that induces TNF $\alpha$  expression [38].

Both our previous work [16] and the current findings show the varying effects of the microenvironments created by ND and T2D myokines on capillary tube formation. What the current work adds is information about the tissue selectivity of responses to specific myokines. Since AT is a highly heterogeneous system, we isolated endothelial cells from AT to determine if they were responsible for the response to GRO $\alpha$  specific to T2D. We found that differential sensitivity to GRO $\alpha$  is an intrinsic property of T2D-EC. Altered behaviour of T2D-EC would agree with multiple *in vitro* studies showing that endothelial progenitor cells (EPC) from individuals with T2D generated EC

with reduced proliferation, adhesion, and capacity to form tubes, thereby inhibiting their ability to revascularize damaged tissue [7]. Additionally, such behaviour is observed in EC isolated from db/db mice [39] and can also be generated *in vitro* by culture of EC under hyperglycaemic conditions [40]. Furthermore, Gealekman and colleagues have reported that the angiogenic capacity and CD of SAT is decreased with morbid obesity [3]. The ND and T2D subjects studied here display a similar extent of adiposity, reducing the impact of that factor.

Angiogenesis is a multistep process that requires EC activation, migration, proliferation, and tube formation [41]. To allow EC migration, degradation of the extracellular matrix (ECM) surrounding the EC is required, mediated by matrix metalloproteinases (MMPs) released from EC. The reduced release of MMP-2 and -9 from T2D-AT and T2D-AT-EC after exposure to [T2D-GRO $\alpha$ ] (Figure 3) could contribute at an early step to the impaired tube formation observed. This behaviour exclusive to T2D-EC would be consistent with reports that the presence of insulin resistance is associated with a relative reduction in MMP-9 gene expression in AT [42]. In addition, MMPs have recently been shown to act on several substrates in addition to ECM proteins [41]. One of these is GRO $\alpha$ , which is inactivated by MMP-9. Thus, reduced MMP-9 release by T2D-AT-EC in response to T2D-related levels of GRO $\alpha$  would lead to reduced GRO $\alpha$  cleavage, elevating GRO $\alpha$  levels even further.

Our results reinforce the concept that T2D can influence ATBF in several ways. One is the well-documented resistance of EC to insulin-induced vasodilation [43]. Another is elevated secretion of GRO $\alpha$ , to levels that could impair angiogenesis in AT. This would result in a reduced vascular reserve leading to reduced delivery of insulin, substrates, and oxygen to adipocytes, leading to metabolic dysfunction and hypoxia, the later triggering recruitment of inflammatory cells [4,21,43]. This could induce adipocyte apoptosis, with further release of inflammatory mediators. The importance of appropriate ATBF to metabolic health is highlighted by numerous reports demonstrating associations between reduced ATBF or CD and insulin resistance, as well as the development of T2D [4,21,43].

Several aspects of this study need to be considered about their impact on the physiologic relevance of our findings. One is a potential limitation of studies with primary cells or cell lines, including the current work, such that they may not be fully reflective of the behaviour of that specific cell type in the context of the intact tissue. Our findings with explant AT in culture confirm the results with EC in a more physiologic system and are consistent with the reduced ATCD and ATBF seen *in vivo* in insulin-resistant humans

and rodents. Second, is that our observations about myokine effects on AT angiogenesis are limited to the factors studied, GRO $\alpha$ , IL8 and IL15, and do not include the primary pro-angiogenic agent, VEGF. We chose to focus on myokines whose release differed between ND and T2D myotubes; unlike with VEGF [11,14]. Of note, VEGF release has been reported to be higher from AT of subjects with T2D [44], suggesting that the microenvironment of T2D AT could include elevated levels of VEGF. Potential T2D-related differential sensitivity to the angiogenic actions of VEGF represents an exciting area for further study. Circulating (serum) GRO $\alpha$  levels in healthy humans have been reported to range over 50–90 pg/mL [32], while levels in individuals with T2D display levels ~50% higher [33]. However, this observation is not universal [34]. Also, there is limited information about absolute levels in skeletal muscle or adipose tissue, especially in humans, although it has been reported that isolated macrophages and pre-adipocytes differentiated from human OAT secrete GRO and that such secretion is elevated several-fold in cells from obese individuals or with exposure to inflammatory agents [45], thereby supporting the supposition that T2D AT may be exposed *in vivo* to elevated levels of GRO $\alpha$ .

In summary, the current report reveals an intrinsic property of T2D-AT-EC whereby elevated levels of GRO $\alpha$  reduce the angiogenic response and, ultimately, CD in AT. Our findings suggest that both the overproduction of GRO $\alpha$  and an altered response of T2D-AT-EC to this factor can contribute to the metabolic and vascular dysfunction of AT in T2D. Meanwhile, both ND and T2D-AT respond similarly to IL8 and IL15.

## Patients/methods/materials

### Patients

AT samples were obtained from 14 ND (ND-AT) and 15 T2D (T2D-AT) subjects. Subjects were classified as ND based on values of either HbA1c < 5.7 or 2 hr OGTT [glucose] < 140 mg/dL within 2 months of biopsy. They also had to be negative for a family history (1<sup>st</sup> degree

relative) of T2D. Subjects with T2D were identified based on an existing clinical diagnosis. Anti-diabetic medications used included: metformin, glipizide, glargine and liraglutide, as well as combinations of the same. Medication use was stable for at least 3 months and maintained up to the time of biopsy. General inclusion criteria for all subjects were: weight stable ( $\pm 2$  kg) over the last 2 months and no medication changes over the last 3 months. Potential subjects were excluded if they were taking medications known to influence carbohydrate metabolism, including antidepressants, and the presence of other conditions known to influence carbohydrate metabolism. Female subjects who were not post-menopausal were biopsied during the early follicular phase of their cycles. Characteristics of the subjects are summarized in Table 1. The Committees on Human Investigation of the University of California, San Diego, and VA San Diego Healthcare System approved the experimental protocol. Informed written consent was obtained from all subjects after explanation of the protocol.

### Adipose tissue biopsy

Subcutaneous adipose tissue (SAT) samples ( $n = 20$ ) were obtained from the lateral abdominal wall region (peri-umbilical) after an overnight (10–12 hr) fast [46]. Briefly, lidocaine (1%) was infiltrated in a square field fashion and the biopsy taken, using a 5-mm side-cutting needle, from the centre of the field. For nine subjects, SAT was obtained during elective gastric bypass surgery. In that instance, SAT samples were collected from the periumbilical region intraoperatively during dissection of the usually placed trocar sites for the laparoscopic operation. All subjects were weight stable before tissue collection.

### Ex vivo angiogenesis assay of adipose tissue

The assay was performed employing established protocols [3,5,19]. Briefly, freshly harvested human AT was cut into  $\sim 1$  mm<sup>3</sup> pieces, matched in size for each individual subject, which were then embedded individually in wells of a 24-well plate, in growth-factor-

**Table 1.** Subject Characteristics.

Group	N (F/M)	Age (yr)	BMI (kg/m <sup>2</sup> )	Fasting glucose (mM)	Fasting insulin (pM)	HOMA-IR
ND	14 (4/10)	48 $\pm$ 4	33.6 $\pm$ 1.6	5.28 $\pm$ 0.16	56 $\pm$ 14	1.2 $\pm$ 0.3
T2D	15 (5/10)	47 $\pm$ 3	37.0 $\pm$ 2.4	7.94 $\pm$ 0.86†	172 $\pm$ 68	2.6 $\pm$ 0.6*

\* $p < 0.05$  vs ND, † $p < 0.01$  vs ND



depleted Matrigel (280  $\mu$ L/well) (BD Bioscience) and cultured in 1:1 EGM-2-MV (Lonza) and  $\alpha$ -MEM (Life Technologies/Gibco.). Explants were treated with rIL8 (#208-IL-010/CF), rGRO $\alpha$  (#275-GR-010), and rIL15 (#247-ILB-005) (all recombinant proteins are human low-endotoxin from R&D Systems). Media and treatments were replaced every other day. Routinely, 3 independent explants per subject for each condition were embedded. Each explant was analysed for capillary outgrowth length by light microscopy and images captured with a 4X objective. The length of capillary branches at the periphery of the growth area was quantified by 3 separate blinded observers (for each explant), using the NIH Image-J program [3,5,19]. In control studies, the basal growth with 1:1 EGM-2-MV or  $\alpha$ -MEM without supplements was minor. It has been demonstrated that >90% of sprouting cells from the AT explants express endothelial cell markers, and thus represent angiogenic growth [3,5,19].

### **Endothelial cell isolation and identification**

Endothelial cells were obtained through two different protocols. With AT collected from abdominal surgeries, the stromal-vascular fraction (SVF) was isolated by collagenase digestion, filtration and centrifugation as described previously [47,48]. CD31 positive cells were isolated with CD31-magnetic microbeads using MACS technology (Miltenyl Biotec, #130-091-935) according to the manufacturer's instructions. With this procedure 96.2% of the cells were CD31 + . Alternatively, freshly harvested human adipose tissue was cut into  $\sim$ 1 mm<sup>3</sup> pieces, which were embedded individually in wells of a 24-well plate, in growth-factor-depleted Matrigel (280  $\mu$ L/well) and cultured in DMEM with D-Valine replacing L-Valine and supplemented with 15% FBS (GE Healthcare) [19,49,50]. After 5 days, the explant was removed, and the cells allowed to propagate for an additional 3–5 days. Cells were released from Matrigel with dispase and transferred to EGM-2-MV media. Endothelial cells were authenticated for each individual preparation using antibodies against human Von Willebrand Factor (Dako, catalogue # A0082) and PECAM-1 (CD31) (Millipore, catalogue #04-1074A). AT-EC were not tested for mycoplasma contamination. The samples were stained with Vectastain Universal Elite Kit (Vector Laboratories, #PK-6200) and DAB Substrate Kit (Vector Laboratories, #SK-4100) according to the manufacturer's instructions.

### **Culture of primary isolated endothelial cell and angiogenesis assay**

Isolated EC were grown according to standard protocols in EGM-2-MV [51]. Measurements of capillary-like tube formation by the isolated cells were achieved using an in vitro assay of endothelial cell tube formation [51]. Matrigel was plated onto 96-well plates (50  $\mu$ L/well) and incubated at 37°C for 60 minutes. Cells were cultured in EGM-2-MV in the absence or presence of rGRO $\alpha$ , and then seeded onto Matrigel at 10,000 cells/well for 8 h. Cells were analysed for capillary number by light microscopy and phase contrast images captured on a Nikon TS100 microscope with a 4X objective. The readout for this assay is the formation of capillary-like tubes. A tube is defined as a closed network unit or an intact loop. Results were obtained by counting the number of tubes in a field. The number of capillaries was quantified by 3–4 separate, blinded observers, using NIH Image-J. In control studies, the extent of capillary formation with serum free EGM-2-MV was negligible. Each experiment was repeated at least 3 times in quadruplicate. In some experiments BiPS (Millipore) a specific inhibitor of MMP-2/MMP-9 [52], was added together with other treatments.

### **Media collection and immunoblotting**

Tissue explants and isolated EC were cultured and treated with or without rGRO $\alpha$  in the presence of EGM-2-MV under the same conditions as described above for the respective angiogenesis assays. Conditioned media was collected after 4d and 8 h, respectively, centrifuged for 10 min at 800  $\times$  g to remove cell debris and stored at  $-80^{\circ}$ C before analysis. Proteins were resolved on 10% SDS-PAGE under reducing conditions, transferred to nitrocellulose membranes, blocked with Odyssey block (LI-COR Biosciences) and incubated with antibodies against MMP2 or MMP9 (Cell Signalling, catalogue #4022 and #3852, respectively). Secondary IRDye antibodies were obtained from LI-COR (catalogue #926-68,020 and #926-68,073). Detection and quantification of band intensity was performed using Odyssey Infrared Imaging System and Image Studio analysis software (version 3.1.4).

### **Data analysis**

Statistical analysis was performed using GraphPad Prism 5.0 (GraphPad). Between group comparisons were evaluated by independent group t test if data was normally distributed and with a Mann-Whitney test for non-normally distributed data. Variances were

similar between groups. Within group comparisons (treatment effects) were evaluated by paired t test. For results that were not normally distributed, data was log-transformed for statistical analysis and then back-transformed and reported in original units as mean  $\pm$  SD. Variation within each group of data was unknown at the initiation of studies. Statistical significance was accepted as  $p < 0.05$ . The number of individual determinations for each measurement is indicated in the legends. The number of individual determinations for each measurement was established after initial measurements revealed the variation for that specific measurement and set to detect a difference of 1.5 SD.

## Acknowledgments

We wish to thank Debra Armstrong and Paivi Burke for assistance with subject care and adipose tissue biopsies, Andrea Gasper for performing many of the biopsies, and Leslie Carter for technical assistance. We also wish to thank Dr. Alan Wittgrove and the staff of the Wittgrove Bariatric Center for providing adipose tissue from elective gastric bypass surgeries.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

This work was supported by Merit Review Award #I01CX00635 (to RRH) from the United States (U.S.) Department of Veterans Affairs Clinical Sciences Research and Development Service. The contents do not represent the views of the U.S. Department of Veterans Affairs or the United States Government.

## Author contributions

Y.A.L. designed studies, researched data and co-wrote the manuscript. T.P.C. researched data and co-wrote/edited the manuscript. S.R.M. contributed to discussion and reviewed/edited the manuscript. S.A.P. collected & processed surgical adipose tissue biopsies, contributed to discussion, and reviewed/edited the manuscript. R.R.H. contributed to study design and provided resources. Y.A.L. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## Data Availability Statement

The data that support the findings of this study are available by request of the corresponding author.

## Ethical approval

The Committees on Human Investigation of the University of California, San Diego, and VA San Diego Healthcare System approved the experimental protocol.

## Informed consent

Informed written consent was obtained from all subjects after explanation of the protocol.

## References

- [1] DeFronzo RA. Pathogenesis of type 2 (non-insulin-dependent) diabetes mellitus: a balanced overview. *Diabetologia*. 1992;35:389–397.
- [2] Manneras-Holm L, Krook A. Targeting adipose tissue angiogenesis to enhance insulin sensitivity. *Diabetologia*. 2012;55:2562–2564.
- [3] Gealekman O, Guseva N, Hartigan C, et al. Depot-specific differences and insufficient subcutaneous adipose tissue angiogenesis in human obesity. *Circulation*. 2011;123:186–194.
- [4] Crewe C, An YA, Scherer PE. The ominous triad of adipose tissue dysfunction: inflammation, fibrosis, and impaired angiogenesis. *J Clin Invest*. 2017;127:74–82.
- [5] Gealekman O, Guseva N, Gurav K, et al. Effect of rosiglitazone on capillary density and angiogenesis in adipose tissue of normoglycemic humans in a randomized controlled trial. *Diabetologia*. 2012;55:2794–2799.
- [6] Ding H, Triggle CR. Endothelial cell dysfunction and the vascular complications associated with type 2 diabetes: assessing the health of the endothelium. 2005; *Vasc Health Risk Management*. 1:55–71.
- [7] Tepper OM, Galiano RD, Capla JM, et al. Human endothelial progenitor cells from type II diabetics exhibit impaired proliferation, adhesion and incorporation into vascular structures. *Circulation*. 2002;106:2781–2786.
- [8] Bonner JS, Lantier L, Hasenour CM, et al. Muscle-specific vascular endothelial growth factor deletion induces muscle capillary rarefaction creating muscle insulin resistance. *Diabetes*. 2013;62:572–580.
- [9] Spencer M, Unal R, Zhu B, et al. Adipose tissue extracellular matrix and vascular abnormalities in obesity and insulin resistance. *J Clin Endocrinol Metab*. 2011;96:E1990–8.
- [10] Marcus MS, Wong GW. Regulation of tissue crosstalk by skeletal muscle-derived myonectin and other myokines. *Adipocyte*. 2012;1:200–202.
- [11] Pedersen BK, Febbraio MA. Muscle, exercise and obesity: skeletal muscle as a secretory organ. *Nat Rev Endocrinol*. 2012;8:457–465.
- [12] Eckel J. Myokines in metabolic homeostasis and diabetes. *Diabetologia*. 2019;62:1523–1528.
- [13] Dimberg A. Chemokines in angiogenesis. *Curr Top Microbiol Immunol*. 2010;341:59–80.
- [14] Ciaraldi TP, Ryan AJ, Mudaliar SR, et al. Altered myokine secretion is an intrinsic property of

- skeletal muscle in type 2 diabetes. *PLoS ONE*. 2016;11:e0158209.
- [15] Bouzakri K, Plomgaard P, Berney T, et al. Bimodal effect on pancreatic  $\beta$ -cells of secretory products from normal or insulin-resistant human skeletal muscle. *Diabetes*. 2011;60:1111–1121.
  - [16] Amir-Levy Y, Ciaraldi TP, Mudaliar SR, et al. Excessive secretion of IL-8 by skeletal muscle in type 2 diabetes impairs tube growth: potential role of PI3K and the Tie2 receptor. *Am J Physiol Endocrinol Metab*. 2015;309:E22–E34.
  - [17] Scott E, Bicknell R. The isolation and culture of microvascular endothelium. *J Cell Sci*. 1993;105:269–273.
  - [18] Bellafiore M, Battaglia G, Bianco A, et al. The involvement of MMP-2 and MMP-9 in heart exercise-related angiogenesis. *J Trans Med*. 2013;11:283.
  - [19] Rojas-Rodriguez R, Gealekman O, Kruse ME, et al. Adipose tissue angiogenesis assay. *Methods Enzymol*. 2014;537:75–91.
  - [20] Pasarica M, Sereda OR, Redman LM, et al. Reduced adipose tissue oxygenation in human obesity: evidence for rarefaction, macrophage chemotaxis, and inflammation without an angiogenic response. *Diabetes*. 2009;58:718–725.
  - [21] Gealekman O, Corvera S. Adipose tissue angiogenesis: impact on obesity and type 2 diabetes. *Biochim Biophys Acta*. 2014;1842:463–472.
  - [22] Goossens GH, Bizzarri A, Ventclef N, et al. Increased adipose tissue oxygen tension in obese compared with lean men is accompanied by insulin resistance, impaired adipose tissue capillarization, and inflammation. *Circulation*. 2011;124:67–76.
  - [23] Senger DR, Davis GE. Angiogenesis. *Cold Spring Harb Perspect Biol*. 2011;3:1–19.
  - [24] van Harmelen V, Skurk T, Rohrig K, et al. Effect of BMI and age on adipose tissue cellularity and differentiation capacity in women. *Int J Obes Relat Metab Disord*. 2003;27(8):889–895.
  - [25] Severinsen MCK, Pedersen BK. Muscle-Organ crosstalk: the emerging roles of myokines. *Endocr Rev*. 2020;41:594–609.
  - [26] Lillioja S, Young AA, Cutler CL, et al. Skeletal muscle capillary density and fiber type are possible determinants of in vivo insulin resistance in man. *J Clin Invest*. 1987;80(2):415–424.
  - [27] Schraufstatter IU, Chung J, Burger M. IL-8 activates endothelial cell CXCR1 and CXCR2 through rho and rac signaling pathways. *Am J Physiol Lung Cell Mol Physiol*. 2001;280:L1094–103.
  - [28] Wang D, Wang H, Brown JA, et al. CXCL1 induced by prostaglandin E2 promotes angiogenesis in colorectal cancer. *J Exp Med*. 2006;203:941–951.
  - [29] Alkhoury H, Moir LM, Armour JCL, et al. CXCL1 is a negative regulator of mast cell chemotaxis to airway smooth muscle cell products in vitro. *Clin Exp Allergy*. 44. 2014:381–392.
  - [30] Al-Alwan LA, Chang Y, Rousseau S, et al. CXCL1 inhibits airway smooth muscle cell migration through the decoy receptor duffy antigen receptor for chemokines. *J Immunol*. 2014;193:1416–1426.
  - [31] Pedersen L, Olsen CH, Pedersen BK, et al. Muscle-derived expression of the chemokine CXCL1 attenuates diet-induced obesity and improves fatty acid oxidation in muscle. *Am J Physiol Endocrinol Metab*. 2012;302:E831–E40.
  - [32] Nunemaker CS, Chung HG, Verrilli GM, et al. Increased serum CXCL1 and CXCL5 are linked to obesity, hyperglycemia, and impaired islet function. *J Endocrinol*. 2014;222:267–276.
  - [33] Sajadi SM, Khoramdelazad H, Hassanshahi G, et al. Plasma levels of CXCL1 (GRO- $\alpha$ ) and CXCL10 (IP-10) are elevated in type 2 diabetic patients: evidence for the involvement of inflammation and angiogenesis/angiostasis in this disease state. *Clin Lab*. 2013;59:133–137.
  - [34] Hong H, Yuan S, Yang Q, et al. Multiplex bead array assay of plasma cytokines in type 2 diabetes mellitus with diabetic retinopathy. *Mol Vis*. 2014;20:1137–1145.
  - [35] Maury E, Ehala-Aleksejev K, Guiot Y, et al. Adipokines oversecreted by omental adipose tissue in human obesity. *Am J Physiol Endocrinol Metab*. 2007;293:E656–E65.
  - [36] Tourniaire F, Romier-Crouzet B, Lee JH, et al. Chemokine expression in inflamed adipose tissue is mainly mediated by NF- $\kappa$ B. *PLoS ONE*. 2013;8:e66515.
  - [37] Pellegrinelli V, Rouault C, Rodriguez-Cuenca S, et al. Human adipocytes induce inflammation and atrophy in muscle cells during obesity. *Diabetes*. 2015;64:3121–3134.
  - [38] Fujiwara K, Matsukawa A, Ohkawara S, et al. Functional distinction between CXC chemokines, interleukin-8 (IL-8), and growth related oncogene (GRO) $\alpha$  in neutrophil infiltration. *Lab Invest*. 2002;82:15–23.
  - [39] Hurley JR, Cho H, Sheikh AQ, et al. Nanofiber micro-environment effectively restores angiogenic potential of diabetic endothelial cells. *Adv Wound Care*. 2014; 3: 717–728.
  - [40] Yiu K-H, Tse H-F. Specific role of impaired glucose metabolism and diabetes mellitus in endothelial progenitor cell characteristics and function. *Arterioscler Thromb Vasc Biol*. 2014;34:1136–1143.
  - [41] Rodriguez D, Morrison CJ, Overall CM. Matrix metalloproteinases: what do they not do? new substrates and biological roles identified by murine models and proteomics. *Biochim Biophys Acta*. 2010;1803(1):39–54.
  - [42] Tinahones FJ, Coin-Araguez L, Mayas MD, et al. Obesity-associated insulin resistance is correlated to adipose tissue vascular endothelial growth factors and metalloproteinase levels. *BMC Physiol*. 2012;12:12.
  - [43] Sotornik R, Brassard P, Martin E, et al. Update on adipose tissue blood flow regulation. *Am J Physiol Endocrinol Metab*. 2012;302:E1157–E70.
  - [44] Schlich R, Willems M, Greulich S, et al. VEGF in the crosstalk between human adipocytes and smooth muscle cells: depot-specific release from visceral and perivascular adipose tissue. *Mediators Inflamm*. 2013;982458.10.1155/2013/982458
  - [45] Bassols J, Ortega FJ, Moreno-Navarrete JM, et al. Study of the proinflammatory role of human differentiated omental adipocytes. *J Cell Biochem*. 2009;15:1107–1117.
  - [46] Ciaraldi TP, Kolterman OG, Olefsky JM. Mechanism of the post-receptor defect in insulin action in human obesity: decrease in the intrinsic activity of the glucose transport system. *J Clin Invest*. 1981;68:875–880.

- [47] Villaret A, Galitzky J, Decaunes P, et al. Adipose tissue endothelial cells from obese human subjects: differences among depots in angiogenic, metabolic, and inflammatory gene expression and cellular senescence. *Diabetes*. 2010;59:2755–2766.
- [48] Curat CA, Miranville A, Sengenès C, et al. From blood monocyte to adipose tissue resident macrophages. *Diabetes*. 2004;53:1285–1292.
- [49] Srinivasan S, Yeh M, Danzinger EC, et al. Glucose regulates monocyte adhesion through endothelial production of interleukin-8. *Circ Res*. 2003;92:371–377.
- [50] Hongpaisan J. Inhibition of contaminating fibroblasts by D-valine in cultures of smooth muscle cells from human myometrium. 2000; *Cell Biol Intl*. 24:1–7.
- [51] Donovan D, Brown NJ, Bishop ET, et al. Comparison of three in vitro human angiogenesis assays with capillaries formed in vivo. *Angiogenesis*. 2001;4:113–121.
- [52] Tamura Y, Watanabe F, Nakatani T, et al. Highly selective and orally active inhibitors of type IV collagenase (MMP-9 and MMP-2): n-sulfonylamino acid derivatives. *J Med Chem*. 1998;41:640–649.