

UCLA

UCLA Previously Published Works

Title

Enhanced CLIP Uncovers IMP Protein-RNA Targets in Human Pluripotent Stem Cells Important for Cell Adhesion and Survival

Permalink

<https://escholarship.org/uc/item/9hv6q5b8>

Journal

Cell Reports, 15(3)

ISSN

2639-1856

Authors

Conway, Anne E

Van Nostrand, Eric L

Pratt, Gabriel A

et al.

Publication Date

2016-04-01

DOI

10.1016/j.celrep.2016.03.052

Peer reviewed



Published in final edited form as:

Cell Rep. 2016 April 19; 15(3): 666–679. doi:10.1016/j.celrep.2016.03.052.

Enhanced CLIP uncovers IMP protein-RNA targets in human pluripotent stem cells important for cell adhesion and survival

Anne E. Conway^{§,1,2,3}, Eric L. Van Nostrand^{§,1,2}, Gabriel A. Pratt^{1,2,4}, Stefan Aigner^{1,2}, Melissa L. Wilbert^{1,2}, Balaji Sundararaman^{1,2}, Peter Freese⁵, Nicole J. Lambert⁵, Shashank Sathe^{1,2}, Tiffany Y. Liang^{1,2}, Anthony Essex³, Severine Landais³, Christopher B. Burge⁵, D. Leanne Jones^{3,6,7,*}, and Gene W. Yeo^{1,2,8,9,*}

¹Department of Cellular and Molecular Medicine, University of California at San Diego, La Jolla, California 92037, USA

²Stem Cell Program and Institute for Genomic Medicine, University of California at San Diego, La Jolla, California 92037, USA

³Laboratory of Genetics, Salk Institute for Biological Studies, La Jolla, California 92037, USA

⁴Department of Bioinformatics and Systems Biology, University of California at San Diego, La Jolla, California 92093, USA

⁵Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts 02142, USA

⁶Department of Molecular, Cellular and Developmental Biology, University of California at Los Angeles, La Jolla, California 90095, USA

⁷Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California- Los Angeles, California 90095, USA

⁸Molecular Engineering Laboratory, A*STAR, Singapore 1190777

⁹Department of Physiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 1190777

SUMMARY

*Correspondence: geneyeo@ucsd.edu, leannejones@ucla.edu.

§These authors contributed equally.

Accession Numbers

IMP eCLIP and knockdown RNA-seq data have been deposited in the Gene Expression Omnibus (GSE78509). Bind-N-Seq datasets have been deposited at the ENCODE DCC (<https://www.encodeproject.org>) under accession IDs ENCSR928XOW (IMP1) and ENCSR588GYZ (IMP2).

Author Contributions

Conceptualization, A.E.C., E.L.V.N., M.L.W., D.L.J. and G.W.Y.; Investigation, A.E.C., E.L.V.N., S.A., B.S., M.L.W., T.Y.L., P.F., N.J.L.; Validation, A.E. and S.L.; Formal Analysis, A.E.C., E.L.V.N., G.A.P. and S.S.; Writing – Original Draft, A.E.C., E.L.V.N., M.L.W., D.L.J. and G.W.Y.; Writing – Review & Editing, A.E.C., E.L.V.N., D.L.J. and G.W.Y.; Funding Acquisition, C.B.B., D.L.J. and G.W.Y.; Supervision, D.L.J. and G.W.Y.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Human pluripotent stem cells (hPSCs) require precise control of post-transcriptional RNA networks to maintain proliferation and survival. Using enhanced UV crosslinking and immunoprecipitation (eCLIP), we identify RNA targets of the IMP/IGF2BP family of RNA-binding proteins in hPSCs. At the broad region- and binding site-level IMP1 and IMP2 show reproducible binding to a large and overlapping set of 3'UTR-enriched targets. RNA Bind-N-Seq applied to recombinant full-length IMP1 and IMP2 reveals CA-rich motifs that are enriched in eCLIP-defined binding sites. We observe that IMP1 loss in hPSCs recapitulates IMP1 phenotypes, including a reduction in cell adhesion and an increase in cell death. For cell adhesion, in hPSCs we find IMP1 maintains levels of integrin mRNA, specifically regulating RNA stability of *ITGB5*. Additionally, we show IMP1 can be linked to hPSC survival via direct target *BCL2*. Thus, transcriptome-wide binding profiles identify hPSC targets modulating well-characterized IMP1 roles.

INTRODUCTION

Human embryonic stem cells (hESCs) are an invaluable model system to address mechanisms of early human development due to the ability to self-renew and differentiate into the majority of cell types in the mammalian embryo. Recent studies profiling RNA regulatory networks controlled by RNA binding proteins (RBPs), including RBFOX2, LIN28A and MBNL, have demonstrated that RBPs play key roles in maintenance of pluripotency through regulating diverse aspects of RNA processing (Han et al., 2013; Wilbert et al., 2012; Yeo et al., 2009).

The IGF2 mRNA binding proteins (IMPs/IGF2BPs) are a highly homologous family of RBPs that are conserved from insects to mammals (Hansen et al., 2004; Nielsen et al., 1999). Humans and mice have three IMPs (IMP1-3/IGF2BP1-3), which are expressed broadly during early development. Protein expression generally decreases in most tissues post-natally, with the exception of sustained expression in the germ line in adults (Hammer et al., 2005; Hansen et al., 2004). IMP2 mRNA, however, remains expressed in adult murine tissues (Bell et al., 2013). IMP1 is necessary for proper embryogenesis (Hansen et al., 2004), and IMPs are upregulated in many different types of cancer including lung, liver, breast, and colon, with expression being tightly correlated with poor patient prognosis (Dimitriadis et al., 2007; Ross et al., 2001)

Molecular mechanisms of how IMP proteins bind and regulate their target RNAs have been studied predominantly *in vitro*. Molecules of IMP1 protein bind RNA cooperatively and sequentially, dimerizing to form a stable complex with bound RNA via the hnRNP K homology (KH) domains 1–4 (Nielsen et al., 2004). All four KH domains contribute to RNA binding and are important for localization of IMP (Nielsen et al., 2002). Furthermore, both IMP2 and IMP3 are able to heterodimerize on a target RNA with IMP1 via the 4 KH domains (Nielsen et al., 2004). Identification of IMP1 RNA targets *in vivo* revealed IMP1 modulates development and differentiation by regulating various stages of RNA processing. The namesake target of the IMP family, *IGF2*, is primarily regulated at the level of translation, but IMP1 can either promote or repress translation of IGF2 depending on cellular context (Dai et al., 2013; Nielsen et al., 1999). IMP1 also controls the localization

and translation of neuron-specific *Tau* mRNA in a differentiation-dependent manner (Atlas et al., 2007) and controls stability of *MYC* RNA (Bernstein et al., 1992).

Although these studies in cell lines and model organisms have provided clues into IMP regulation of a small number of RNAs, our understanding of how the IMP-RNA target orchestra is conducted transcriptome-wide in human development is incomplete. In HEK293 cells, Hafner and colleagues surveyed the genome-wide binding preferences of all three IMPs over-expressed using Photoactivatable-Ribonucleoside-Enhanced Crosslinking and Immunoprecipitation (PAR-CLIP) (Hafner et al., 2010) and Jonson and colleagues surveyed the RNAs in IMP1 RNP complexes using RIP-Chip (Jonson et al., 2007). However, whether over-expression recapitulates endogenous binding is always a concern with RBPs, and indeed it was recently shown that exogenous expression of IMP1 results in aberrant sedimentation in polysomal gradient centrifugation when compared with endogenous protein (Bell et al., 2013). Therefore, to study the normal roles of endogenous IMP proteins in hESCs we integrated two recently developed approaches: enhanced UV crosslinking and immunoprecipitation followed by high-throughput sequencing (eCLIP) to identify the endogenous RNA targets of IMP1, IMP2 and IMP3 *in vivo*, and RNA Bind-n-seq (RBNS) to uncover the *in vitro* binding preferences of full length IMP1 and IMP2 proteins. These approaches revealed highly overlapping binding for IMP1 and IMP2 that was distinct from IMP3, suggesting the IMP family plays both redundant and distinct functions in hPSCs. Further, loss of IMP1 leads to defects in cell survival and adhesion in hPSCs that can be partially explained through its effects on direct targets *BCL2* and *ITGB5*, respectively. Thus, profiling of endogenous IMP1 targets in hPSCs reveals insight into the pathways through which well-characterized IMP1 functions are achieved in stem cells.

RESULTS

Enhanced CLIP identifies targets of IMP1, IMP2 and IMP3 proteins in human embryonic stem cells

The human IMP family of RNA binding proteins (RBPs) consists of three members (IMP1, IMP2 and IMP3) that contain two RNA recognition motifs (RRMs) and four KH domains each (Figure 1A). Previous reports have observed significant expression of all three IMP proteins in pluripotent and cancer cell lines, with expression in differentiated tissues mostly limited to IMP2 (Bell et al., 2013). Analyzing public RNA-seq datasets (Marchetto et al., 2013), we confirmed that all three members are highly expressed at the mRNA level in PSCs relative to differentiated tissues (Figure 1B). At the protein level, we validated that IMP1, IMP2, and IMP3 are all expressed in undifferentiated human ESC lines H9 and HUES6 and an induced pluripotent stem cell (iPSC) line, whereas IMP2 is also expressed in the parental fibroblasts from which the iPSC line was generated (Figure 1C). Further, immunohistochemical staining (Figure 1D) and subcellular fractionation (Figure 1E) in H9 hESCs demonstrated dominant cytoplasmic localization of all three IMP proteins. Thus, we selected H9 hESC to identify the RNA targets of IMP proteins in pluripotent stem cells.

To uncover molecular pathways in PSCs regulated by IMP proteins, we utilized an enhanced iCLIP (eCLIP) protocol to identify transcriptome-wide RNA targets of the IMP proteins (Konig et al., 2011; Van Nostrand et al., 2016). Briefly, H9 hESCs were subjected to UV-

mediated crosslinking, lysis and treatment with limiting amount of RNase, followed by immunoprecipitation (IP) of protein-RNA complexes using commercially available antibodies that specifically recognize IMP1, IMP2 or IMP3 (Figures 2A and S1A). RNA fragments protected from RNase digestion by IMP protein occupancy were subjected to 3' RNA linker ligation, reverse-transcription and 3' DNA linker ligation to generate eCLIP libraries for high-throughput Illumina sequencing. eCLIP improves these ligations to greater than 70% efficiency, significantly increasing the number of non-PCR duplicate reads that can be obtained after high-throughput sequencing (Van Nostrand et al., 2016). Specificity of the antibodies was evaluated by Western blotting with recombinant human IMP1, IMP2 and IMP3 proteins (Figure S1A). Co-immunoprecipitation experiments in H1 hESCs demonstrate that the IMP1 and IMP2 antibodies do not enrich any of the other family members, while IMP3 appears to slightly co-immunoprecipitate IMP1 (Figure S1B).

We generated biological replicate eCLIP libraries for IMP1 and IMP2, and single replicates for IMP3, a negative control (IgG-only IP) and an unrelated RBP (RBFox2) (Figures S1C–D). The improved efficiency of eCLIP enabled us to generate a Size-Matched Input (SMInput) library for each biological sample, in which 2% of the pre-immunoprecipitation sample was subjected to identical library generation steps including ribonuclear protein complex size-selection on nitrocellulose membranes. In total, ten eCLIP (including SMInput) libraries were sequenced to ~15 million reads, of which ~70% mapped uniquely to the human genome, and ~58–93% of the uniquely mapped reads are distinct after accounting for PCR duplicates (Table S1).

Enrichment of IMP1 and IMP2 binding to 3' untranslated regions

To identify which annotated gene regions are preferentially bound by the IMP proteins, we first evaluated the biological reproducibility of eCLIP (including SMInput) data. Read density within full-length annotated gene regions, namely coding exons (CDS), introns, and 3' untranslated regions (3'UTRs), were highly correlated across replicates for the IMP proteins (R^2 values of ~0.9; Figures S2A and S2B), as well as between IMP proteins and their individually paired SMInput (R^2 values of ~0.7; an example IMP1 replicate is shown in Figure S2C) and IgG (R^2 values of ~0.5; example IMP1 replicate shown in Figure S2D). These high correlations were expected due to intrinsic biological biases in gene expression and shared technical variations in shearing and amplification. Therefore, as a more accurate measure of enrichment (signal) over background, we computed the fold-enrichment in the IP compared to the paired SMInput within each region (Figure 2B, Figures S2E–H, and Table S2). Biological replicates of IMP1 and IMP2 showed significant correlation in fold-enrichment ($R^2 > 0.46$) (Figure 2C and Figure S2I). In contrast, neither IMP1 nor IMP2 showed correlation of binding signal at CDS or 3'UTR regions with either IgG or unrelated RBP RBFox2 (all $R^2 < 0.04$; Figures 2D and S2J–L).

The region fold-enrichments between the paralogs IMP1 and IMP2 were also highly correlated ($R^2 > 0.37$; Figures 2E and S2M). Unexpectedly, despite having the same domain architecture and cytoplasmic localization as IMP1 and IMP2, IMP3 binding within coding exon regions was not correlated with either IMP1 or IMP2 ($R^2 = .03$) (Figures 2F and S2N). Furthermore, we observed that rather than interacting with a specific limited set of RNA

substrates, there was widespread enrichment in binding to particular regions within genes, as 3'UTRs were enriched for IMP1 and IMP2 binding by 2.7 to 4.4-fold (median values) above SMInput IP (Figures 2B, 2G, S2E). This was notably higher than coding exons, which were enriched by only 1.2 to 1.7-fold. Analysis of IMP3 binding revealed an opposite trend, as coding exons were 2.9-fold enriched while 3'UTRs were only 2.3-fold enriched (Figures 2G, S2F). Consistent with their cytoplasmic localization, depletion of intronic signal globally across all introns was observed for all three IMP family members (depleted 1.9 and 2.3-fold for IMP1, 1.8 and 2.1-fold for IMP2, and 1.7-fold for IMP3 respectively) (Figures 2B, 2G, S2E–H). These global enrichments for coding exons (depleted in both) or 3'UTR regions were not observed for IgG or RBFOX2 (1.3-fold enriched in IgG, depleted in RBFOX2). We conclude that the IMP1 and IMP2 proteins exhibit remarkably similar binding preferences to 3'UTRs of mature mRNAs, and IMP3 binding is enriched for coding exons.

Discovery of reproducible IMP1 and IMP2 binding sites by input normalization

Due to their similarity in 3'UTR preferences, we chose to continue evaluating IMP1 and IMP2. To identify high resolution IMP1 and IMP2 binding sites, standard CLIP-seq cluster discovery was performed using CLIPper (Lovci et al., 2013). We identified 62,784 and 95,577 clusters for two biological replicates of IMP1 and 57,648 and 66,928 clusters for IMP2 replicates, with cluster sizes 36–40 bases on average (Figures 3A and S3A, Table S1). Next, as SMInput normalization significantly improves signal-to-noise in identifying true binding sites (Van Nostrand et al., 2016), we compared the read density in IP and SMInput within clusters to compute the enrichment of each cluster above SMInput (Figure S3B). As in the regional comparisons, fold-enrichment at the cluster level exhibited high correlation across biological replicates (R values 0.54 and 0.51 for IMP1 and IMP2, respectively), whereas low correlation was observed when IMP1 (or IMP2) was compared to IgG or RBFOX2 (Figure S3C–F). Additionally, ranking clusters by fold-enrichment makes them amenable to Irreproducible Discovery Rate (IDR) analysis, a standard metric to evaluate the reproducibility of binding sites across biological replicates (Li et al., 2011). We observed that IMP1-IMP1 or IMP2-IMP2 comparisons yielded thousands of reproducible clusters at a 0.01 IDR threshold (where 1% of peaks do not reproduce), whereas identical comparisons with IgG or RBFOX2 yielded less than ten reproducible clusters (Figure 3B). These orthogonal computational approaches indicate that eCLIP of IMP1 and IMP2 yield highly reproducible binding at both the cluster and read-density-within-cluster levels.

For further analysis, we identified a set of 1,884 and 7,004 high-confidence peaks in IMP1 and 1,572 and 4,494 in IMP2 that meet stringent enrichment criteria ($p < 10^{-5}$ and 8-fold enriched versus SMInput) (Figures 3A and S3A). In contrast, IgG eCLIP identified only 142 clusters that satisfy these criteria. These stringent binding sites were highly reproducible, as over 66% of stringent clusters identified in the first biological replicate overlapped clusters in the second for both IMP1 and IMP2 (Figures 3A and S3A). Consistent with our regional analyses, IMP1 and IMP2 binding sites were generally located within the 3'UTR and to a lesser extent, within coding exons (Figures 3C and S3I–J). Thus eCLIP identifies thousands of highly confident and reproducible IMP1 and IMP2 binding sites.

High-resolution IMP1 and IMP2 binding is highly correlated

As we observed substantial correlation between IMP1 and IMP2 binding at the region level we next compared IMP1 and IMP2 at the binding site level. Pairwise comparisons indicated 2,495 and 4,301 peaks (at the 0.01 IDR threshold), on par with that observed for biological replicates (Figure 3B). We further observed high correlation of input-normalized signal intensity ($R = 0.42$ and 0.47 for IMP1_{Rep1} versus IMP2_{Rep2} and IMP1_{Rep2} versus IMP2_{Rep1} respectively), indicating that the association of IMP1 and IMP2 is highly similar across thousands of binding sites (Figure S3G–H). To test whether these factors were associating to the same short regions or simply binding nearby each other, we next considered the distribution of read intensity around peak centers. Considering a window of 600-nt centered on the midpoint of IMP1 peaks from an independent biological replicate, we observed that both IMP1 and IMP2 read density are enriched at the peak centers (Figure 3D). Thus, despite the IMP1 and IMP2 antibodies showing very little cross-immunoprecipitation (Figure S1B), analysis of our eCLIP data at both the region- and cluster-level indicates that IMP1 and IMP2 binding signals are as highly correlated as biological replicates of IMP1 or IMP2.

RNA Bind-N-Seq identifies CA-rich motifs enriched in coding and 3'UTR binding sites

To characterize the sequence specificity of IMP proteins, we applied RNA Bind-N-Seq (RBNS) (Lambert et al., 2014) to purified full-length human IMP1 and IMP2. After incubation of protein with randomized RNA pools, affinity purification, and high-throughput sequencing, we performed motif analysis to calculate enrichment over input (R) values (Figures 4A, S4A–B). This identified two CA-rich motifs for each IMP: a primary motif exemplified by AY(A)1YA and secondary motif exemplified by Y(A)2YA (Figure 4B–C), with many enriched 6-mers (52% for IMP1, 49% for IMP2) containing one of four 4-mers (CACA, UACA, AACA, CAUA), similar to previously identified IMP motifs (Alipanahi et al., 2015; Hafner et al., 2010; Ray et al., 2013). We observed a high correlation between IMP1 and IMP2 6-mer enrichments ($R^2=0.788$) (Figure 4D), whereas IMP1 and RBFOX2 were uncorrelated ($R^2=0.018$) (Figure 4E), indicating that the RBNS assay captured IMP-specific binding signatures.

Next we interrogated whether hESC IMP binding sites identified by eCLIP were enriched for the RBNS-identified *in vitro* motifs. We found that hexamers containing the CACA core sequence were shifted towards higher IMP1 RBNS enrichments, and they also showed greater enrichments among IMP1 eCLIP 3'UTR and CDS binding sites than other hexamers (Figures 4F, S4C). Other RBNS motifs (UACA, AACA, CAUA) showed more variable enrichment, suggesting a distinguishable difference between *in vitro* and *in vivo* binding preferences (Figure 4F–H). The enrichment for the CACA motif was more significant when using the subset of stringent CLIP-enriched peaks, consistent with these peaks having improved signal-to-noise (Figure 4G, S4D). IMP2 showed similar enrichment for CA-rich sequences in both RBNS and eCLIP binding sites (Figure 4H). Thus, these results demonstrate that IMP1 and IMP2 interact with CA-rich sequences *in vitro*, and this preference can be observed for coding and 3'UTR binding sites *in vivo*.

Integrated IMP1 eCLIP and RNA-seq data implicates integrin mRNAs in IMP1-mediated cell adhesion defects

Next, we utilized our IMP1 binding data to provide insight into the direct regulatory roles of IMP1. To evaluate if loss of IMP1 affected mRNA expression, IMP1 was depleted in H9 hESCs using lentiviral transduction of independent short-hairpin RNAs (shRNAs) that specifically target IMP1 (hereafter referred to as IMP1 knock-down (IMP1^{KD}) cells)(Figure S5A–B). Total RNA was extracted from three biologically independent transductions of IMP1 shRNA and two transductions of a non-targeting shRNA to generate RNA-seq libraries, which were then sequenced (Figure S5C). Of the 17,013 expressed genes analyzed, we identified 257 decreased and 467 increased genes with significantly altered expression (two-fold & $p < 0.05$ versus non-targeting control) upon IMP1 depletion (Figure S5D). When we compared global IMP1 RNA targets to genes affected by IMP1 depletion, we did not observe any meaningful correlation between genes that were bound and trends in gene expression changes (Figure S5E).

Nevertheless, when we considered the most CLIP-enriched 3'UTRs for IMP1, we observed significant enrichment for genes involved in cell-cell and cell-extracellular matrix (ECM) adhesion (Figure 5A). As IMP1 regulation of cell adhesion has been well characterized in cancer cell-lines (Gu et al., 2012; Vikesaa et al., 2006), we performed a quantitative crystal violet adherence assay following IMP1 depletion to determine whether loss of IMP1 affected adhesion in H9 hESCs. We found that IMP1^{KD} cells showed significantly decreased adherence compared to control shRNA treated cells ($p < 0.05$), even within the first hour after plating, indicating that cell adhesion is also affected by loss of IMP1 in hESCs (Figures 5B–D). We also observed that the actin cytoskeleton and cytoskeletal organization appeared disrupted upon IMP1 depletion (Figure S5F–G). However, analysis of our RNA-seq data did not indicate altered RNA levels of known IMP1 targets previously shown to modulate cell adhesion, such as *CTNNB1* (in breast cancer cells) (Gu et al., 2008) or *CD44* (in adenocarcinoma cells)(Vikesaa et al., 2006). In particular, despite having enriched IMP1 3'UTR binding (Figure S5H), we could observe no mRNA or protein change for *CTNNB1* in IMP1^{KD} cells (Figure S5I). Similarly, although the F-actin anchoring, cell adhesion protein Vinculin (*VCL*) was one of the most enriched genes in our eCLIP dataset (over 20-fold enriched relative to SMInput), *VCL* did not appear to be affected at the RNA or protein level upon loss of IMP1 (Figures S5J–K). Thus, we observe cell adhesion defects as a consequence of IMP1 depletion in hESCs, but the phenotype cannot be explained simply by regulation through previously characterized IMP1 targets.

As IMP1 binding alone was insufficient to predict mRNA level effects, we next focused on IMP1 target genes that not only harbored enriched binding sites in the 3'UTR and whose levels were affected in the IMP1^{KD} RNA-seq data. Surprisingly, we observed that multiple genes in the integrin family were bound by IMP1 and were downregulated upon depletion of IMP1 (Figures 5E–F). Integrins are known to have significant roles in extracellular signaling and cell adhesion across various systems, with *ITGB5* and *ITGB1* specifically described to play key roles in human stem cell maintenance and cell adhesion (Braam et al., 2008). Interestingly, *ITGB5* was the most down-regulated of all of the integrin genes and contained regions of enriched read density within the 3'UTR compared to SMInput or RBFOX2

(Figure 5G). We performed RNA Immunoprecipitation (RIP) followed by RT-PCR in an independent hESC line (HUES6) and observed IMP1 enrichment on *ITGB5*, with *ACTB* mRNA as a positive control (Ross et al., 1997)(Figure 5H), validating the interaction between IMP1 and *ITGB5* mRNA. Additionally, we confirmed down-regulation of *ITGB5* and *ITGB1* mRNAs in IMP1^{KD} cells using two independent shRNAs (Figure 5I) and further observed significant depletion of ITGB5 protein upon IMP1 loss (Figure 5J).

Next, we considered potential mechanisms for IMP1 regulation of *ITGB5*. To test whether IMP1 affects *ITGB5* mRNA levels post-transcriptionally at the level of mRNA turnover, we treated hESCs with Actinomycin D (ActD) to inhibit the transcription of newly transcribed RNA and collected total RNA after 60 and 120 minutes. Quantification of mRNA levels by qRT-PCR revealed that *ITGB5* was destabilized more quickly in the IMP1^{KD} cells compared to cells treated with a control shRNA (Figure 5K). Interestingly, *ITGB1* did not show this destabilization upon IMP1 depletion (Figure S5L), confirming specificity of this approach and indicating that IMP1 may regulate various integrins differently. Taken together, these results indicate that the well-characterized role of IMP1 in maintaining proper cell-cell interactions is conserved in hESCs, but that the downstream effectors in hESCs include unanticipated integrin targets like *ITGB5*, which is regulated at the level of mRNA turnover.

IMP1 target *BCL2* enhances survival of IMP1-depleted hESCs

In addition to cytoskeletal defects, depletion of IMP1 also led to a drastic reduction in hESC colony size (Figure 6A). Embryoid bodies derived from IMP1^{KD} cells were substantially and consistently smaller than those derived from controls (Figure S6A). In order to determine whether depletion of IMP1 led to a decrease in proliferation, which could explain a decrease in colony size, we performed fluorescence activated cell sorting (FACS) analysis using the proliferation antigen KI-67. We observed only a slight, but insignificant, difference between IMP1 depleted and control cells using two independent shRNAs targeting IMP1 (Figure S6B). To further analyze a potential role for IMP1 in hESC proliferation, cell cycle analysis was conducted by BrdU and propidium iodide (PI) staining followed by FACS. IMP1^{KD} cells exhibited a moderate, but significant decrease in the S phase population ($p < 0.01$), along with an increase in the number of cells in G2 ($p < 0.05$) (Figure S6C). Supporting the hypothesis that a loss of IMP1 leads to an increase in cell death, we detected a statistically significant increase in AnnexinV-positive IMP1^{KD} cells compared to controls by FACS ($p < 0.05$)(Figure 6B). Additionally, we observed a significant increase in levels of cleaved-Caspase 3 upon reduction of IMP1, but not in control treated hESCs (Figure 6C). Together, these results strongly indicate that IMP1 plays an important role in hPSC survival.

To determine the molecular mechanism by which IMP1 impacts hESC survival, we examined anti-apoptotic factors in our IMP1^{KD} RNA-seq data as candidate targets of IMP1 and found that *BCL2* (B-cell lymphoma 2) was decreased by two-fold (Table S4). This decrease in *BCL2* was confirmed at the mRNA level by qRT-PCR (Figure 6D) and at the protein level by Western blot analysis (Figure 6C, S6D). Interestingly, the *BCL2* 3'UTR contains CA-rich sequences (Figure S6E) and was greater than two-fold enriched over SMInput in both IMP1 eCLIP datasets; however, *BCL2* fell below our standard significance cutoffs due to low RNA expression (Table S2). Nevertheless, we confirmed IMP1 binding by

performing RIP followed by RT-PCR in HUES6 hESCs (Figure 6E). To further investigate IMP binding preferences to CA-rich sequences, as previously shown by RBNS, we performed an electrophoretic mobility shift assay (EMSA) with full-length recombinant human IMP1 protein and both wild-type (WT) and mutated versions of the *BCL2* 3'UTR (88bp segments, see Methods). The wild-type *ACTB* zipcode (IMP binding site) and a truncated *ACTB* zipcode were used as positive and negative controls, respectively. We were able to detect an interaction between IMP1 and the wild type *BCL2* 3'UTR *in vitro* using 200nM recombinant IMP1 protein (Figure 6F). Upon mutation of the CACA motif to GAGA, the binding affinity was dramatically reduced, indicating that IMP1 interacts specifically with the CA-rich motif that constitutes an IMP1 binding site within *BCL2* 3'UTR. Finally, to evaluate whether restoring levels of *BCL2* can suppress cell death as a consequence of IMP1 depletion, we utilized a doxycycline-inducible lentiviral system (Ardehali et al., 2011) to ectopically express *BCL2* in IMP1^{KD} hESCs. Using the Caspase-Glo assay we measured apoptosis following a titration of *BCL2* over-expression with doxycycline and found that *BCL2* is able to rescue the IMP1^{KD} cell death phenotype (Figures 6G, S6F–G). Therefore, our data indicate that one pro-survival function of IMP1 in hESCs is to maintain adequate levels of *BCL2* mRNA and, consequently, maintain its anti-apoptotic activity.

DISCUSSION

Using systematic, transcriptome-wide mapping with eCLIP we identified thousands of IMP1, IMP2 and IMP3 binding sites within RNA targets in hESCs. IMP1 and IMP3 are typically viewed as the most related family members, with greater similarity at the protein sequence level (Nielsen et al., 1999), expression patterns across tissues and development (Bell et al., 2013), and co-immunoprecipitation during CLIP (Figure S1B), whereas IMP2 has been associated with more distinct roles, such as in metabolism (Dai et al., 2011; Dai et al., 2015; Janiszewska et al., 2012). Thus, hESCs (which express IMP1, IMP2, and IMP3) present a unique opportunity to observe redundant or co-regulation of RNA targets by multiple IMP family members. Surprisingly, we observed substantial overlap between IMP1 and IMP2 binding that was not observed between IMP1 and IMP3, indicating it is not simply an artifact of analyzing cytoplasmic factors with CLIP. Despite the large number of IMP1-bound mRNAs and quite dramatic phenotypes upon knockdown of IMP1 in hESCs, we observed relatively few transcripts strongly bound by IMP1 to be altered when RNA-seq was performed in hESCs depleted of IMP1. In contrast, previous studies in HEK293 cells observed a small but significant shift towards decreased expression (presumably by decreased RNA stability) of IMP targets when all three IMP proteins were depleted (Hafner et al., 2010). These results suggest that IMP family members may share redundant regulatory roles, particularly during development and in cancer when multiple family members are expressed at high levels. The distinct binding to CDS regions observed for IMP3 suggests that further studies may yet reveal additional regulatory roles distinct from IMP1, although the severe phenotypes observed upon individual knockdown by shRNA of IMP1 and IMP2 presents a challenge to detailed characterization of redundancy among IMP family members in hESCs.

At the region-level, we observed significant IMP1 binding to a substantial fraction of all 3'UTRs. Although such widespread binding has been described for core RNA processing factors, such as the nonsense-mediated decay regulator UPF1 (Lee et al., 2015), previous studies of IMP have largely focused on a small number of specific targets. Although the degree of widespread binding is unexpected, it may help to explain why detailed studies of individual IMP targets have not identified one specific mechanism or pathway of regulation; rather, IMP binding has been shown to participate in a broad range of RNA processing regulation steps, including mRNA stability (Leeds et al., 1997), mRNA localization (Atlas et al., 2007; Ross et al., 1997), both inhibition of and enhancement of translation (Dai et al., 2011; Nielsen et al., 1999), and even potentially nuclear export (Huttelmaier et al., 2005; Wu et al., 2015). Thus, considering IMPs as broad regulators may provide insight into how these factors can achieve these various roles. Future work will be needed to better characterize how different IMP targets are directed towards distinct regulatory mechanisms.

Although standard UV₂₅₄ crosslinking provides specificity by requiring interacting molecules to have reactive groups within one bond length apart (Wagenmakers et al., 1980), the frequency of crosslinking is substantially decreased at non-Uridine bases (Sugimoto et al., 2012), limiting the ability of IMPs to crosslink at a CA-rich motif. Thus, to complement our eCLIP profiling data we performed RBNS to characterize the *in vitro* binding motif for full-length IMP1 and IMP2 proteins (each including all six RNA binding domains). This method revealed that recombinant IMP1 and IMP2 proteins harbor strong preferences for CA-rich motifs with a degenerate CAU within them, correlating well with the MAHWCA motifs identified for IMP2 and IMP3 using an independent *in vitro* method RNACompete (Ray et al., 2013) as well as the CA motif identified in DeepBind re-analysis of RNACompete data (Alipanahi et al., 2015). Profiling *in vivo* by PAR-CLIP identified a similar but slightly altered motif (CAU) (Hafner et al., 2010), possibly influenced by crosslinking at 4SU nucleotides in PAR-CLIP. These motifs all share strong similarity with motifs described by detailed molecular studies of individual IMP targets (Chao et al. (2010)). The CA-rich motif was significantly enriched in eCLIP peaks, providing further validation that this motif likely represents a major component of IMP1 and IMP2 binding. As IMP binding is often complex, requiring specific spacing of associated motifs to drive dimerization (potentially of multiple IMP family members) (Nielsen et al., 2004), more detailed biochemical studies should provide insight into whether the CA repeat is more critical for binding initiation or stabilization of IMP complexes, and what role these other motifs play in directing IMP target recognition.

Given the relatively small transcriptome change observed upon IMP1 knockdown in hESCs, we were surprised to observe that loss of IMP1 led to dramatic cellular phenotypes, including increased apoptosis and a loss of cell adhesion and cytoskeletal integrity. As IMP roles in maintenance of cell adhesion have been previously described (Gu et al., 2012; Vikesaa et al., 2006), we asked whether the cell adhesion defect in hESCs could be explained by known or novel targets. We observed that there was a specific enrichment for downregulation of integrin mRNAs, particularly those most strongly bound by IMP1 (Figure 5). Further analysis validated IMP regulation of the stability of *ITGB5*. These results indicate that in addition to known IMP1 regulatory targets such as CTNBN1 and CD44 that play critical roles in maintenance of proper cell adhesion (Gu et al., 2012; Vikesaa et al.,

2006), the integrin family represents an additional cell adhesion regulatory mechanism for IMP1. Future work to determine whether integrin regulation by IMP1 is specific to hESCs or affects cell adhesion in other systems and the direct mechanisms (including additional co-factors) through which IMP1 binding modulates stability of *ITGB5* will provide further insight into the cell-type-specificity of the mechanisms through which IMPs regulate cell adhesion.

Although loss of cell adhesion partially explains the dramatic hESC cell viability defect upon IMP1 knockdown, we noted that apoptotic markers were also increased. Global analyses of either eCLIP-bound or differential transcripts did not show general enrichments for apoptosis or related pathways. However, closer inspection of IMP1 bound and responsive targets identified *BCL2*, which encodes a critical anti-apoptotic protein, as another IMP1 target that decreases upon IMP1 depletion in hPSCs. Pursuing this further due to the well-characterized roles of *BCL2* in mediating apoptotic signals, we found that re-expression of *BCL2* in hESCs partially rescued the cell death phenotype resulting from IMP1 depletion in a dose-dependent manner, confirming the contribution of *BCL2* to IMP1 knockdown phenotypes in hESCs. The linkage of IMP1 with direct regulation of *BCL2* further associates IMP1 with known oncogenic pathways and may provide an interesting avenue for further studies of IMP1 in other cell types, particularly with respect to understanding its roles in modulating tumorigenesis and metastasis. Thus, our results indicate that we are far from an exhaustive list of functional IMP family targets, and that further elucidation of the direct and regulated targets of IMP proteins in their various cellular contexts (whether in normal or cancerous cell-types) may provide insights into the distinct and shared roles these proteins play in development and tumorigenesis.

Experimental Procedures

eCLIP-seq Experimental Procedures

10×10^6 UV-crosslinked (400 mJ/cm^2 constant energy) H9ES (IMPs, IgG) or H1ES (RBFOX2) cells were lysed in iCLIP lysis buffer and sonicated (BioRuptor). Lysate was treated with RNase I (Ambion) to shear RNA, after which IMP1 (MBL, #RN007P), IMP2 (MBL, #RN008P), IMP3 (MBL, #RN009P), RBFOX2 (Bethyl Laboratories, #A300-864A), and rabbit IgG (Life Technologies) protein-RNA complexes were immunoprecipitated using the indicated antibody. In addition to the RBP-IPs a parallel Size-Matched Input (SMInput) library was generated; these samples were not immunoprecipitated with anti-RBP antibodies but were otherwise treated identically (to aid in the removal of false positives). One SMInput was used for each biological replicate grouping of all IMP proteins due to their similarity in molecular weight, with a separate SMInput generated for RBFOX2. Stringent washes were performed as described in iCLIP, during which RNA was dephosphorylated with FastAP (Fermentas) and T4 PNK (NEB). Subsequently, a 3' RNA adapter was ligated onto the RNA with T4 RNA ligase (NEB). Protein-RNA complexes were run on an SDS-PAGE gel, transferred to nitrocellulose membranes, and RNA was isolated off the membrane identically to standard iCLIP. After precipitation, RNA was reverse transcribed with AffinityScript (Agilent), free primer was removed (ExoSap-IT, Affymetrix), and a 3' DNA adapter was ligated onto the cDNA product with T4 RNA ligase (NEB). Libraries were then amplified

with Q5 PCR mix (NEB). See Van Nostrand, *et al.* (2016) for further details regarding standardized eCLIP experimental workflows.

eCLIP-seq Read Processing and Cluster Analysis

Briefly, reads were adapter trimmed (cutadapt), mapped against repetitive elements (with repeat-mapping reads discarded), and then mapped to the human genome with STAR. PCR duplicate reads were removed, and the second (paired-end) read was used to perform peak-calling with CLIPper (Lovci et al., 2013). Region-level analysis was performed by counting reads overlapping regions annotated in Gencode (v19). Input normalization of peaks was performed by counting reads mapping to CLIPper-identified peaks in eCLIP and paired SMInput datasets, with significance thresholds of $p < 10^{-5}$ and fold-enrichment > 8 . See Supplemental Experimental Procedures for further details, and Van Nostrand, *et al.* (2016) for software packages used and additional description of processing steps.

hPSC Cell Culture

All hPSC lines (including H9, H1, HUES6, and iPSC) lines were grown on Matrigel (BD Biosciences) using mTeSR1 medium (Stem Cell Technologies). Cells were routinely passaged using Dispase (2mg/ml) and scraping the colonies with a glass pipet. For assays requiring single-cell dissociation, Accutase (Innovative Cell Technologies, Inc) was used followed by culture medium supplemented with 10 μ M Rock Inhibitor Y-26732 (Calbiochem) for 24 hours. See Supplemental Experimental Procedures for further details.

Lentiviral Vectors, Production and hESC Infection

If not otherwise indicated, experiments were performed using pLKO lentivirus constructs TRCN0000075149 for IMP1, TRCN0000255463 for IMP2, TRCN0000074675 for IMP3, and non-target control Sigma #SHC002 (which targets turboGFP). Two additional shRNAs were tested for IMP1, TRCN0000218079 which targeted the CDS (shRNA 2) and TRCN0000230114 which targets the IMP1 3'UTR (shRNA 3). Unless otherwise noted, shRNA 3 (TRCN0000230114) was used as the second shRNA for phenotypic experiments. See Supplemental Experimental Procedures for further details.

Adhesion Assay

After virus transduction and puromycin selection, hES cells were plated out at 50,000 cells per well in a 96 well plate coated with matrigel, incubated for 1 hour at 37°C with 5% CO₂, vortexed at 2000rpm for 15 seconds, washed three times with .1% BSA in DMEM/F12, and fixed with 4% paraformaldehyde for 10 minutes at room temperature. Following fixation, cells were washed with .1% BSA in DMEM/F12 and stained with Crystal Violet (5mg/ml in 2% EtOH) for 10 minutes. Cells were then rinsed with H₂O, and left to completely dry for 15 minutes. 2% SDS was added for 20 minutes followed by absorbance reading on a plate reader. Data are represented as mean \pm SD, with statistical significance calculated by unpaired *t*-test.

BCL2 Rescue Apoptosis Assay

H9 hESCs expressing control and IMP1 shRNAs were split into 4 biological replicates each and transduced with dox-inducible BCL2 viruses. After 24 hours, 3 replicates were each replated into 4 wells of a 96 well plate at a density of 20,000 cells per well. A titration of doxycycline was added for 24 hours after which apoptosis was measured by the Caspase-Glo 3/7 assay (Promega). The Caspase-Glo 3/7 assay was performed according to the manufacturer's instructions. The 4th replicate was collected in parallel for Western blot analysis of BCL2 induction. Luminescence and fluorescence data were averaged across the 4 technical replicate wells, and are represented as mean \pm SD of biological replicates, with statistical significance calculated by unpaired *t*-test.

Western Blot

Cells were washed with PBS and lysed with lysis buffer (10 mM Tris-HCl (pH 8), 150 mM NaCl, 1% Triton X100 and complete protease inhibitor mixture (Roche)). Total protein extracts were run on 4–12% NuPAGE Bis-Tris gels in NuPAGE MOPS running buffer (Thermo Fisher), transferred to nitrocellulose membranes (Amersham Biosciences) and analyzed using primary antibodies. Primary antibodies were incubated overnight at 4°C and secondary HRP conjugated antibodies (Jackson ImmunoResearch, 1:10,000) were incubated for 1 hour at room temperature. Thermo Pierce ECL detection reagents were used. See Supplemental Experimental Procedures for full list of antibodies used.

RNA extraction and RT-qPCR analysis

Total RNA was isolated using Trizol Reagent (Invitrogen) according to the manufacturer's recommendations, DNase treated with Turbo DNA-free kit (Ambion), and cDNA synthesized from 2 μ g total RNA using the SuperScript III Reverse Transcriptase kit for RT-qPCR (Invitrogen). Both random hexamers and oligo(dT) primers were used for reverse transcription. Real-time PCR was performed using the SYBR-Green FAST qPCR Master mix (Applied Biosystems) on a CFX384 Touch Real-Time PCR Detection System (Bio-Rad). Values of gene expression were normalized using *18s* and/or *GAPDH* and *HMBS* (see figure legends for details) expression and are shown as fold-change relative to the value of the control shRNA treated sample. All experiments were performed in technical and biological triplicates. Bars indicate mean \pm SEM as measured by the Ct method. Significance was determined by unpaired *t*-test between the Control shRNA sample and either IMP1 shRNA sample unless otherwise described in figure legend. See Supplemental Experimental Procedures for primer sequences used.

Actinomycin D RNA Stability Assay

H9 hESCs were treated with 10 μ g/ml Actinomycin D (Sigma) and RNA was isolated using Trizol at time 0 (no treatment), 60 minutes and 120 minutes after treatment. RNA decay was measured using RT-qPCR normalized to the amount of RNA at time 0 (see "RNA extraction and RT-qPCR analysis" above). Values of gene expression were normalized using *PPIA* and *RPLP0*, genes determined not to change over the given time period following addition of Actinomycin D.

RNA Immunoprecipitation (RIP) Assay in hESCs

RIP was performed using an input of $\sim 20 \times 10^6$ uncrosslinked HUES6 hES cells lysed with CLIP lysis buffer (Wilbert et al., 2012). 5 μ g of each antibody, Rabbit IgG (Santa Cruz, #SC-2027) and IMP1 (MBL, #RN007P), were coupled to Protein G Dynabeads (Invitrogen) and incubated with pre-cleared cell lysate overnight on rotation at 4°C. Immunoprecipitated RNA was isolated from beads using 1mL Trizol according to the manufacturer's instructions and reverse-transcribed using the SuperScript III Reverse Transcriptase kit. Endpoint RT-PCR was performed using Crimson Taq DNA Polymerase (NEB).

Electrophoretic Mobility Shift Assay (EMSA)

IMP1 EMSA was performed based on the conditions used in (Farina et al., 2003). Binding reactions contained the following: 10 mM Tris pH 7.5, 50 mM NaCl, 1 mM MgCl₂, 0.5 mM EDTA, 0.5 mM DTT, 4% glycerol, 100ng *E. coli* tRNA (Roche), 10,000 cpm of the ³²P-labeled target RNA, and increasing amounts of recombinant human IMP1 protein (Origene # TP316226) to a final volume of 20 μ l. Reactions were incubated for 30 min at room temperature followed by 15 minutes on ice. Reactions were loaded on 5% nondenaturing polyacrylamide TGE gels containing 5% glycerol and run at 150V for 2 hours. Following electrophoresis, gels were dried and exposed to phosphorimager film. See Supplemental Experimental Procedures for further details.

RNA Bind-N-Seq (RBNS)

RBNS was performed on full-length GST-SBP-tagged IMP1 and IMP2 as previously described (Lambert et al., 2014). The motif frequency in the RBP-selected pool divided by the frequency in the input RNA library was calculated for all *k*-mers (*k*=4, 5, 6) and defined as the motif enrichment (R) value. Mean and standard deviation of R values were calculated across all *k*-mers for each *k* (4-, 5-, and 6-mers), with significance thresholds set at z-score 2. Enrichments from the protein concentration with the greatest enrichment were used for comparison with CLIP peaks, compared against same-sized regions randomly selected from the same genomic region (e.g. 3'UTRs or CDS.). See Supplemental Experimental Procedures for further details.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank members of the Jones and Yeo labs for constructive discussions and comments on the manuscript, the Salk Institute Stem Cell Core, especially Dr. Travis Berggren, Margaret Lutz, and Ken Diffenderfer for their invaluable assistance. We thank Patrick Liu for computational analysis assistance, and Reza Ardehali, Matt Inlay, and Irving Weissman for sharing BCL2 plasmids. This work was supported by the Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at the University of California- Los Angeles and a California Institute of Regenerative Medicine (CIRM) New Faculty Award (RN1-00544) to D.L.J. and by grants from the NIH (HG004659, NS075449 and U54HG007005) and CIRM (RB1-01413 and RB3-05009) to G.W.Y., and was partially supported by NIH grant HG007005 to C.B.B. A.E.C and M.L.W. were supported in part by the University of California, San Diego, Genetics Training Program through an institutional training grant from the NIGMS, T32 GM008666. S.L. and A.E. were supported by Pioneer postdoctoral fellowship awards from the Salk Institute for Biological Studies. M.L.W. was a CIRM pre-doctoral fellow, E.L.V.N. is a Merck Fellow of the Damon Runyon Cancer Research Foundation (DRG-2172-13) and G.W.Y. was an Alfred P. Sloan Research Fellow.

References

- Alipanahi B, Delong A, Weirauch MT, Frey BJ. Predicting the sequence specificities of DNA- and RNA-binding proteins by deep learning. *Nature biotechnology*. 2015; 33:831–838.
- Ardehali R, Inlay MA, Ali SR, Tang C, Drukker M, Weissman IL. Overexpression of BCL2 enhances survival of human embryonic stem cells during stress and obviates the requirement for serum factors. *PNAS*. 2011; 108:3282–3287. [PubMed: 21300885]
- Atlas R, Behar L, Sapoznik S, Ginzburg I. Dynamic association with polysomes during P19 neuronal differentiation and an untranslated-region-dependent translation regulation of the tau mRNA by the tau mRNA-associated proteins IMP1, HuD, and G3BP1. *Journal of neuroscience research*. 2007; 85:173–183. [PubMed: 17086542]
- Bell JL, Wachter K, Muhleck B, Pazaitis N, Kohn M, Lederer M, Huttelmaier S. Insulin-like growth factor 2 mRNA-binding proteins (IGF2BPs): post-transcriptional drivers of cancer progression? *Cellular and molecular life sciences: CMLS*. 2013; 70:2657–2675. [PubMed: 23069990]
- Bernstein PL, Herrick DJ, Prokipcak RD, Ross J. Control of c-myc mRNA half-life in vitro by a protein capable of binding to a coding region stability determinant. *Genes & development*. 1992; 6:642–654. [PubMed: 1559612]
- Braam SR, Zeinstra L, Litjens S, Ward-van Oostwaard D, van den Brink S, van Laake L, Lebrin F, Kats P, Hochstenbach R, Passier R, et al. Recombinant vitronectin is a functionally defined substrate that supports human embryonic stem cell self-renewal via alphavbeta5 integrin. *Stem cells*. 2008; 26:2257–2265. [PubMed: 18599809]
- Chao JA, Patskovsky Y, Patel V, Levy M, Almo SC, Singer RH. ZBP1 recognition of beta-actin zipcode induces RNA looping. *Genes & development*. 2010; 24:148–158. [PubMed: 20080952]
- Dai N, Christiansen J, Nielsen FC, Avruch J. mTOR complex 2 phosphorylates IMP1 cotranslationally to promote IGF2 production and the proliferation of mouse embryonic fibroblasts. *Genes & development*. 2013; 27:301–312. [PubMed: 23388827]
- Dai N, Rapley J, Angel M, Yanik MF, Blower MD, Avruch J. mTOR phosphorylates IMP2 to promote IGF2 mRNA translation by internal ribosomal entry. *Genes & development*. 2011; 25:1159–1172. [PubMed: 21576258]
- Dai N, Zhao L, Wrighting D, Kramer D, Majithia A, Wang Y, Cracan V, Borges-Rivera D, Mootha VK, Nahrendorf M, et al. IGF2BP2/IMP2-Deficient mice resist obesity through enhanced translation of Ucp1 mRNA and Other mRNAs encoding mitochondrial proteins. *Cell metabolism*. 2015; 21:609–621. [PubMed: 25863250]
- Dimitriadis E, Trangas T, Milatos S, Foukas PG, Gioulbasanis I, Courtis N, Nielsen FC, Pandis N, Dafni U, Bardi G, et al. Expression of oncofetal RNA-binding protein CRD-BP/IMP1 predicts clinical outcome in colon cancer. *International journal of cancer Journal international du cancer*. 2007; 121:486–494. [PubMed: 17415713]
- Farina KL, Huttelmaier S, Musunuru K, Darnell R, Singer RH. Two ZBP1 KH domains facilitate beta-actin mRNA localization, granule formation, and cytoskeletal attachment. *The Journal of cell biology*. 2003; 160:77–87. [PubMed: 12507992]
- Gu W, Katz Z, Wu B, Park HY, Li D, Lin S, Wells AL, Singer RH. Regulation of local expression of cell adhesion and motility-related mRNAs in breast cancer cells by IMP1/ZBP1. *J Cell Sci*. 2012; 125:81–91. [PubMed: 22266909]
- Gu W, Wells AL, Pan F, Singer RH. Feedback regulation between zipcode binding protein 1 and beta-catenin mRNAs in breast cancer cells. *Mol Cell Biol*. 2008; 28:4963–4974. [PubMed: 18490442]
- Hafner M, Landthaler M, Burger L, Khorshid M, Hausser J, Berninger P, Rothballer A, Ascano M Jr, Jungkamp AC, Munschauer M, et al. Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell*. 2010; 141:129–141. [PubMed: 20371350]
- Hammer NA, Hansen T, Byskov AG, Rajpert-De Meyts E, Grondahl ML, Bredkjaer HE, Wewer UM, Christiansen J, Nielsen FC. Expression of IGF-II mRNA-binding proteins (IMPs) in gonads and testicular cancer. *Reproduction*. 2005; 130:203–212. [PubMed: 16049158]
- Han H, Irimia M, Ross PJ, Sung HK, Alipanahi B, David L, Golipour A, Gabut M, Michael IP, Nachman EN, et al. MBNL proteins repress ES-cell-specific alternative splicing and reprogramming. *Nature*. 2013; 498:241–245. [PubMed: 23739326]

- Hansen TVO, Hammer NA, Nielsen J, Madsen M, Dalbaeck C, Wewer UM, Christiansen J, Nielsen FC. Dwarfism and Impaired Gut Development in Insulin-Like Growth Factor II mRNA-Binding Protein 1-Deficient Mice. *Molecular and Cellular Biology*. 2004; 24:4448–4464. [PubMed: 15121863]
- Huttelmaier S, Zenklusen D, Lederer M, Dichtenberg J, Lorenz M, Meng X, Bassell GJ, Condeelis J, Singer RH. Spatial regulation of beta-actin translation by Src-dependent phosphorylation of ZBP1. *Nature*. 2005; 438:512–515. [PubMed: 16306994]
- Janiszewska M, Suva ML, Riggi N, Houtkooper RH, Auwerx J, Clement-Schatlo V, Radovanovic I, Rheinbay E, Provero P, Stamenkovic I. Imp2 controls oxidative phosphorylation and is crucial for preserving glioblastoma cancer stem cells. *Genes & development*. 2012; 26:1926–1944. [PubMed: 22899010]
- Jonson L, Vikesaa J, Krogh A, Nielsen LK, Hansen Tv, Borup R, Johnsen A, Christiansen J, Nielsen FC. Molecular Composition of IMP1 Ribonucleoprotein Granules. *Molecular and Cellular Proteomics*. 2007; 6:798–811. [PubMed: 17289661]
- Konig J, Zarnack K, Rot G, Curk T, Kayikci M, Zupan B, Turner D, Luscombe N, Ule J. iCLIP - Transcriptome-wide Mapping of Protein-RNA Interactions with Individual Nucleotide Resolution. *Journal of Visualized Experiments*. 2011
- Lambert N, Robertson A, Jangi M, McGeary S, Sharp PA, Burge CB. RNA Bind-n-Seq: quantitative assessment of the sequence and structural binding specificity of RNA binding proteins. *Molecular cell*. 2014; 54:887–900. [PubMed: 24837674]
- Lee SR, Pratt GA, Martinez FJ, Yeo GW, Lykke-Andersen J. Target Discrimination in Nonsense-Mediated mRNA Decay Requires Upf1 ATPase Activity. *Molecular cell*. 2015; 59:413–425. [PubMed: 26253027]
- Leeds P, Kren B, Boylan J, Betz N, Steer C, Gruppuso P, Ross J. Developmental regulation of CRD-BP, an RNA-binding protein that stabilizes c-myc mRNA in vitro. *Oncogene*. 1997; 14:1279–1286. [PubMed: 9178888]
- Li Q, Brown JB, Huang H, Bickel PJ. Measuring reproducibility of high-throughput experiments. *The Annals of Applied Statistics*. 2011; 5:1752–1779.
- Lovci MT, Ghanem D, Marr H, Arnold J, Gee S, Parra M, Liang TY, Stark TJ, Gehman LT, Hoon S, et al. Rbfox proteins regulate alternative mRNA splicing through evolutionarily conserved RNA bridges. *Nature structural & molecular biology*. 2013; 20:1434–1442.
- Marchetto MC, Narvaiza I, Denli AM, Benner C, Lazzarini TA, Nathanson JL, Paquola AC, Desai KN, Herai RH, Weitzman MD, et al. Differential L1 regulation in pluripotent stem cells of humans and apes. *Nature*. 2013; 503:525–529. [PubMed: 24153179]
- Nielsen FC, Nielsen J, Kristensen MA, Koch G, Christiansen J. Cytoplasmic trafficking of IGF-II mRNA-binding protein by conserved KH domains. *Journal of Cell Science*. 2002; 115:2087–2097. [PubMed: 11973350]
- Nielsen J, Christiansen J, Lykke-Andersen J, Johnsen AH, Wewer UM, Nielsen FC. A Family of Insulin-Like Growth Factor II mRNA-Binding Proteins Represses Translation in Late Development. *Molecular and Cellular Biology*. 1999; 19:1262–1270. [PubMed: 9891060]
- Nielsen J, Kristensen MA, Willemoes M, Nielsen FC, Christiansen J. Sequential dimerization of human zipcode-binding protein IMP1 on RNA: a cooperative mechanism providing RNP stability. *Nucleic acids research*. 2004; 32:4368–4376. [PubMed: 15314207]
- Ray D, Kazan H, Cook KB, Weirauch MT, Najafabadi HS, Li X, Gueroussov S, Albu M, Zheng H, Yang A, et al. A compendium of RNA-binding motifs for decoding gene regulation. *Nature*. 2013; 499:172–177. [PubMed: 23846655]
- Ross AF, Oleynikov Y, Kislaukis EH, Taneja KL, Singer RH. Characterization of a beta-actin mRNA zipcode-binding protein. *Molecular and Cellular Biology*. 1997; 17:2158–2165. [PubMed: 9121465]
- Ross J, Lemm I, Berberet B. Overexpression of an mRNA-binding protein in human colorectal cancer. *Oncogene*. 2001; 20:6544–6550. [PubMed: 11641779]
- Sugimoto Y, Konig J, Hussain S, Zupan B, Curk T, Frye M, Ule J. Analysis of CLIP and iCLIP methods for nucleotide-resolution studies of protein-RNA interactions. *Genome biology*. 2012; 13:R67. [PubMed: 22863408]

- Van Nostrand EL, Pratt GA, Shishkin AA, Gelboin-Burkhart C, Fang MY, Sundararaman B, Blue SM, Nguyen TB, Surka C, Elkins K, et al. Robust transcriptome-wide discovery of RNA binding protein binding sites with enhanced CLIP (eCLIP). 2016 In Press.
- Vikesaa J, VOHT, Jonson L, Borup R, Weaver UM, Christiansen J, Nielsen FC. RNA-Binding IMPs promote cell adhesion and invadopodia formation. *EMBO*. 2006; 25:1456–1468.
- Wagenmakers A, Reinders RJ, Van Venrooij W. Cross-linking of mRNA to Proteins by Irradiation of Intact Cells with Ultraviolet Light. *European Journal of Biochemistry*. 1980; 112:323–330. [PubMed: 7460925]
- Wilbert ML, Huelga SC, Kapeli K, Stark TJ, Liang TY, Chen SX, Yan BY, Nathanson JL, Hutt KR, Lovci MT, et al. LIN28 binds messenger RNAs at GGAGA motifs and regulates splicing factor abundance. *Molecular cell*. 2012; 48:195–206. [PubMed: 22959275]
- Wu B, Buxbaum AR, Katz ZB, Yoon YJ, Singer RH. Quantifying Protein-mRNA Interactions in Single Live Cells. *Cell*. 2015; 162:211–220. [PubMed: 26140598]
- Yeo GW, Coufal NG, Liang TY, Peng GE, Fu XD, Gage FH. An RNA code for the FOX2 splicing regulator revealed by mapping RNA-protein interactions in stem cells. *Nature structural & molecular biology*. 2009; 16:130–137.




Figure 1. Expression patterns of IMP1, IMP2, and IMP3 RNA binding proteins
 (A) Domain structure of IMP protein family members, with RNA-Recognition Motif (RRM) 1–2, hnRNPk-homology (KH) 1–2 and 3–4 domains, and nuclear export signal (NES). (B) Illumina Bodymap tissue RNA-seq data of *IMP1-3*, *GAPDH*, and *ACTB* mRNA expression (RPKM) in comparison to H1, H9, and HUES6 human embryonic stem cells (hESCs). (C) IMP protein expression in human fibroblasts, induced pluripotent (iPS) and hESCs by Western blot analysis. (D) Immunofluorescence displaying IMP localization in hESCs, scale bar represents 10 microns. (E) Cellular fractionation into nuclear and cytoplasmic expression of IMP1–3 by Western blot analysis.




Figure 2. Identification of RNA binding targets of IMP1, IMP2, and IMP3 in hESCs by eCLIP (A) Schematic of enhanced CLIP protocol. Briefly, H9 hESCs were subjected to UV-mediated crosslinking, lysis and treatment with limiting amount of RNase, followed by immunoprecipitation (IP) of protein-RNA complexes. RNA fragments protected from RNase digestion were subjected to 3' RNA linker ligation, reverse-transcription and 3' DNA linker ligation to generate eCLIP libraries for high-throughput Illumina sequencing. In addition, a Size-Matched Input (SMInput) library was generated for each IMP protein (See Experimental Procedures for further details). (B) Identification of IMP1-bound regions. For each gene in Gencode v19, read density was counted separately for 3'UTR (blue), CDS (red), or introns (teal). Scatter plot indicates fold-enrichment for each region in IMP1 eCLIP relative to paired SMInput (y-axis), plotted against read density in SMInput (x-axis). Open circles indicate significant enrichment ($p < 10^{-5}$ and 4-fold) in eCLIP relative to SMInput. (C–F) Scatter plots indicate correlation between region-based fold enrichment in eCLIP for (C) IMP1 biological replicates, (D) IMP1 versus RBFOX2, (E) IMP1 versus IMP2 and (F) IMP1 versus IMP3. For each, regions from all genes meeting a minimal read depth criteria are shown, with least-squares regression line indicated by the dotted line. (G) Histogram of region-based fold-enrichment for IMP family members IMP1, IMP2, and IMP3, unrelated splicing regulator RBFOX2, and an IgG negative control (each compared to its paired SMInput). See also Figure S1 and S2.




Figure 3. Reproducible peaks reveal correlated binding of IMP1 and IMP2

(A) Clusters of enriched read density within IMP1 eCLIP are identified for each biological replicate by CLIPper (grey). Comparison of cluster read density in eCLIP and paired SMInput identifies a subset of clusters enriched above SMInput (red/orange; also see Figure S3B), which show high overlap between replicates (center). See Figure S3A for IMP2. (B) Irreproducible Discovery Rate (IDR) analysis comparing peak fold-enrichment across various datasets are indicated. Rep1 was used unless otherwise indicated. (C) Pie chart indicates the distribution of significantly enriched IMP1 peak locations across the transcriptome for replicating peaks described in (A). See Figure S3I for all (including non-significant) IMP1-enriched clusters, and S3J for IMP2. (D) Plot indicates the normalized read density for indicated eCLIP experiments (Rep2 for IMP1 and IMP2), centered around the midpoint of significantly enriched IMP1 peaks (described in (A)) identified from an independent biological replicate (Rep1). Read densities across each peak region were normalized within the 600nt window, and mean (points) and SEM (error bars) were calculated across all peaks for both eCLIP and paired SMInput datasets as indicated. See also Figure S3.




Figure 4. RNA Bind-N-Seq (RBNS) identifies an AC-rich preference for IMP1 and IMP2 binding (A) Schematic of RBNS protocol. (B, C) Motif logos with corresponding probability bar graphs made from aligning enriched 5-mers for IMP1 (B) and IMP2 (C). All 5-mers with an enrichment Z-score ≥ 2 with two or fewer mismatches to the most-enriched 5-mer were aligned to create top logo; remaining 5-mers with enrichment Z-score ≥ 2 were aligned to create bottom logo. Probabilities in left bar graph are proportional to the summed enrichments of all 5-mers aligned in each logo. (D, E) Comparison of 6-mer enrichments (RBNS R values) in IMP2 (D) or RBFOX2 (E) versus IMP1. 6-mers containing one of the top four non-overlapping IMP1 4-mers are colored, and significant enrichment (Z-score ≥ 2) is indicated by dotted lines. (F–H) Comparison of RBNS and eCLIP k-mer enrichment. 6-mers containing a CACA 4-mer (red), TACA 4-mer (navy), AACA 4-mer (light blue), and TACA 4-mer (magenta) are highlighted. (F–G) IMP1 RBNS enrichment of all 6-mers (x-axis) is plotted against (F) enrichment in all reproducible eCLIP 3'UTR clusters, or (G) stringent reproducible 3'UTR peaks only (as described in Figure 3A). (H) IMP2 RBNS enrichment plotted against motif enrichment in IMP2 stringent eCLIP clusters. See also Figure S4.




Figure 5. IMP1 controls Integrin RNA stability and cell adhesion in hESC
 (A) Gene ontology analysis of genes with significantly enriched IMP1 binding in their 3'UTR region in both replicates. (B) Western blot displaying levels of IMP1 depletion in the cell-adhesion assay. (C) Quantification and (D) Phase contrast images of H9 hESCs stained with Crystal Violet one hour after plating. Data are shown as mean \pm SD. (E) RNA-seq analysis of integrin RNA expression changes following loss of IMP1 in hESC. (F) eCLIP 3'UTR binding (\log_2 fold-enrichment over SMInput) for the integrins shown in (E). (G) Read density tracks show read density for IMP1 (Rep2; red), RBFOX2 (blue), and paired SMInputs (gray) across the 3'UTR of *ITGB5*. Clusters are indicated as boxes below tracks, with significantly enriched peaks indicated as darkly colored boxes. (H) RNA immunoprecipitation for *ITGB5* in HUES6 hESCs. *ACTB* is shown as a positive control. (I) qRT-PCR validation of *ITGB1* and *ITGB5* expression changes by RNA-seq, normalized to *HMBS*, *GAPDH* and *18s* (Error bars = mean \pm SEM, N=3). (J) *ITGB1* and *ITGB5* protein levels following depletion of IMP1 in H9 hESCs. An average of three independent experiments quantified by densitometry is shown on the right. (K) ActinomycinD RNA stability assay. Expression of *ITGB5* was measured by qRT-PCR at different time points following addition of 10 μ M Actinomycin D in H9 hESCs and normalized to *PPIA* and *RPLP0*, genes determined not to change over the given time period (Error bars = mean \pm SEM, N=3; A single asterisk indicates significance of $p < 0.05$ by unpaired *t*-test). See also Figure S5.




Figure 6. IMP1 promotes cell survival through regulation of BCL2

(A) Phase contrast images of H9 hESCs following depletion of IMP1. Scale bar represents 1mm. (B) Shown is a representative image of flow cytometry analysis and quantification of AnnexinV expression following depletion of IMP1 (Error bars = mean \pm SD, N=3). (C) Cleaved-Caspase 3 and BCL2 protein expression following depletion of IMP1 in H9 hESCs by Western blot. (D) qRT-PCR validation of *BCL2* expression following depletion of IMP1, with expression normalized to average of *HMBS*, *GAPDH* and *18s* (Error bars = mean \pm SEM, N=3). (E) RNA immunoprecipitation (RIP) for *BCL2* in HUES6 hESCs using antibodies that target either IMP1 or IgG. (F) EMSA results depicting IMP1 binding preferences for *ACTB* WT, *ACTB* truncated, *BCL2* WT, and *BCL2* RNAs. One representative experiment is shown. (G) Caspase-Glo Apoptosis assay. BCL2 protein was induced with increasing amounts of doxycycline after which apoptosis was measured via luminescence using the Caspase-Glo 3/7 assay. A single asterisk indicates significance of $p < 0.05$, and two asterisks indicate significance of $p < 0.01$, determined by unpaired t -test. Data are shown as mean \pm SD. See also Figure S6.