Lawrence Berkeley National Laboratory

Recent Work

Title

THE SPINS OF SILVER-105, SILVER-106, AND SILVER-110m

Permalink

<https://escholarship.org/uc/item/9j81n100>

Authors

Ewbank, W.B.
Nierenberg, W.A.
Shugart, H.A.
et al.

Publication Date

1958-01-16

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

THE SPINS OF SILVER-105, SILVER-106, AND SILVER-110m
W.B. Ewbank, W.A. Nierenberg, H.A. Shugart, and H.B. Silsbee
January 16, 1958

THE SPINS OF SILVER-105, SILVER-106, AND SILVER-110m

W.B. Ewbank, W.A. Nierenberg, H.A. Shugart, and H.B. Silsbee

**Department of Physics and Radiation Laboratory
University of California, Berkeley, California**

January 16, 1958

ABSTRACT

The spins of three radioactive silver isotopes have been measured by the atomic-beam magnetic-resonance method. The results are: for 40-day Ag^{105} , spin $\frac{1}{2}$; for 8.3-day Ag^{106} , spin 6; for 253-day $\text{Ag}^{110\text{m}}$, spin 6.

THE SPINS OF SILVER-105, SILVER-106, AND SILVER-110m*

W. B. Ewbank, W. A. Nierenberg, † H. A. Shugart, and H. B. Silsbee

Department of Physics and Radiation Laboratory
University of California, Berkeley, California

January 16, 1958

EXPERIMENTAL METHOD

The spins of three radioactive silver isotopes, Ag^{105} ($T_{1/2} = 40$ days, ¹ $I = \frac{1}{2}$), Ag^{106} ($T_{1/2} = 8.3$ days, ² $I = 6$), and Ag^{110m} ($T_{1/2} = 253$ days, ³ $I = 6$), have been measured by atomic-beam techniques similar to those previously described.⁴

Ag^{105} and Ag^{106} were produced in the Berkeley 60-inch Crocker cyclotron, both by the reaction $\text{Rh}(\alpha, \text{kn})\text{Ag}$ and by $\text{Pd}(p, n)\text{Ag}$. After rhodium bombardments, the target material was simply transferred to the atomic-beam oven and heated until atomic silver diffused out at a satisfactory rate. Difficulty was sometimes encountered in getting a sufficiently intense beam without melting the rhodium. If the rhodium melts, the silver comes out too rapidly for use, and the rhodium flows into the oven slits and plugs them. The same technique was used for some palladium targets, but chemical separation of the silver proved more satisfactory. The target was dissolved and the active silver (plus stable carrier) was precipitated as AgCl . This was washed and dissolved in ammonium hydroxide. Then metallic silver was recovered by electroplating and transferred to the atomic beam apparatus.

Ag^{110m} was produced in a 4-week neutron bombardment of natural silver in the Arco reactor. The silver target evidently contained appreciable mercury, since simple qualitative analysis using radioactive detection showed the presence of 46-day Hg^{203} . The mercury tended to come from the atomic beam oven in

*Research supported in part by the United States Atomic Energy Commission and the Office of Naval Research.

†Professor in the Miller Institute for Basic Research in Science.

¹J. R. Gum and M. L. Pool, Phys. Rev. 80, 315 (1950).

²Gum, Thompson, and Pool, Phys. Rev. 76, 184 (1949).

³K. W. Geiger, Phys. Rev. 105, 1539 (1957).

⁴Hobson, Hubbs, Nierenberg, Silsbee, and Sunderland, Phys. Rev. 104, 101 (1956).

bursts and thus to give occasional spurious signals--which could be identified, however, by a crude analysis of their γ -ray spectrum.

Since the stable silver carrier is not readily detectable, it could not be used to monitor the intensity of the atomic beam. Instead, the radioactive beam was collected for short periods when the stops in the apparatus were removed. Under such conditions, the fast tail of the Maxwell distribution reaches the collector position in spite of the deflecting magnets, and the counting rates from such "half-beam" exposures were used to normalize spin counting rates.

A small amount of an appropriate alkali compound, e.g. CsCl, was included with the radioactive material in the oven. Surface-ionization detection of this material facilitated lining up of the beam apparatus at the beginning of the run.

Calibration of the transition magnetic field was accomplished by observing the resonances of an atomic alkali beam from a second oven located behind the radioactive one. The latter was simply moved to one side to permit the observations.

SILVER-105

Table I indicates the results of the first spin search for Ag^{105} . There is a large signal corresponding to spin $\frac{1}{2}$. There is also an above-background rate at spin $3/2$, because the frequency appropriate to spin $3/2$ has for its second harmonic the spin- $\frac{1}{2}$ frequency. This spurious signal disappeared when proper precautions were taken.

Table I. Results of spin search for Ag^{105} .

Spin	Counting rate (cpm)	Spin	Counting rate (cpm)
0	0.15 ± 0.05	3	0.07 ± 0.07
$1/2$	4.8 ± 0.2	4	0.5 ± 0.1
1	0.15 ± 0.05	5	0.4 ± 0.1
$3/2$	1.4 ± 0.1	6	2.0 ± 0.2
2	0.10 ± 0.07	7	0.5 ± 0.1
$5/2$	0.10 ± 0.05		

Figure 1 compares the decay of a spin- $\frac{1}{2}$ exposure with that of a full-beam sample from the same rhodium bombardment. The latter shows a mixture of 40-day Ag^{105} activity with shorter-lived material (Ag^{106}), whereas the spin button shows a clean 40-day decay indicating a significant enrichment in Ag^{105} . The spin- $\frac{1}{2}$ signal shown in Fig. 2 is observed from activity produced by proton bombardment of palladium.

This isotope has been observed repeatedly at a variety of frequencies from 7.4 Mc to 15.4 Mc. No significant quadratic Zeeman shift was observable at these frequencies with the present line width.

SILVER-106

Table I (from a rhodium bombardment) gives a hint that the Ag^{106} activity has spin 6. This was confirmed by the results of a later palladium bombardment, as indicated in Table II. The decay of the spin-6 sample is compared in Fig. 2 with that of the full beam, and clearly shows enrichment of the 8.3-day activity. No quadratic Zeeman shift was observed between 2.4 and 15.7 Mc.

Table II. Results of spin search for Ag^{106} .

Spin	Counting rate (cpm)	Spin	Counting rate (cpm)
$\frac{1}{2}$	9.94 ± 0.22	6	4.86 ± 0.16
5	0.33 ± 0.09	7	0.56 ± 0.09

SILVER-110m

Results of a spin search on the Arco material are indicated in Table III. There is a clear indication of spin 6, which must be attributed to 253-day Ag^{110m} , since no other activity of comparable life can be produced in such a bombardment. The long-term stability of the counters may not be adequate for a reliable determination of the half-life; however, a check of two spin-6 buttons after 300 days showed a decrease in measured counting rates by a factor 2.1, which may be regarded as consistent with a 253-day half-life. Signals corresponding to spin 6 were observed at seven frequencies, from 2.4 Mc to 15.7 Mc; the quadratic Zeeman term, if any is small.

Table III. Results of spin search for $\text{Ag}^{110\text{m}}$.

Spin	Counting rate (cpm)	Spin	Counting rate (cpm)
1	2.8 ± 0.3	5	2.6 ± 0.3
2	1.7 ± 0.3	6	31.1 ± 0.5
3	2.2 ± 0.3	7	2.0 ± 0.3
4	1.7 ± 0.3	8	2.7 ± 0.2

REMARKS

The result, spin $\frac{1}{2}$ for Ag^{105} , is not surprising, since the stable isotopes Ag^{107} and Ag^{109} also have this spin; it is also consistent with the simple shell model⁵ that calls for the odd proton to be in a $p_{1/2}$ level, although the $g_{9/2}$ level lies close. The odd neutron in Ag^{106} and $\text{Ag}^{110\text{m}}$ is expected to lie in a $d_{5/2}$ state. (However, the ground state of Cd^{111} , with the same neutron-number as $\text{Ag}^{110\text{m}}$, exhibits spin $\frac{1}{2}$; the first excited state, spin $5/2$.) The observed spins of 6 could thus be attributed to the coupling of a $g_{9/2}$ proton with a $d_{5/2}$ neutron to give one less than the maximum spin, in agreement with Nordheim's "weak" rule, although other possibilities exist. In any event, one would expect a large positive magnetic moment, consistent with our failure to observe appreciable quadratic Zeeman terms.

⁵M. G. Mayer and J. H. D. Jensen, *Elementary Theory of Nuclear Shell Structure* (Wiley, New York, 1955), pp. 71-92.

FIGURE CAPTIONS

Fig. 1. The decay of spin- $\frac{1}{2}$ sample from a rhodium bombardment is compared with a full-beam sample. The essentially pure half-life of the spin- $\frac{1}{2}$ sample serves to identify Ag^{105} as responsible for the resonance signal.

Fig. 2. The decay of an $I = 6$ sample from a palladium bombardment identifies Ag^{106} as responsible for the spin 6 resonance. The Ag^{105} spin- $\frac{1}{2}$ resonance is observed by use of a different method of production from that shown in Fig. 1.

1744-2