

UCSF

UC San Francisco Previously Published Works

Title

Low Dose Carbon Monoxide Exposure in Idiopathic Pulmonary Fibrosis Produces a CO Signature Comprised of Oxidative Phosphorylation Genes.

Permalink

<https://escholarship.org/uc/item/9jc7f9m3>

Journal

Scientific Reports, 9(1)

Authors

Casanova, Nancy
Zhou, Tong
Gonzalez-Garay, Manuel
et al.

Publication Date

2019-10-15

DOI

10.1038/s41598-019-50585-3

Peer reviewed

OPEN

Low Dose Carbon Monoxide Exposure in Idiopathic Pulmonary Fibrosis Produces a CO Signature Comprised of Oxidative Phosphorylation Genes

Nancy Casanova¹, Tong Zhou², Manuel L. Gonzalez-Garay¹, Ivan O. Rosas³, Hilary J. Goldberg³, Stefan W. Ryter¹⁰, Harold R. Collard⁴, Souheil El-Chemaly³, Kevin R. Flaherty⁷, Gary M. Hunninghake³, Joseph A. Lasky⁸, David J. Lederer⁹, Roberto F. Machado⁵, Fernando J. Martinez¹⁰, Imre Noth¹¹, Ganesh Raghu⁶, Augustine M. K. Choi¹⁰ & Joe G. N. Garcia¹

Compelling preclinical studies indicate that low-dose carbon monoxide (CO) abrogates experimental lung fibrosis. We recently reported the results of a multicenter, double-blinded, clinical trial of inhaled CO in patients with idiopathic pulmonary fibrosis (IPF). Identifying no significant changes in metalloproteinase-7 (MMP7) serum concentration, or secondary endpoints of physiologic measurements, hospitalization, death, or patient-reported outcomes. In the present study, we evaluated the effect of low dose CO exposure (100–200 ppm) for 12 weeks on genome-wide gene expression in peripheral blood mononuclear cells (PBMC) derived from these IPF study subjects. We conducted transcriptome profiling on 38 IPF subjects with time points available at 0, 12, and 24 weeks. Total RNA isolated from PBMCs was hybridized onto the Affymetrix Human Gene 2.0 ST Array. We identified 621 genes significantly upregulated in the 24-week CO exposed group compared with the 12-week. Pathway analysis demonstrated association with Oxidative Phosphorylation (adjusted $P < 0.05$). We identified a clear CO signature dominated with 23 oxidative phosphorylation-related genes (FDR $< 10\%$). We confirmed the expression of nine selected gene products using Nanostring's nCounter analysis system. These findings suggest this signature may serve as a potential genomic biomarker for CO exposure and for potential titration of dosage to allow precision testing of therapies in future low dose CO therapeutic studies in IPF.

Idiopathic pulmonary fibrosis (IPF) is a relentlessly progressive interstitial lung disease characterized by the excessive formation of scar tissue in the absence of any known provocation¹. IPF patients typically experience a progressive decline in lung function leading to a fatal respiratory failure². In the absence of treatment, IPF is usually fatal within 2–3 years of the onset of symptoms. Lung transplantation is the only cure for IPF patients, but

¹Department of Medicine, University of Arizona Health Sciences, Tucson, AZ, USA. ²Department of Physiology and Cell Biology, University of Nevada School of Medicine, Reno, NV, USA. ³Division of Pulmonary and Critical Care Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA. ⁴Division of Pulmonary and Critical Care Medicine, University of California San Francisco, San Francisco, CA, USA. ⁵Division of Pulmonary, Critical Care, Indiana University, Indianapolis, IN, USA. ⁶Division of Pulmonary and Critical Care Medicine, University of Washington Medical Center, Seattle, WA, USA. ⁷Division of Pulmonary and Critical Care Medicine, University of Michigan, Ann Arbor, MI, USA. ⁸Pulmonary and Critical Care Medicine Section, Tulane University Medical School, New Orleans, LA, USA. ⁹Division of Pulmonary and Critical Care Medicine, Columbia University Medical Center, New York, NY, USA. ¹⁰Department of Medicine, Weill Cornell Medical College, New York, NY, USA. ¹¹Division of Pulmonary and Critical Care Medicine, University of Chicago, Chicago, IL, USA. Correspondence and requests for materials should be addressed to J.G.N.G. (email: skipgarcia@email.arizona.edu)

many IPF patients expire before receiving a lung transplant and only 20% to 30% of IPF patients survive 5 years after diagnosis. The approvals of Pirfenidone and Nintedanib^{3,4} provides additional, much needed therapeutic options. However, 20% of IPF patients discontinue treatment as a consequence of adverse events, and the high cost of these new drugs prohibit their wider use⁵. More sobering yet is that these costly drugs have a modest impact on disease progression and survival. Thus, IPF remains an incurable disease with a dismal prognosis and there is a continuing search for better tolerated, safer and economical treatments.

The pleiotropic biological functions of carbon monoxide (CO) include protection against oxidative injury^{6,7}, inhibition of cell proliferation⁸, suppression of matrix production⁹, repression of fibrinolysis¹⁰, modulation of apoptosis, and inflammation, and protection against other environmental insults^{11–16}. CO can bind to hemoproteins resulting in modulation signal transduction pathways affecting gene regulation that result in significant reduction of processes associated with the pathogenesis of lung fibrosis. Numerous pre-clinical studies have examined the protective effect of low CO concentrations on the lung parenchyma and vasculature. Mice exhibit an increased tolerance to lethal concentrations of oxygen (hyperoxia) with increased survival and attenuation of lung injury¹². CO provides lung protection for lung transplantation¹⁷, aeroallergen-induced inflammation¹⁸ and lethal ischemic lung injury¹⁰. Low levels of CO suppress bleomycin-induced lung fibrosis and CO-exposed cells displayed impaired production of extracellular matrix proteins such as fibronectin and collagen⁹. CO (100–125 ppm) was well tolerated in patients with chronic obstructive pulmonary disease (COPD) and was shown to reduce sputum eosinophilia and improves methacholine responsiveness¹⁹. CO has antivaso-occlusive and immunomodulatory effects beneficial to sickle cell disease patients to prevent cardiovascular complications²⁰.

The use of transcriptomic data to characterize biological effects of small molecules has become popular in drug discovery projects²¹ to determine biological effects of a drug at gene expression level. We recently completed a multicenter, double-blinded, clinical trial of inhaled CO in IPF which demonstrated the safety and tolerability of low doses (100–200 ppm) of CO in IPF²². However, despite modest increases in carboxy-hemoglobin (CO-Hb) blood levels, low dose CO exposure failed to significantly affect the primary study endpoint of changes in metalloproteinase-7 (MMP7) serum concentration, or secondary endpoints of physiologic measurements, hospitalization, death, or patient-reported outcomes. The aim of the present study was to determine the effect of CO exposure on genome-wide gene expression in peripheral blood mononuclear cells from room air (RA) and CO-exposed IPF study subjects.

Methods

Study overview. We conducted a multicenter phase II randomized, double blind, placebo-controlled of clinical trial of inhaled CO compared to placebo, which was approved by each participating center's Institutional Review Board (Brigham and Women's IRB #210P001676). Research was performed in accordance with relevant guidelines and regulations. Written informed consent was obtained from all subjects enrolled at Brigham and Women's Hospital-Harvard Medical School, University of Illinois, University of Chicago, University of California San Francisco, University of Michigan, Tulane University, Columbia University, and University of Washington. Participants had IPF diagnosed according to ATS/ERS/ALAT guidelines for IPF diagnosis and management², mild to moderate lung disease (FVC greater than or equal to 50% predicted). Inclusion/exclusion criteria, randomization, carbon monoxide dosing and administration were previously described²². Briefly, subjects were randomized to inhaled CO treatment with an initial dose of 100 ppm for one week followed by dose escalation to 200 ppm, or to placebo with RA (21% inhaled oxygen). Inhaled CO was administered twice weekly for a total of 12 weeks. Peripheral blood samples were collected on specified research visits. A total of 51 subjects completed 12 weeks of randomized therapy (CO or placebo) and 45 subjects completed follow-up period. Complete enrolled cohort baseline demographics, imaging, biopsy findings and pulmonary function testing of randomized subjects were balanced; statistically significant differences were not noted between subjects treated with CO compared to treatment with placebo for any element²². We excluded 9 subjects in the CO arm and 11 in the RA due to early termination or sample issues, additional details are depicted in Fig. 1. There was no clinical significant differences between these and the subjects included in the gene expression analysis.

Genomic analysis of CO trial participants. For this genomic analysis, we assessed the treatment effects in IPF subjects with at base line (week 0), end of treatment dosing period (week 12), and follow up visit (week 24). We included the subjects with isolated mononuclear cells available on these time points PBMC isolation was performed on each of the centers using the Ficoll-Paque isolation method; samples were stored at the BWH-HMS and transferred to the University of Arizona where the total RNA was isolated from PBMCs using RNeasy MiniKit Qiagen™ following manufacturer's protocol. RNA concentration and quality (RIN > 7) was assayed by Nanodrop™ (Thermo Fisher) and 2100 Bioanalyzer RNA™ (Agilent).

Transcriptome profiling. RNA was hybridized onto the Affymetrix GeneChip® Human Gene 2.0 ST Array (Thermo Fisher Scientific), to conduct transcriptome profiling. The expression microarrays were analyzed using the Affymetrix Power Tools v.1.15.1. Probeset expression signals were summarized with the robust multi-array average (RMA) algorithm²³ and log₂ transformed with a median polish. Transcripts were considered to be reliably expressed in the samples if the Affymetrix implemented DABG (detection above ground) *P*-value was less than 0.01 in at least 67% of the samples. SAM (Significance Analysis of Microarrays)²⁴, implemented in the *samr* library of the R Statistical Package, was used to compare log₂-transformed gene expression levels between 0-week and 12-week samples and between 12-week and 24-week samples, respectively. False discovery rate (*FDR*) was controlled using the *q*-value method²⁵. Transcripts with *FDR* less than 10% were deemed differentially expressed. We searched for any enriched Kyoto Encyclopedia of Genes and Genomes (KEGG)²⁶ physiological pathways among the differential genes relative to the final analysis set using the NIH/DAVID²⁷. An adjusted *P*-value < 0.05 after the *Benjamini-Horchberg* procedure was used as the cutoff.

CONSORT 2010 Flow Diagram




Figure 1. Flowchart of total study enrollment and subjects included in transcriptome and validation analysis.

Computing geneset scores using human transcriptomic data. The Functional Analysis of Individual Microarray Expression (FAIME) algorithm was applied to assign a geneset score for each KEGG pathway²⁸. FAIME computes geneset scores using rank-weighted gene expression of individual samples, which provides a translation of each sample's transcriptomic information to molecular mechanisms²⁸. Higher geneset score indicates overall upregulation of a given KEGG pathway.

Validation. Transcriptomic results obtained by microarray were validated in the 18 subjects from the CO arm using the NanoString's nCounter[®] analysis system (NanoString Technologies, Seattle, WA, USA)²⁹. We selected oligonucleotide probes for molecular barcoding of the nine microarray-selected genes and three housekeeping genes (OAZ1, RPL24, RPS29). After hybridization and magnetic bead purification, barcodes were counted for each target molecule. Following normalization, we calculated the fold-change for the genes of interest. Using R Statistical Package, one tailed paired t-test (P-value < 0.05), was used to compare gene expression levels between 0-week and 12-week samples and between 12-week and 24-week samples, respectively.

Characteristics	CO (n = 20)	Room Air (n = 18)	P value
Age	65 ± 6.9	65.8 ± 8.2	0.81
Gender (Male/Female)	17(85)/3(15)	13(72)/5(28)	0.48
Race/ Ethnicity			0.11
White	15	16	
Black	2	0	
Asian	3	0	
Latino	0	2	
PFT			
FVC, L	3.08 ± 0.65	2.92 ± 0.96	0.435
FVC % predicted	75.6 ± 17.1	69.05 ± 13.6	0.147
TLC % predicted	70.4 ± 11	66.23 ± 13.6	0.305
Dlco % predicted	44.2 ± 11.9	41.06 ± 15.7	0.26

Table 1. Baseline demographics and pulmonary function testing by study arm. Values are No. (%), mean ± SD, or as otherwise indicated. CO = carbon monoxide; Dlco = diffusing capacity for carbon monoxide; FVC = forced vital capacity; TLC = total lung capacity.

Results

Human subjects. We conducted transcriptome analysis and validation analysis on 38 from the original 58 subjects enrolled in the clinical trial, selected based upon study completion, sample availability and sample quality Fig. 1. 20 subjects were randomized in the CO arm and 18 subjects received RA. Baseline demographics and pulmonary function testing results are outlined in Table 1. No significant differences on baseline demographic and clinical characteristics were present in the two study arms.

Differential gene expression induced by CO exposure. At the specified significance level ($FDR < 10\%$), we failed to identify any differentially-expressed genes between the 0-week and 12-week groups and between the 0-week and the 24 week under CO treatment. However, 621 genes were significantly upregulated in the 24-week CO exposed group compared with the 12-week CO samples (Fig. 2A). The increase at week 24 represented a normalization to baseline values. Pathway analysis based on the KEGG database demonstrated that the top KEGG pathway associated with the 621 upregulated genes is “Oxidative Phosphorylation” (adjusted $P < 0.05$) (Fig. 2B). In addition, we found that due to the large proportion of overlapped genes ($>40\%$) with the “Oxidative Phosphorylation” pathway; the “Parkinson’s disease”, “Huntington’s disease”, and “Alzheimer’s disease” pathways were also significantly enriched by the upregulated genes (adjusted $P < 0.05$) (Fig. 2B). We further computed the “Oxidative Phosphorylation” pathway score (see Methods for details) for both the CO and RA samples at each time point. Paired comparison revealed that for the patients with CO treatment, the “Oxidative Phosphorylation” pathway score is significantly lower at 12-week compared with 0-week (paired t -test: $P = 0.035$) (Fig. 2C), while this pathway is significantly upregulated at 24-week relative to 12-week (paired t -test: $P = 0.009$) (Fig. 2C). These results suggest that the dysregulation caused by CO treatment is remarkably recovered by discontinuation of CO treatment. As expected, patients with RA treatment did not demonstrate any significant difference in the “Oxidative Phosphorylation” pathway score among the three time points (Fig. 2C).

Oxidative phosphorylation pathway-based gene signature. To understand the CO-induced gene dysregulation pattern, we developed a gene signature based on the genes within the “Oxidative phosphorylation” pathway. Only the genes differentially expressed between 12-week and 24-week for the CO treated patients ($FDR < 10\%$) were retained. In total, a gene signature with 23 oxidative phosphorylation-related genes was identified (Fig. 3).

Candidate gene biomarkers confirmed by nanostring. We confirmed the expression of nine selected gene products using Nanostring in 18 subjects from the same cohort of CO treated patients (Fig. 4). All nine genes were downregulated at 12-week compared with 0-week (one-tailed paired t -test: $P < 0.05$). Two genes, CYC1 and NDUFA6, were significantly upregulated at 24-week compared with 12-week (one-tailed paired t -test: $P < 0.05$). In the RA control group, five of the 11 genes were confirmed by Nanostring at nominal p -value < 0.05 .

Discussion

The present study identified changes in gene expression in PBMCs of Idiopathic pulmonary fibrosis patients following exposure to low dose (100–200 ppm) for 12 weeks using the Affymetrix microarray platform. A validation study was conducted in the same cohort of patients. Despite the absence of significant difference in the pulmonary function testing and functional clinical assessment in the CO-receiving group²², we detected significant dysregulation of gene expression in subjects receiving CO twice a week. We analyzed gene expression levels between 0-week and 12-week samples and between 12-week and 24-week samples, and 0 week and 24 week respectively. Significant differential expression was observed at week 12 and week 24 with gene dysregulation response abolished after treatment termination. The differentially-expressed genes were dominantly enriched in mitochondrial-related signaling pathways with oxidative phosphorylation, Parkinson’s disease and Huntington’s disease pathways representing the top dysregulated pathways during CO administration.




Figure 2. (A) Means of expression (log₂) comparison of 621 genes significantly upregulated at 12 week and 24-week in the CO-exposed cohort. (B) Oxidative phosphorylation was the top KEGG pathway associated with the 621 upregulated genes (adjusted $P < 0.05$). A proportion of genes (40%) in this pathway overlapped with Parkinson's disease, Huntington's disease and Alzheimer's disease pathways (adjusted $P < 0.05$). (C) Oxidative Phosphorylation pathway score according to FAIME algorithm indicates that the dysregulation caused by CO treatment recovers following discontinuation of CO treatment (24 wk).




Figure 3. The 23 oxidative phosphorylation-related CO gene signature. Only genes within the Oxidative phosphorylation with differential expression between 12-week and 24-weeks in the CO-treated patients (FDR < 10%) were retained. In total, a gene signature with 23 oxidative phosphorylation-related genes was identified.

There is evidence of mitochondrial dysfunction in IPF includes a decreased efficiency of electron transport and increases in reactive oxygen species (ROS) production³⁰. Oxidative phosphorylation is the coordinated transfer of electrons and protons leading to the ATP production. The mitochondrial electron transport chain (ETC) is vital for cellular energy production, but may serve as a source of aberrant ROS production from ETC complexes I and III during pathophysiological states. Complex I (NADH dehydrogenase) is a site of ROS leakage under pathophysiological conditions. CO inhibits Toll-like receptor signaling by suppressing the ROS generated through




Figure 4. Nanostring validation confirmed the downregulation at 12-week of nine selected microarray-derived gene products (p-value < 0.05).

NADPH oxidase activation²⁹. Mitochondria heme functional groups have been strongly implicated as a primary targets of CO action³¹. Notably, we identified that CO therapy decreased the expression of genes encoding ETC complex I genes: NDUFB4, NDUFS2, NDUFAB1, NDUFA6, NDUFB9, also enriched in Parkinson's, Huntington's and Alzheimer disease pathways. Similarly, complex IV, cytochrome c oxidase (COX) subunits: COX4I1 and COX7A2 within the terminal enzyme respiratory chain appear to be modulated by CO exposure with a downregulation that appear to return to baseline at week 24. CYC1, from the complex III, an ubiquinol-cytochrome c reductase complex subunit, also followed the same dysregulation pattern.

In the present study, we were able to identify the downregulation of gene expression at the mitochondrial level secondary to CO modulation at COX and NADH levels. These findings raise the possibility that CO exposure in IPF, targets modulation of mitochondrial energy production and potentially impacts mitochondrial ROS production.

Despite the reproducibility of our findings, we recognize the limitations of our study. Additional dose-titration studies are required to determine the CO dose and duration that can evoke changes on other clinical markers of disease progression. Despite these limitations, similar to other molecular profiling strategies designed to subphenotype patients with chronic lung disease^{32,33}, our study suggests that a microarray-derived molecular signature successfully identifies the transcriptomic profile of CO in IPF patients dominated by significant dysregulation of genes in the oxidative phosphorylation pathway. These findings demonstrate interesting transcriptional effects associated to the CO administration and suggest this signature may serve as a potential genomic biomarker for CO exposure that can be used for dose stratification to potentially allow future precision testing of low dose CO therapies in IPF.

References

1. Travis, W. D. *et al.* An official American Thoracic Society/European Respiratory Society statement: Update of the international multidisciplinary classification of the idiopathic interstitial pneumonias. *Am J Respir Crit Care Med* **188**, 733–748, <https://doi.org/10.1164/rccm.201308-1483ST> (2013).
2. Raghu, G. *et al.* An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med* **183**, 788–824, <https://doi.org/10.1164/rccm.2009-040GL> (2011).
3. Karimi-Shah, B. A. & Chowdhury, B. A. Forced vital capacity in idiopathic pulmonary fibrosis—FDA review of pirfenidone and nintedanib. *N Engl J Med* **372**, 1189–1191, <https://doi.org/10.1056/NEJMp1500526> (2015).
4. Noble, P. W. *et al.* Pirfenidone in patients with idiopathic pulmonary fibrosis (CAPACITY): two randomised trials. *Lancet* **377**, 1760–1769, [https://doi.org/10.1016/S0140-6736\(11\)60405-4](https://doi.org/10.1016/S0140-6736(11)60405-4) (2011).
5. Fletcher, S. *et al.* The safety of new drug treatments for idiopathic pulmonary fibrosis. *Expert Opin Drug Saf* **15**, 1483–1489, <https://doi.org/10.1080/14740338.2016.1218470> (2016).
6. Otterbein, L. E. & Choi, A. M. Heme oxygenase: colors of defense against cellular stress. *Am J Physiol Lung Cell Mol Physiol* **279**, L1029–1037 (2000).

7. Otterbein, L. E. *et al.* MKK3 mitogen-activated protein kinase pathway mediates carbon monoxide-induced protection against oxidant-induced lung injury. *Am J Pathol* **163**, 2555–2563, [https://doi.org/10.1016/S0002-9440\(10\)63610-3](https://doi.org/10.1016/S0002-9440(10)63610-3) (2003).
8. Song, R. *et al.* Carbon monoxide inhibits human airway smooth muscle cell proliferation via mitogen-activated protein kinase pathway. *Am J Respir Cell Mol Biol* **27**, 603–610, <https://doi.org/10.1165/rcmb.4851> (2002).
9. Zhou, Z. *et al.* Carbon monoxide suppresses bleomycin-induced lung fibrosis. *Am J Pathol* **166**, 27–37, [https://doi.org/10.1016/S0002-9440\(10\)62229-8](https://doi.org/10.1016/S0002-9440(10)62229-8) (2005).
10. Fujita, T. *et al.* Paradoxical rescue from ischemic lung injury by inhaled carbon monoxide driven by derepression of fibrinolysis. *Nat Med* **7**, 598–604, <https://doi.org/10.1038/87929> (2001).
11. Morita, T., Mitsialis, S. A., Koike, H., Liu, Y. & Kourembanas, S. Carbon monoxide controls the proliferation of hypoxic vascular smooth muscle cells. *J Biol Chem* **272**, 32804–32809 (1997).
12. Otterbein, L. E., Mantell, L. L. & Choi, A. M. Carbon monoxide provides protection against hyperoxic lung injury. *Am J Physiol* **276**, L688–694 (1999).
13. Brouard, S. *et al.* Carbon monoxide generated by heme oxygenase 1 suppresses endothelial cell apoptosis. *J Exp Med* **192**, 1015–1026 (2000).
14. Otterbein, L. E. *et al.* Carbon monoxide has anti-inflammatory effects involving the mitogen-activated protein kinase pathway. *Nat Med* **6**, 422–428, <https://doi.org/10.1038/74680> (2000).
15. Morse, D. & Choi, A. M. Heme oxygenase-1: from bench to bedside. *Am J Respir Crit Care Med* **172**, 660–670, <https://doi.org/10.1164/rccm.200404-465SO> (2005).
16. Ryter, S. W., Alam, J. & Choi, A. M. Heme oxygenase-1/carbon monoxide: from basic science to therapeutic applications. *Physiol Rev* **86**, 583–650, <https://doi.org/10.1152/physrev.00011.2005> (2006).
17. Song, R. *et al.* Carbon monoxide induces cytoprotection in rat orthotopic lung transplantation via anti-inflammatory and anti-apoptotic effects. *Am J Pathol* **163**, 231–242, [https://doi.org/10.1016/S0002-9440\(10\)63646-2](https://doi.org/10.1016/S0002-9440(10)63646-2) (2003).
18. Chapman, J. T., Otterbein, L. E., Elias, J. A. & Choi, A. M. Carbon monoxide attenuates aeroallergen-induced inflammation in mice. *Am J Physiol Lung Cell Mol Physiol* **281**, L209–216 (2001).
19. Bathoorn, E. *et al.* Anti-inflammatory effects of inhaled carbon monoxide in patients with COPD: a pilot study. *Eur Respir J* **30**, 1131–1137, <https://doi.org/10.1183/09031936.00163206> (2007).
20. Gomperts, E. *et al.* The role of carbon monoxide and heme oxygenase in the prevention of sickle cell disease vaso-occlusive crises. *Am J Hematol* **92**, 569–582, <https://doi.org/10.1002/ajh.24750> (2017).
21. Verbist, B. *et al.* Using transcriptomics to guide lead optimization in drug discovery projects: Lessons learned from the QSTAR project. *Drug Discov Today* **20**, 505–513, <https://doi.org/10.1016/j.drudis.2014.12.014> (2015).
22. Rosas, I. O. *et al.* A Phase II Clinical Trial of Low-Dose Inhaled Carbon Monoxide in Idiopathic Pulmonary Fibrosis. *Chest* **153**, 94–104, <https://doi.org/10.1016/j.chest.2017.09.052> (2018).
23. Irizarry, R. A. *et al.* Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* **4**, 249–264, <https://doi.org/10.1093/biostatistics/4.2.249> (2003).
24. Tusher, V. G., Tibshirani, R. & Chu, G. Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci USA* **98**, 5116–5121, <https://doi.org/10.1073/pnas.091062498> (2001).
25. Tibshirani, R., Hastie, T., Narasimhan, B. & Chu, G. Diagnosis of multiple cancer types by shrunken centroids of gene expression. *Proc Natl Acad Sci USA* **99**, 6567–6572, <https://doi.org/10.1073/pnas.082099299> (2002).
26. Kanehisa, M., Goto, S., Kawashima, S., Okuno, Y. & Hattori, M. The KEGG resource for deciphering the genome. *Nucleic Acids Res* **32**, D277–280, <https://doi.org/10.1093/nar/gkh063> (2004).
27. Dennis, G. Jr. *et al.* DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol* **4**, P3 (2003).
28. Yang, X. *et al.* Single sample expression-anchored mechanisms predict survival in head and neck cancer. *PLoS Comput Biol* **8**, e1002350, <https://doi.org/10.1371/journal.pcbi.1002350> (2012).
29. Geiss, G. K. *et al.* Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nat Biotechnol* **26**, 317–325, <https://doi.org/10.1038/nbt1385> (2008).
30. Zanj, D. C., Bueno, M., Mora, A. L. & Rojas, M. Idiopathic Pulmonary Fibrosis: Aging, Mitochondrial Dysfunction, and Cellular Bioenergetics. *Frontiers in medicine* **5**, 10, <https://doi.org/10.3389/fmed.2018.00010> (2018).
31. Ryter, S. W., Ma, K. C. & Choi, A. M. K. Carbon monoxide in lung cell physiology and disease. *Am J Physiol Cell Physiol* **314**, C211–C227, <https://doi.org/10.1152/ajpcell.00022.2017> (2018).
32. Zhou, T. *et al.* Peripheral blood gene expression as a novel genomic biomarker in complicated sarcoidosis. *PLoS One* **7**, e44818, <https://doi.org/10.1371/journal.pone.0044818> (2012).
33. Herazo-Maya, J. D. *et al.* Peripheral blood mononuclear cell gene expression profiles predict poor outcome in idiopathic pulmonary fibrosis. *Sci Transl Med* **5**, 205ra136, <https://doi.org/10.1126/scitranslmed.30059645/205/205ra136> (2013).

Acknowledgements

This work was supported by NHLBI grant 5U01HL105371–03 and SA 106476. The authors thank the study participants and the Data Safety Monitoring Board members and the clinical research staff at participating sites. The authors have reported the following: I.O.R. reports personal fees from Genentech, Inc, Boehringer Ingelheim, and Three Lakes Partners. I.O.R. has a patent for a novel panel of biomarkers to predict and diagnose idiopathic pulmonary fibrosis with royalties paid to University of Pittsburgh. A.M.K.C. is a co-founder of Proterris, Inc and served on the TEVA Advisory Board meeting as a consultant on July 12–13, 2018.

Author Contributions

N.C., T.Z. and J.G.N.G. conceived of the study. T.Z. and N.C. collected and processed the microarray and nanostring data. T.Z. performed the statistical analysis. J.G.N.G. helped interpret the results. N.C., T.Z., M.L.G.-G., drafted the manuscript. I.O.R., S.W.R., A.M.K.C. and J.G.N.G. helped to critically revise the manuscript. I.O.R., N.C., H.J.G. S.E.-C., R.F.M., K.R.F., G.M.H., H.R.C., D.L.J., J.A.L., F.J.M., I.N., G.R. and A.M.K.C. contributed to study conception and implementation of the clinical study. All authors read and approved the final manuscript.

Additional Information

Competing Interests: The authors declare no competing interests.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019