

UC Merced

UC Merced Previously Published Works

Title

High-quality permanent draft genome sequence of the Bradyrhizobium elkanii type strain USDA 76T, isolated from Glycine max (L.) Merr

Permalink

<https://escholarship.org/uc/item/9jj7j9wp>

Journal

Environmental Microbiome, 12(1)

ISSN

1944-3277

Authors

Reeve, Wayne
van Berkum, Peter
Ardley, Julie
et al.

Publication Date

2017

DOI

10.1186/s40793-017-0238-2

Peer reviewed

EXTENDED GENOME REPORT

Open Access



High-quality permanent draft genome sequence of the *Bradyrhizobium elkanii* type strain USDA 76^T, isolated from *Glycine max* (L.) Merr

Wayne Reeve^{1*}, Peter van Berkum², Julie Ardley¹ , Rui Tian¹, Margaret Gollagher³, Dora Marinova³, Patrick Elia², T. B. K. Reddy⁴, Manoj Pillay⁵, Neha Varghese⁴, Rekha Seshadri⁴, Natalia Ivanova⁴, Tanja Woyke⁴, Mohamed N. Baeshen⁶, Nabih A. Baeshen⁷ and Nikos Kyrpides^{4,7}

Abstract

Bradyrhizobium elkanii USDA 76^T (INSCD = ARAG00000000), the type strain for *Bradyrhizobium elkanii*, is an aerobic, motile, Gram-negative, non-spore-forming rod that was isolated from an effective nitrogen-fixing root nodule of *Glycine max* (L. Merr) grown in the USA. Because of its significance as a microsymbiont of this economically important legume, *B. elkanii* USDA 76^T was selected as part of the DOE Joint Genome Institute 2010 *Genomic Encyclopedia for Bacteria and Archaea-Root Nodule Bacteria* sequencing project. Here the symbiotic abilities of *B. elkanii* USDA 76^T are described, together with its genome sequence information and annotation. The 9,484,767 bp high-quality draft genome is arranged in 2 scaffolds of 25 contigs, containing 9060 protein-coding genes and 91 RNA-only encoding genes. The *B. elkanii* USDA 76^T genome contains a low GC content region with symbiotic *nod* and *fix* genes, indicating the presence of a symbiotic island integration. A comparison of five *B. elkanii* genomes that formed a clique revealed that 356 of the 9060 protein coding genes of USDA 76^T were unique, including 22 genes of an intact resident prophage. A conserved set of 7556 genes were also identified for this species, including genes encoding a general secretion pathway as well as type II, III, IV and VI secretion system proteins. The type III secretion system has previously been characterized as a host determinant for *Rj* and/or *rj* soybean cultivars. Here we show that the USDA 76^T genome contains genes encoding all the type III secretion system components, including a translocon complex protein NopX required for the introduction of effector proteins into host cells. While many bradyrhizobial strains are unable to nodulate the soybean cultivar Clark (*rj1*), USDA 76^T was able to elicit nodules on Clark (*rj1*), although in reduced numbers, when plants were grown in Leonard jars containing sand or vermiculite. In these conditions, we postulate that the presence of NopX allows USDA 76^T to introduce various effector molecules into this host to enable nodulation.

Keywords: Root-nodule bacteria, GEBA-RNB, Nitrogen fixation, *Bradyrhizobium*, Soybean, Type III secretion system

Introduction

Soybean (*Glycine max*) (L.) Merr. is the dominant and the most important commercial legume crop species, yielding food oil and animal meal as well as nutritious vegetable protein [1–3]. The plant was first introduced into USA agriculture during the mid-18th century and was mainly used as a forage crop until the 1920s [4].

The development of new cultivars, along with technological advances in soybean processing and increased demand for soybean products, has led to major increases in production during the 20th century [4].

As with most papilionoid legumes, soybean engages in a symbiotic relationship with dinitrogen-fixing soil bacteria known as rhizobia and is able to obtain on average 50–60% of its required nitrogen through symbiotic nitrogen fixation [5]. A greater understanding of the symbiosis between soybean and its cognate rhizobia is of direct relevance for maintaining environmentally sustainable high

* Correspondence: W.Reeve@murdoch.edu.au

¹School of Veterinary and Life Sciences, Murdoch University, Murdoch, Australia

Full list of author information is available at the end of the article



crop yields, which significantly contributes to the Sustainable Development Goals adopted in September 2015 as part of the UN's development agenda 'Transforming our world: the 2030 Agenda for Sustainable Development' [6].

The soybean-nodulating bacteria, known as *Rhizobium japonicum* according to a 1929 classification scheme [7], were reclassified as *Bradyrhizobium japonicum* in 1982 because of several fundamental morphological and physiological differences with the genus *Rhizobium* [8]. The bacteria isolated from nodules of soybean had previously been shown to be phenotypically diverse, even though they were grouped together in the species *Bradyrhizobium japonicum*. One of the major methods that demonstrated this diversity was serology, which was used to classify individual isolates into 17 distinct serogroups [9]. This was accomplished by generating antisera to specific strains in the USDA collection in Beltsville and then using the sera to generate a serological scheme. One of the strains used to generate antisera was USDA 76^T and all isolates that cross-reacted with the antiserum generated with this serotype strain were combined together in the 76 serogroup. The strain USDA 76^T deposited in the Beltsville collection was a re-isolate from a greenhouse-grown plant inoculated with USDA 74 in Maryland. In turn, USDA 74 was a re-isolate of USDA 8 from a plant passage field test in California in 1956. The original parent culture of USDA 76^T is USDA 8, which was isolated from soybean grown at the Arlington Farm, Virginia in 1915.

Differences among the soybean root nodule bacteria classified as *B. japonicum* were also demonstrated using molecular methods. Hollis et al. [10] reported the presence of three DNA homology groupings by analysis of 28 strains within the soybean rhizobia. Using this approach, nine of the 17 serogroups were assigned to three DNA homology groupings: group I, the closely related group Ia and the more divergent group II. Supporting evidence for these three groupings was obtained by Kuykendall et al. [11]. By sequence analysis of the 16S rRNA genes, each of the 17 serotype strains representing

the serogroups were also placed into three closely related groups [12] that matched their separation by DNA homology. Since soybean strains could be distinguished phenotypically and by several approaches in molecular biology, Kuykendall et al. [13] proposed that DNA homology group II strains be separated from *B. japonicum* as the species *Bradyrhizobium elkanii*, with USDA 76^T as the type strain.

Because of these distinguishing characteristics and its significance as a microsymbiont of the economically important legume soybean, *B. elkanii* USDA 76^T was selected as part of the DOE Joint Genome Institute 2010 *Genomic Encyclopedia for Bacteria and Archaea-Root Nodule Bacteria* sequencing project [14, 15]. Here we present a summary classification and a set of general features for *B. elkanii* strain USDA 76^T, together with a description of its genome sequence and annotation.

Organism information

Classification and features

Bradyrhizobium elkanii USDA 76^T is a motile, non-sporulating, non-encapsulated, Gram-negative strain in the order *Rhizobiales* of the class *Alphaproteobacteria*. The rod shaped form has dimensions of approximately 0.5 μm in width and 1.0–2.0 μm in length (Fig. 1 *Left* and *Center*). It is relatively slow growing, forming colonies after 6–7 days when grown on $\frac{1}{2}$ Lupin Agar [16], Modified Arabinose Gluconate [17] and modified Yeast Mannitol Agar [18] at 28 °C. Colonies on $\frac{1}{2}$ LA are opaque, slightly domed and moderately mucoid with smooth margins (Fig. 1 *Right*).

Sequence divergence among the 16S rRNA genes of the 33 type strains within the genus *Bradyrhizobium* was limited and ranged from no differences in many cases to a similarity of 98% between *B. elkanii* USDA 76^T and *B. neotropicale* (Fig. 2) after accounting for 40 bp in gaps along the alignment length. Such high similarity values would question the reliability of defining species limits within the genus based on divergence of the 16S rRNA genes [19]. Bootstrap values for each of the nodes of the





Fig. 1 Images of *Bradyrhizobium elkanii* USDA 76^T using scanning (*Left*) and transmission (*Center*) electron microscopy as well as light microscopy to visualize colony morphology on solid media (*Right*)



branches were low and none of the confidence values reached or exceeded 95%. Therefore, the placement of each of the taxa relative to the others in the tree is inconclusive.

Genetic recombination resulting in a reticulate evolutionary history of the 16S rRNA gene is perhaps a likely explanation for the low bootstrap values. Therefore, an analysis for recombination was done with the aligned 33 *Bradyrhizobium* 16S rRNA genes using the pairwise homoplasy index test [20]. By using this test, statistically significant evidence for recombination among the 33 16S rRNA genes was detected ($P = 0.003$). The detection

of genetic recombination within the *rrn* loci of rhizobia is not unprecedented since reticulate evolutionary histories of the 16S rRNA genes and the Internally Transcribed Spacer between the 16S and 23S rRNA genes has been described before [21, 22]. The 16S rRNA sequence of *B. pachyrhizi* was identical with those of the *B. elkanii* serogroup strains USDA 31, USDA 94 and USDA 130, which differed from *B. elkanii* USDA 76^T by one bp (99.999% similar). The most divergent 16S rRNA gene within *B. elkanii* was that of the serogroup strain USDA 46 (99.996% similar), while the most divergence among the soybean serogroup strains was that between

USDA 46 and USDA 110, which were 98.4% similar. Since the divergence of the 16S rRNA genes of the genus *Bradyrhizobium* is narrow, with evidence for the presence of a history of genetic recombination, it may be necessary to more precisely establish their phylogeny by comparing their entire genomes rather than individual genes. Such an approach may provide more fundamental insight into the evolutionary history of this class of symbiotic bacteria as well as impacting potential changes in their current proposed taxonomy. Minimum Information about the Genome Sequence of USDA 76^T is provided in Table 1 and Additional file 1: Table S1.

Symbiotaxonomy

An investigation of the symbiotic properties of soybean began with the work of Brooks [23] in the late 19th century, when he observed that soybean grown in the fields of his experiment station in Massachusetts only nodulated when supplied with dust he had brought with him from Japan. This led to the theory that soybean-nodulating bacteria in the soils of the USA were imported from the Far East. Cotrell et al. [24] and Hopkins [25] reported the supporting evidence that soybean in Kansas nodulated with soil taken from the Massachusetts Experiment station, or in Illinois from soil collected from fields with a history of soybean cultivation. However, several decades later it became evident that rhizobia that nodulated native American legumes within the genera *Apios*, *Amphicarpa*, *Crotalaria*, *Desmodium*, *Lespedeza*, *Baptisia*, *Cassia*, *Genista* and *Wisteria* also nodulated soybean [26–28]. With the exception of USDA 6 and USDA 38, which are from Japan, all the remaining soybean serotype strains were recovered from nodules of soybeans grown in the USA, including USDA 8 (the original parent of USDA 76^T). Consequently, it is unclear whether these rhizobia obtained from nodules of USA-grown soybean originate from the Far East or are in fact native to the soils of America. Therefore, the possibility exists that USDA 76^T may be able to nodulate and form a symbiosis with a wide variety of legumes, but this has not been thoroughly investigated. Unfortunately, the communication that included the proposal of USDA 76^T as the type strain for *B. elkanii* did not include results of plant tests to describe its symbiotic range, but instead relied on distinction by phenotype and genotype [11]. An indication of the possible American origin of USDA 76^T is its reported effectiveness in symbiosis with the native *Apios americana* Medik. and use as an inoculum for this potential leguminous crop [29]. Further evidence for this theory is the ability of USDA 76^T to nodulate and fix nitrogen with the native American *Amphicarpaea bracteata* (L.) Fernald [30]. USDA 76^T effectively nodulates the promiscuous *Vigna unguiculata* (L.) Walp.

Table 1 Classification and general features of *Bradyrhizobium elkanii* USDA 76^T in accordance with the MIGS recommendations [71] published by the Genome Standards Consortium [72]

MIGS ID	Property	Term	Evidence code
	Classification	Domain Bacteria	TAS [73]
		Phylum <i>Proteobacteria</i>	TAS [74, 75]
		Class <i>Alphaproteobacteria</i>	TAS [74, 76]
		Order <i>Rhizobiales</i>	TAS [77]
		Family <i>Bradyrhizobiaceae</i>	TAS [78]
		Genus <i>Bradyrhizobium</i>	TAS [8, 78]
		Species <i>elkanii</i>	IDA
	Gram stain	Negative	IDA
	Cell shape	Rod	IDA
	Motility	Motile	IDA
	Sporulation	Non-sporulating	NAS
	Temperature range	Mesophile	NAS
	Optimum temperature	28°C	NAS
	pH range; Optimum	Unknown	NAS
	Carbon source	Arabinose, gluconate	TAS [17]
MIGS-6	Habitat	Soil, root nodule of <i>Glycine max</i> (L. Merr)	NAS
MIGS-6.3	Salinity	0 to <2% (w/v) NaCl	TAS [78]
MIGS-22	Oxygen requirement	Aerobic	NAS
MIGS-15	Biotic relationship	Free living, symbiotic	TAS
MIGS-14	Pathogenicity	Non-pathogenic	TAS [79]
MIGS-4	Geographic location	Alexandria, Virginia, USA	NAS
MIGS-5	Sample collection date	1915	NAS
MIGS-4.1	Latitude	38.8047	NAS
MIGS-4.2	Longitude	-77.0472	NAS
MIGS-4.3	Depth	5 cm	NAS
MIGS-4.4	Altitude	13 m	NAS

Evidence codes—*IDA* Inferred from Direct Assay, *TAS* Traceable Author Statement (i.e., a direct report exists in the literature), *NAS* Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). Evidence codes are from the Gene Ontology project [80, 81]

The original isolation location and date indicated is that of the parent culture USDA 8

(cowpea), but is unable to nodulate the tropical American legume *Phaseolus lunatus* L. (Lima bean), which forms nodules with various other strains of bradyrhizobia [31]. To our knowledge, the only other reported information is that USDA 74 (parent of USDA 76^T) forms an effective symbiosis with *Macroptilium atropurpureum* (DC.) Urb. (Siratiro) and *Vigna unguiculata* (L.) Walp [32].

In soybean, the *Rj*(s) or *rj*(s) genetic loci have been identified as controlling the ability of compatible rhizobia to nodulate with a particular cultivar (reviewed by Hayashi et al. [33]). USDA 76^T is reported to form nodules (albeit in reduced numbers) on the cultivar Clark (*rj1*) and to nodulate and fix N₂ with the isogenic lines BARC-2 and BARC-3, harboring the *Rj4* and *rj4* alleles, respectively, when tested in Leonard jars with sterile vermiculite or sand [30]. The symbiotic characteristics of *B. elkanii* USDA 76^T on a range of selected hosts are summarized in Additional file 2: Table S2.

Genome sequencing information

Genome project history

This organism was selected for sequencing at the U.S. Department of Energy funded Joint Genome Institute as part of the *Genomic Encyclopedia of Bacteria and Archaea-Root Nodule Bacteria* project project [14, 15]. The root nodule bacteria in this project were selected on the basis of environmental and agricultural relevance to issues in global carbon cycling, alternative energy production, and biogeochemical importance. In particular, strain USDA 76^T was chosen since it is a microsymbiont of the economically important legume soybean, but can also form symbioses with several legumes native to the USA. The USDA 76^T genome project is deposited in the Genomes Online Database [34] and a high-quality permanent draft genome sequence is deposited in IMG [35]. Sequencing, finishing and annotation were performed by the JGI [36] and a summary of the project information is shown in Table 2.

Growth conditions and genomic DNA preparation

After recovery from permanent storage, the *B. elkanii* USDA 76^T was streaked onto MAG solid medium and grown at 28 °C for 6 days to obtain well grown, well separated colonies, then a single colony was selected and used to inoculate 5 ml MAG broth. The culture was grown on a gyratory shaker (200 rpm) at 28 °C for 6 days. Subsequently 1 ml was used to inoculate 50 ml MAG broth and grown on a gyratory shaker (200 rpm) at 28 °C until an OD_{600nm} of 0.6 was reached. DNA was isolated from the cells according to van Berkum [17]. Final concentration of the DNA was set to 0.5 mg ml⁻¹. Culture identity was confirmed by partial sequence analysis of several housekeeping genes and the 16S rRNA gene using the prepared DNA as template for PCR.

Genome sequencing and assembly

The draft genome of *B. elkanii* USDA 76^T was generated at the DOE Joint genome Institute (JGI) using the Illumina technology [37]. An Illumina short-insert paired-end library was constructed with an average insert size of 200 bp that when sequenced generated

Table 2 Genome sequencing project information of *Bradyrhizobium elkanii* strain USDA 76^T

MIGS ID	Property	Term
MIGS-31	Finishing quality	High-quality permanent draft
MIGS-28	Libraries used	2× Illumina libraries; Std short PE & CLIP long PE
MIGS-29	Sequencing platforms	Illumina HiSeq2000, PacBio
MIGS-31.2	Fold coverage	3,560x
MIGS-30	Assemblers	Velvet version 1.1.05; Allpaths-LG version r38445; phrap, version 4.24
MIGS-32	Gene calling methods	Prodigal 1.4; GenePRIMP
	Locus Tag	Brael [82]
	GenBank ID	ARAG000000000
	GenBank Date of Release	Apr 22, 2013
	GOLD ID	Gp0009610
	NCBI BIOPROJECT	162247
MIGS-13	Source Material Identifier	USDA 76, USDA 8, USDA 74, ATCC 49852, DSM 11554, IFO (now NBRC) 14791, LMG 6134
	Project relevance	Symbiotic N ₂ fixation, agriculture

312,796,730 reads. An Illumina long-insert paired-end library with an average insert size of 6505.78 +/- 3679.88 bp also was constructed that when sequenced generated 19,315,434 reads. The total amount of sequence data obtained with the Illumina was 34,177 Mbp. Library construction and sequence analysis were done at the JGI according to the protocols outlined on their website [38]. The first of two initial drafts, assembled with Allpaths version r38445 [39], contained 81 contigs in 17 scaffolds and subsequently a consensus was computationally shredded into 10 Kbp overlapping fake reads (shreds). The second draft assembled with Velvet, version 1.1.05 [40], resulted in consensus sequences that were computationally shredded into 1.5 Kbp overlapping fake reads (shreds). The data were assembled again with Velvet using the shreds from the first Velvet assembly to guide the next assembly. The consensus from this second Velvet assembly was shredded into 1.5 Kbp overlapping fake reads. The fake reads from the Allpaths and both Velvet assemblies together with a subset of the Illumina CLIP paired-end reads were assembled using parallel Phrap, version 4.24 (High Performance Software, LLC). Potential errors in the assemblies were corrected by manual editing with Consed [41–43]. Gap closure was accomplished using repeat resolution software (Wei Gu, unpublished) and sequence analysis of bridging PCR fragments with PacBio technology (Cliff Han, unpublished). Gaps were closed and the quality of the final sequence was

improved with 35 PCR PacBio consensus sequences. The total size of the genome is 9.5 Mbp and the final assembly is based on 34,177 Mbp of Illumina draft data, which provides an average 3560x coverage of the genome.

Genome annotation

Genes were identified using Prodigal [44] that was followed by a round of manual curation using GenePRIMP [45] as part of the DOE-JGI genome annotation pipeline [46, 47]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) non-redundant, UniProt, TIGRFam, Pfam, KEGG, COG, and InterPro databases. The tRNAScanSE tool [48] was used to find tRNA genes. Ribosomal RNA genes were found by searches against models of the ribosomal RNA genes built from SILVA [49]. Other non-coding RNAs such as the RNA components of the protein secretion complex and the RNase P were identified by searching the genome for the corresponding Rfam profiles using INFERNAL [50]. Additional gene prediction analysis and manual functional annotation were done within the Integrated Microbial Genomes-Expert Review system [51] developed by the Joint Genome Institute, Walnut Creek, CA, USA.

Genome properties

The genome of *B. elkanii* USDA 76^T is 9,484,767 nucleotides long with a GC content of 63.70% (Table 3) and has been assembled into two scaffolds. Of the 9151 genes identified, 9060 are protein encoding and 91 are RNA only encoding genes. Of the 9151 total genes identified in USDA 76^T, the majority (73.28%) were assigned

Table 3 Genome statistics for *Bradyrhizobium elkanii* USDA 76^T

Attribute	Value	% of Total
Genome size (bp)	9,484,767	100.00
DNA coding (bp)	8,070,200	85.09
DNA G + C (bp)	6,041,732	63.70
DNA scaffolds	2	100.00
Total genes	9151	100.00
Protein coding genes	9060	99.01
RNA genes	91	0.99
Pseudo genes	408	4.46
Genes in internal clusters	789	8.62
Genes with function prediction	6706	73.28
Genes assigned to COGs	5665	61.91
Genes with Pfam domains	7004	76.54
Genes with signal peptides	864	9.44
Genes with transmembrane helices	2055	22.46
CRISPR repeats	2	


a putative function and the remaining genes were annotated as hypothetical. The distribution of genes into COGs functional categories is presented in Table 4.

Insights from the genome sequence

Scaffold 1.1 of *B. elkanii* USDA 76^T contains a low GC content for the region ~3,000,000–3,800,000 and the presence of symbiotic *nod*, *nif* and *fix* genes in this region indicates a symbiotic island integration (Fig. 3). Using the Phylogenetic Profiler tool within IMG, 356 genes were found to be unique to USDA 76^T in a

Table 4 Number of protein coding genes of *Bradyrhizobium elkanii* USDA 76^T associated with the general COG functional categories

Code	Value	Percent	COG Category
J	235	3.63	Translation, ribosomal structure and biogenesis
A	0	0.00	RNA processing and modification
K	514	7.93	Transcription
L	175	2.70	Replication, recombination and repair
B	2	0.03	Chromatin structure and dynamics
D	40	0.62	Cell cycle control, cell division, chromosome partitioning
V	165	2.55	Defense mechanisms
T	253	3.90	Signal transduction mechanisms
M	313	4.83	Cell wall/membrane/envelope biogenesis
N	80	1.23	Cell motility
U	135	2.08	Intracellular trafficking, secretion, and vesicular transport
O	267	4.12	Posttranslational modification, protein turnover, chaperones
C	439	6.77	Energy production and conversion
G	392	6.05	Carbohydrate transport and metabolism
E	685	10.57	Amino acid transport and metabolism
F	94	1.45	Nucleotide transport and metabolism
H	317	4.89	Coenzyme transport and metabolism
I	423	6.53	Lipid transport and metabolism
P	381	5.88	Inorganic ion transport and metabolism
Q	295	4.55	Secondary metabolite biosynthesis, transport and catabolism
R	663	10.23	General function prediction only
S	399	6.16	Function unknown
-	3486	38.09	Not in COGS



comparison with four other strains (587 [52], CCBAU43297, CCBAU05737 [53] and USDA 94) ascribed to the *B. elkanii* IMG clique. Of those that were unique, the majority (223 genes, representing 62.6%) were annotated as encoding hypothetical proteins. Out of the remainder, a significant number were phage related. Using the PHASTER algorithm [54], 22 of these genes were found to be co-located genes of an intact resident prophage (Fig. 4). Using this algorithm another incomplete phage gene set on the same scaffold was also identified.

Extended insights

Using the Phylogenetic Profiler tool, 7556 genes were found to be conserved in five *B. elkanii* strains (587, CCBAU43297, CCBAU05737, USDA 76^T, USDA 94), including genes encoding a general secretion pathway and type II, III, IV and VI secretion system proteins. The Type III secretion system (T3SS) [55] can either promote or impair the establishment of symbiosis, depending on the legume host [56], and has been characterized as a host determinant for *rj1*, *Rfg1*, *Rj2* and *Rj4* soybean cultivars [33, 57, 58]. The dominant soybean genes *Rj2* and *Rj4* restrict nodulation with specific strains of *Bradyrhizobium* [33]. Most investigations of soybean host genes controlling the symbiosis have focused on the *Rj4* soybean line that was originally identified by its inability to nodulate with USDA 61 (*B. elkanii*, serogroup 31) [59]. The predicted *Rj4* thaumatin-like protein is thought to be involved in



conferring resistance to *Bradyrhizobium* strains producing specific T3SS effector proteins [60]. However, USDA 76^T was reported to nodulate and form an effective nitrogen-fixing symbiosis with the isogenic lines BARC-2 (*Rj4*) and BARC-3 (*rj4*) [30, 61], suggesting that this strain does not produce the interacting T3SS effector protein(s). Conversely, the recessive soybean gene *rj1rj1* [62], encoding a putative truncated Nod factor receptor protein [63], restricts nodulation by many *Bradyrhizobium* and *Ensifer* strains, although specific strains of *B. elkanii*, including USDA 76^T, can form a limited number of nodules when tested with plants in Leonard jars containing sterilized vermiculite or sand [30, 59, 61].

USDA 76^T genes encoding components required for a functional T3SS were identified within the integrated symbiotic island (Figs. 5 and 6). Although the *nopA* and *nopC* genes were not annotated in the USDA 76^T genome, by using TBLASTN these genes were identified in the intergenic region between BraelDRAFT_3047 (*sctD*) and BraelDRAFT_3048 (hypothetical) that share 100% sequence similarity with *nopA* and *nopC* of the characterized *Bradyrhizobium elkanii* strain USDA 61 [57]. Although T3SS components can also be found in *Bradyrhizobium* strain USDA 110, this strain lacks the *nopX* gene encoding the translocon required to introduce effector molecules into host cells [56, 64]. This is in contrast to the presence of *nopX* in USDA 76^T, which could extend its host range to otherwise incompatible hosts.

Belk USDA 76^T: BraelDRAFT_scaffold1.1

Prophage Region 1

Start: 5947991
 End: 5968342
 # CDS: 22
 Predicted Type: intact
 GC%: 63.53



Prophage Region 2

Start: 7061278
 End: 7071123
 # CDS: 16
 Predicted Type: incomplete
 GC%: 58.57




Fig. 4 Resident prophage present in *Bradyrhizobium elkanii* USDA 76^T imaged using PHASTER [54]. Prophage maps are not drawn to scale. Reference locus tag for Prophage Region 1 is BraelDRAFT_5594 terminase; ter; reference locus tag for Prophage Region 2 is BraelDRAFT_6751 (terminase; ter). Coat protein (coa), fiber protein (fib), phage-like protein (plp), portal protein (por), tail shaft protein (sha), and terminase (ter). All other genes encode hypothetical proteins





Fig. 5 Comparison of the gene neighbourhood regions containing loci that encode type III secretion system components in the genomes of *Ensifer fredii* NGR234 and the *Bradyrhizobium* strains USDA 76^T and USDA 110. The colour scheme is as follows: green, structural component; orange, pilus component; purple, regulatory component; red, translocon component; uncoloured, other genes; and yellow, effector component



Conclusions

B. elkanii USDA 76^T originated from strain USDA 8, which was obtained in 1915 from an effective nodule of soybean grown on the USDA Arlington farm in Virginia. Its ability to nodulate the native North American legumes *Apios americana* Medik. and *Amphicarpaea bracteata* (L.) Fernald indicates a possible North American origin for this strain. USDA 76^T was selected for genome sequencing [14] because of its significance as a microsymbiont of soybean. The genome size of USDA 76^T was established as 9.5 Mbp, which falls within the range of 7.7 to 10.5 Mbp observed for other bradyrhizobial genomes. The genome of this N₂-fixing microsymbiont contains *nod*, *nif* and *fix* genes located on an integrated symbiotic island, and genes encoding both an intact and an incomplete phage. According to ANI values, strain USDA 76^T formed an ANI clique with four other *B. elkanii* soybean strains: USDA 94, 587, CCBAU 43297 and CCBAU 05737. Of particular interest was the discovery that these strains contain a T3SS that contains the NopCA pilus genes and the NopX translocon protein, which are essential for introducing effector molecules into host cells [55]. The T3SS has been shown to be an important host range determinant that enables the nodulation of some soybean cultivars and is detrimental to symbiosis with other cultivars [56]. Here we postulate that the presence of a functional T3SS is important in

determining the host range of USDA 76^T and enables it to form some nodules on the soybean cultivar Clark (*rj1*) when grown in Leonard jars with sterilized vermiculite or sand [65, 66]. Further analyses of *Bradyrhizobium* genomes, including that of USDA 76^T, will increase our understanding of determinants that lead to the establishment and functioning of different *Bradyrhizobium* symbioses.

Additional files

Additional file 1: Associated MIGS record. Table S1. Associated MIGS record for *Bradyrhizobium elkanii* USDA 76^T. (DOCX 19 kb)

Additional file 2: Symbiotic properties of USDA 76^T. Table S2. Nodulation and N₂-fixation properties of *Bradyrhizobium elkanii* USDA 76^T on selected legume hosts. (DOCX 16 kb)

Abbreviations

½ LA: ½ Lupin Agar; ANI: Average Nucleotide Identity; GEBA-RNB: Genomic Encyclopedia for Bacteria and Archaea-Root Nodule Bacteria; IMG: Integrated Microbial Genomes; MAG: Modified Arabinose Gluconate; T3SS: Type 3 Secretion System

Acknowledgements

We thank Gordon Thompson (Murdoch University) for the preparation of SEM and TEM photos.

Funding

This work was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory

under contract No. DE-AC02-05CH11231. We gratefully acknowledge the funding received from the Curtin University Sustainability Policy Institute, and the funding received from Murdoch University Small Research Grants Scheme in 2016.

Authors' contributions

PVB supplied the strain, background information for this project and the DNA to the JGI; TR performed all imaging; PVB, TR, JA and WR drafted the paper; MNB and NAB provided financial support and MG, DM, PE, TBKR, VM, NI, TW, RS and NK were involved in sequencing the genome and/or editing the final paper. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Author details

¹School of Veterinary and Life Sciences, Murdoch University, Murdoch, Australia. ²U.S. Department of Agriculture, Soybean Genomics and Improvement Laboratory, Beltsville Agricultural Research Center, 10300 Baltimore Avenue, Bldg. 006, Beltsville, MD 20705, USA. ³Curtin University Sustainability Policy Institute, Curtin University, Bentley, WA, Australia. ⁴DOE Joint Genome Institute, Walnut Creek, CA, USA. ⁵Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. ⁶Department of Biology, Faculty of Science, University of Jeddah, Jeddah, Saudi Arabia. ⁷Department of Biological Sciences, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia.

Received: 14 October 2016 Accepted: 21 February 2017

Published online: 04 March 2017

References

- Graham PH, Vance CP. Legumes: Importance and constraints to greater use. *Plant Physiol.* 2003;131:872–7.
- Herridge DF, Peoples MB, Boddey RM. Global inputs of biological nitrogen fixation in agricultural systems. *Plant Soil.* 2008;311:1–18.
- Soybean International Commodity Profile. http://siteresources.worldbank.org/INTAFRICA/Resources/257994-1215457178567/Soybean_Profile.pdf. Accessed 24 Feb 2017.
- Hymowitz T. Speciation and cytogenetics. In: Boerma HR, Specht JE, editors. *Soybeans: Improvement, Production, and Uses*. Madison: American Society of Agronomy, Crop Science Society of America, Soil Science Society of America; 2005. p. 110–1.
- Salvagioti F, Cassman KG, Specht JE, Walters DT, Weiss A, Dobermann A. Nitrogen uptake, fixation and response to fertilizer N in soybeans: A review. *Field Crops Res.* 2008;108:1–13.
- UNDP 2015. <http://www.undp.org/content/undp/en/home/sdgoverview/post-2015-development-agenda.html>. Accessed 24 Feb 2017.
- Fred EB, Baldwin IL, McCoy E. Root nodule bacteria and leguminous plants. Madison: University of Wisconsin-Madison Libraries Parallel Press; 1932.
- Jordan DC. Transfer of *Rhizobium japonicum* Buchanan 1980 to *Bradyrhizobium* gen. nov., a genus of slow-growing, root nodule bacteria from leguminous plants. *Int J Syst Bacteriol.* 1982;32:136–9.
- Date RA, Decker AM. Minimal antigenic constitution of 28 strains of *Rhizobium japonicum*. *Can J Microbiol.* 1965;11:1–8.
- Hollis AB, Kloos WE, Elkan GH. DNA:DNA hybridization studies of *Rhizobium japonicum* and related *Rhizobiaceae*. *J Gen Microbiol.* 1981;123:215–22.
- Kuykendall LD, Saxena B, Devine TE, Udell SE. Genetic diversity in *Bradyrhizobium-japonicum* Jordan 1982 and a proposal for *Bradyrhizobium elkanii* sp. nov. *Can J Microbiol.* 1992;38:501–5.
- van Berkum P, Fuhrmann JJ. Evolutionary relationships among the soybean bradyrhizobia reconstructed from 16S rRNA gene and internally transcribed spacer region sequence divergence. *Int J Syst Evol Microbiol.* 2000;50(Pt 6):2165–72.
- Kuykendall LD, Roy MA, O'Neill JJ, Devine TE. Fatty acids, antibiotic resistance, and deoxyribonucleic acid homology groups of *Bradyrhizobium japonicum*. *Int J Syst Evol Microbiol.* 1988;38:358–61.
- Reeve WG, Ardley J, Tian R, Eshragi L, Yoon JW, Ngamwisetkun P, Seshadri R, Ivanova NN, Kyrpidis NC. A genomic encyclopedia of the root nodule bacteria: Assessing genetic diversity through a systematic biogeographic survey. *Stand Genomic Sci.* 2015;10:14.
- Seshadri R, Reeve WG, Ardley JK, Tennesen K, Woyke T, Kyrpidis NC, Ivanova NN. Discovery of novel plant interaction determinants from the genomes of 163 Root Nodule Bacteria. *Sci Rep.* 2015;5:16825.
- Howieson JG, Ewing MA, D'Antuono MF. Selection for acid tolerance in *Rhizobium meliloti*. *Plant Soil.* 1988;105:179–88.
- van Berkum P. Evidence for a third uptake hydrogenase phenotype among the soybean bradyrhizobia. *Appl Environ Microbiol.* 1990;56:3835–41.
- Vincent JM. A manual for the practical study of the root-nodule bacteria. International Biological Programme. UK: Blackwell Scientific Publications, Oxford; 1970.
- Stackebrandt E, Goebel BM. Taxonomic note: A place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol.* 1994;44:846–9.
- Bruen TC, Philippe H, Bryant D. A simple and robust statistical test for detecting the presence of recombination. *Genetics.* 2006;172:2665–81.
- van Berkum P, Fuhrmann J. Evidence from internally transcribed spacer sequence analysis of soybean strains that extant *Bradyrhizobium* spp. are likely the products of reticulate evolutionary events. *Appl Environ Microbiol.* 2009;75:78–82.
- van Berkum P, Terefework Z, Paulin L, Suomalainen S, Lindstrom K, Eardly BD. Discordant phylogenies with the *rrn* loci of rhizobia. *J Bacteriol.* 2003;185:2988–98.
- Brooks WP. *Agriculture, vol II. Manures, Fertilizers and Farm Crops, including green manuring and crop rotation.* MA, USA: The Home Correspondence School. Springfield; 1901.
- Cottrell HM, Otis DH, Haney JG. Soil inoculation for soybeans. *Kans Agric Exp Stn Bull.* 1900;97:116.
- Hopkins CG. Nitrogen bacteria and legumes. *Ill Agric Exp Station Bull.* 1904; 306:28.
- Bushnell OA, Sarles WB. Studies on the root-nodule bacteria of wild leguminous plants in Wisconsin. *Soil Sci.* 1937;44:409–23.
- Conklin ME. Studies of the root nodule organisms of certain wild legumes. *Soil Sci.* 1936;41:167–85.
- Wilson JK. Over five hundred reasons for abandoning the cross-inoculation groups of the legumes. *Soil Sci.* 1944;58:61–9.
- Putnam DH, Heichel GH, Field LA. Response of *Apios americana* to nitrogen and inoculation. *Hort Sci.* 1991;26:853–5.
- Marr DL, Devine TE, Parker MA. Nodulation restrictive genotypes of *Glycine* and *Amphicarpaea*: A comparative analysis. *Plant Soil.* 1997;189:181–8.
- López-López A, Negrete-Yankelevich S, Rogel MA, Ormeño-Orrillo E, Martínez J, Martínez-Romero E. Native bradyrhizobia from Los Tuxtlas in Mexico are symbionts of *Phaseolus lunatus* (Lima bean). *Syst Appl Microbiol.* 2013;36:33–8.
- Keyser HH, van Berkum P, Weber DF. A comparative study of the physiology of symbioses formed by *Rhizobium japonicum* with *Glycine max*, *Vigna unguiculata*, and *Macroptilium atropurpureum*. *Plant Physiol.* 1982;70:1626–30.
- Hayashi M, Saeki Y, Haga M, Harada K, Kouchi H, Umehara Y. *Rj* (*rj*) genes involved in nitrogen-fixing root nodule formation in soybean. *Breed Sci.* 2012;61:544–53.
- Pagani I, Liolios K, Jansson J, Chen IM, Smirnova T, Nosrat B, Markowitz VM, Kyrpidis NC. The Genomes OnLine Database (GOLD) v.4: Status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res.* 2012;40:D571–9.
- Markowitz VM, Chen IA, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A, Huang J, Woyke T, Huntemann M, et al. IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res.* 2014;42:D560–7.
- Mavromatis K, Land ML, Brettin TS, Quest DJ, Copeland A, Clum A, Goodwin L, Woyke T, Lapidus A, Klenk HP, et al. The fast changing landscape of sequencing technologies and their impact on microbial genome assemblies and annotation. *PLoS One.* 2012;7:e48837.
- Bennett S. Solexa Ltd. *Pharmacogenomics.* 2004;5:433–8.
- Joint Genome Institute. <http://jgi.doe.gov/>. Accessed 24 Feb 2017.
- Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, et al. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A.* 2011;108:1513–8.
- Zerbino DR. Using the Velvet *de novo* assembler for short-read sequencing technologies. *Curr Protoc Bioinformatics.* 2010;Chapter 11:Unit 11–15.
- Ewing B, Green P. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res.* 1998;8:186–94.

42. Ewing B, Hillier L, Wendl MC, Green P. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res.* 1998;8:175–85.
43. Gordon D, Abajian C, Green P. Consed: A graphical tool for sequence finishing. *Genome Res.* 1998;8:195–202.
44. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: Prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics.* 2010;11:119.
45. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. GenePRIMP: A gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods.* 2010;7:455–7.
46. Chen IMA, Markowitz VM, Chu K, Anderson I, Mavromatis K, Kyrpides NC, Ivanova NN. Improving microbial genome annotations in an integrated database context. *PLoS One.* 2013;8:e54859.
47. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Palaniappan K, Szeto E, Pillay M, Chen IM-A, Pati A, et al. The standard operating procedure of the DOE-JGI microbial genome annotation pipeline (MGAP v.4). *Stand Genomic Sci.* 2015;10:86.
48. Lowe TM, Eddy SR. tRNAscan-SE: A program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 1997;25:955–64.
49. Pruesse E, Quast C, Knittell K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO. SILVA: A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* 2007;35:7188–96.
50. Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics.* 2013;29:2933–5.
51. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. IMG ER: A system for microbial genome annotation expert review and curation. *Bioinformatics.* 2009;25:2271–8.
52. de Souza JAM, Tieppo E, Magnani Gds, Alves LM, Cardoso RL, Cruz LM, de Oliveira LF, Raittz RT, de Souza EM, Pedrosa FdO, Lemos EGdM. Draft genome sequence of the nitrogen-fixing symbiotic bacterium *Bradyrhizobium elkanii*. *J Bacteriol.* 2012;194:3547–8.
53. Tian CF, Zhou YJ, Zhang YM, Li QQ, Zhang YZ, Li DF, Wang S, Wang J, Gilbert LB, Li YR, Chen WX. Comparative genomics of rhizobia nodulating soybean suggests extensive recruitment of lineage-specific genes in adaptations. *Proc Natl Acad Sci U S A.* 2012;109:8629–34.
54. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res.* 2016;44:W16–21.
55. Galán JE, Lara-Tejero M, Marlovits TC, Wagner S. Bacterial type III secretion systems: specialized nanomachines for protein delivery into target cells. *Annu Rev Microbiol.* 2014;68:415–38.
56. Staehelin C, Krishnan HB. Nodulation outer proteins: double-edged swords of symbiotic rhizobia. *Biochem J.* 2015;470:263–74.
57. Okazaki S, Zehner S, Hempel J, Lang K, Gottfert M. Genetic organization and functional analysis of the type III secretion system of *Bradyrhizobium elkanii*. *FEMS Microbiol Lett.* 2009;295:88–95.
58. Tsurumaru H, Hashimoto S, Okizaki K, Kanesaki Y, Yoshikawa H, Yamakawa T. A putative type III secretion system effector encoded by the MA20_12780 gene in *Bradyrhizobium japonicum* Is-34 causes incompatibility with Rj4 genotype soybeans. *Appl Environ Microbiol.* 2015;81:5812–9.
59. Williams LF, Lynch DL. Inheritance of the non-nodulating character in the soybean. *Agron J.* 1954;46:28–9.
60. Indrasumunar A, Searle I, Lin MH, Kereszt A, Men A, Carroll BJ, Gresshoff PM. Nodulation factor receptor kinase 1alpha controls nodule organ number in soybean (*Glycine max* L. Merr). *Plant J.* 2011;65:39–50.
61. Devine TE, Kuykendall LD. Host genetic control of symbiosis in soybean (*Glycine max* L.). *Plant Soil.* 1996;186:173–87.
62. Devine TE, O'Neill JJ, Kuykendall LD. Near isogenic lines of soybeans as tools to identify nodulation specific mutants of *Bradyrhizobium elkanii*. *Plant Soil.* 1993;149:205–9.
63. Hayashi M, Shiro S, Kanamori H, Mori-Hosokawa S, Sasaki-Yamagata H, Sayama T, Nishioka M, Takahashi M, Ishimoto M, Katayose Y, et al. A thaumatin-like protein, Rj4, controls nodule symbiotic specificity in soybean. *Plant Cell Physiol.* 2014;55:1679–89.
64. Krause A, Doerfel A, Göttfert M. Mutational and transcriptional analysis of the type III secretion system of *Bradyrhizobium japonicum*. *Mol Plant Microbe Interact.* 2002;15:1228–35.
65. Devine TE, Breithaupt BH. Phenotypic thermal-stability of rhizobitoxine-induced chlorosis and the nodulation controlling gene, rj1. *Crop Sci.* 1980;20:394–6.
66. Devine TE, Kuykendall LD, Breithaupt BH. Nodulation of soybeans carrying the nodulation-restrictive gene, rj1, by an incompatible *Rhizobium japonicum* strain upon mixed inoculation with a compatible strain. *Can J Microbiol.* 1980;26:179–82.
67. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 2011;28:2731–9.
68. Nicholas KB, Nicholas HMB, Deerfield DW. GeneDoc: analysis and visualization of genetic variation. *EMBnet News.* 1997;4:14.
69. Felsenstein J. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution.* 1985;39:783–91.
70. Reddy TB, Thomas AD, Stamatis D, Bertsch J, Isbandi M, Jansson J, Mallajosyula J, Pagani I, Lobos EA, Kyrpides NC. The Genomes OnLine Database (GOLD) v.5: A metadata management system based on a four level (meta)genome project classification. *Nucleic Acids Res.* 2015;43:D1099–106.
71. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen M, Angiuoli SV, et al. Towards a richer description of our complete collection of genomes and metagenomes “Minimum Information about a Genome Sequence” (MIGS) specification. *Nat Biotechnol.* 2008;26:541–7.
72. Field D, Amaral-Zettler L, Cochrane G, Cole JR, Dawyndt P, Garrity GM, Gilbert J, Glöckner FO, Hirschman L, Karsch-Mizrachi I, et al. The Genomic Standards Consortium. *PLoS Biol.* 2011;9:e1001088.
73. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A.* 1990;87:4576–9.
74. Euzéby J. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Evol Microbiol.* 2005;55:2235–8.
75. Garrity GM, Bell JA, Lilburn T. Phylum XIV. *Proteobacteria* phyl. nov. In: Garrity GM, Brenner DJ, Kreig NR, Staley JT, editors. *Bergey's Manual of Systematic Bacteriology*, vol. 2. Second ed. New York: Springer-Verlag; 2005. p. 1.
76. Garrity GM, Bell JA, Lilburn T. Class I. *Alphaproteobacteria* class. In: Garrity GM, Brenner DJ, Kreig NR, Staley JT, editors. *Bergey's Manual of Systematic Bacteriology*. Second ed. New York: Springer-Verlag; 2005.
77. Kuykendall LD. Order VI. *Rhizobiales* ord. nov. In: Garrity GM, Brenner DJ, Kreig NR, Staley JT, editors. *Bergey's Manual of Systematic Bacteriology*. Second ed. New York: Springer-Verlag; 2005. p. 324.
78. Garrity GM, Bell JA, Lilburn T. Family VII. *Bradyrhizobiaceae* fam. nov. In: Bren DJ, editor. *Bergey's Manual of Systematic Bacteriology*, vol. 2. 2nd ed. New York: Springer-Verlag; 2005. p. 438.
79. Biological Agents: Technical rules for biological agents. <http://www.baua.de/en/Topics-from-A-to-Z/Biological-Agents/TRBA/TRBA.html>. Accessed 24 Feb 2017.
80. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene Ontology: Tool for the unification of biology. *Nat Genet.* 2000;25:25–9. The Gene Ontology Consortium.
81. Christie PJ, Atmakuri K, Krishnamoorthy V, Jakubowski S, Cascales E. Biogenesis, architecture, and function of bacterial type IV secretion systems. *Annu Rev Microbiol.* 2005;59:451–85.
82. *Bradyrhizobium elkanii* USDA 76 locus tag. <http://www.ncbi.nlm.nih.gov/bioproject/162247>. Accessed 24 Feb 2017.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

