UC Davis

UC Davis Previously Published Works

Title

Co-segregation analysis and mapping of the anthracnose Co-10 and angular leaf spot Phg-ON disease-resistance genes in the common bean cultivar Ouro Negro.

Permalink

<https://escholarship.org/uc/item/9kd27574>

Journal

TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik, 126(9)

ISSN

0040-5752

Authors

Gonçalves-Vidigal, MC
Cruz, AS
Lacanallo, GF
et al.

Publication Date

2013-09-01

DOI

10.1007/s00122-013-2131-8

Peer reviewed

Co-segregation analysis and mapping of the anthracnose *Co-10* and angular leaf spot *Phg-ON* disease-resistance genes in the common bean cultivar Ouro Negro

M. C. Gonçalves-Vidigal · A. S. Cruz · G. F. Lacanallo · P. S. Vidigal Filho · L. L. Sousa · C. M. N. A. Pacheco · P. McClean · P. Gepts · M. A. Pastor-Corrales

Departamento de Agronomia, Universidade Estadual de Maringá, Av. Colombo, 5790, 87020-900, Maringá, Paraná, Brazil; Genomics and Bioinformatics Program, Department of Plant Sciences, North Dakota State University, Fargo, ND 58105, USA; Section of Crop and Ecosystem Sciences, Department of Plant Sciences/MS1, University of California, Davis, CA 95616-8780, USA; Soybean Genomics and Improvement Laboratory, USDA-ARS, BARC-West, Beltsville, MD 20705, USA.

*Corresponding author: talo.pastor-corrales@ars.usda.gov

Received:

Abstract Anthracnose (ANT) and angular leaf spot (ALS) are devastating diseases of common bean (*Phaseolus vulgaris* L.). Ouro Negro is a highly productive common bean cultivar, which contains the *Co-10* and *Phg-ON* genes for resistance to ANT and ALS, respectively. In this study, we performed a genetic co-segregation analysis of resistance to ANT and ALS using an F₂ population from the Rudá × Ouro Negro cross and the F_{2:3} families from the AND 277 × Ouro Negro cross. Ouro Negro is resistant to races 7 and 73 of the ANT and race 63-39 of the ALS pathogens. Conversely, cultivars AND 277 and Rudá are susceptible to races 7 and 73 of ANT, respectively. Both cultivars are susceptible to race 63-39 of ALS. Co-segregation analysis revealed that *Co-10* and *Phg-ON* were inherited together, conferring resistance to races 7 and 73 of ANT and race 63-39 of ALS. The *Co-10* and *Phg-ON* genes co-segregated and were tightly linked at a distance of 0.0 cM on chromosome Pv04. The molecular marker g2303 was linked to *Co-10* and *Phg-ON* at a distance of 0.0 cM. Because of their physical linkage in a *cis* configuration, the *Co-10* and *Phg-ON* resistance alleles are inherited together and can be monitored with great efficiency using g2303. The close linkage between the *Co-10* and *Phg-ON* genes and prior evidence are consistent with the existence of a resistance gene cluster at one end of chromosome Pv04, which also contains the *Co-3* locus and ANT resistance quantitative trait loci. These results will be very useful for breeding programs aimed at developing bean cultivars with ANT and ALS resistance using marker-assisted selection.

Keywords: Co-segregation analysis, linkage mapping, common bean, anthracnose, angular leaf spot

26 Introduction

27

28 The common bean (*Phaseolus vulgaris* L.) is a very important human food, particularly
 29 in many countries of the Americas and eastern and southern Africa. In these countries,
 30 the common bean is recognized as an especially important source of protein, complex
 31 carbohydrates, fiber, and minerals in the diet of millions of individuals (Broughton et al.
 32 2003; Gepts et al. 2008). Moreover, the common bean is the most widely cultivated
 33 species of the genus *Phaseolus* and accounts for approximately 95% of the world's
 34 *Phaseolus* bean production. However, this crop is affected by several biotic, edaphic,
 35 and climatic factors that decrease its yields (Schwartz and Pastor-Corrales 1989).
 36 Among the biotic factors, diseases can cause severe yield losses and reduce the quality
 37 of dry and snap beans worldwide (Singh and Schwartz 2010). Anthracnose (ANT),
 38 caused by *Colletotrichum lindemuthianum* (Sacc. & Magnus) Lams.-Scrib., and angular
 39 leaf spot (ALS), caused by *Pseudocercospora griseola* (Sacc.) Crous and Braun (Crous
 40 et al. 2006), are among the most widespread diseases of the common bean in the tropics.
 41 Both ANT and ALS are widespread in Brazil and eastern and southern Africa, causing
 42 crop losses of up to 100% due to ANT and up to 70% due to ALS (Correa-Victoria et al.
 43 1989; Pastor-Corrales and Tu 1989; Singh and Schwartz 2010; Mahuku et al. 2009,
 44 2011; Oblessuc et al. 2012). The pathogens that cause ANT and ALS are characterized
 45 by extensive diversity in their virulence; each pathogen has many virulent strains, which
 46 are known as races.

47 Host resistance is the most cost-effective strategy for controlling ANT and ALS in
 48 the common bean. Disease-resistant common bean cultivars are easily adopted by
 49 farmers and do not generate environmental risks. Resistance to ANT and ALS is
 50 conditioned primarily by single dominant genes. Fourteen genes for resistance to ANT
 51 with the *Co* designation have been identified (Kelly and Vallejo 2004; Gonçalves-
 52 Vidigal et al. 2008, 2009, 2011, 2012a). Similarly, seven independent dominant genes
 53 identified by the *Phg* symbol have been reported to confer resistance to *P. griseola*
 54 (Caixeta et al. 2003, 2005; Gonçalves-Vidigal et al. 2011; Mahuku et al. 2009, 2011).
 55 However, the names of many of these genes have not been submitted to the Bean
 56 Improvement Cooperative Genetics Committee (<http://bic.css.msu.edu/Genetics.cfm>).
 57 Ouro Negro is a black-seeded Mesoamerican cultivar derived from the CIAT accession
 58 G 3680, also known as Honduras 35. This cultivar, which was introduced in Brazil in
 59 1991, is highly productive and exhibits desirable agronomic and cooking characteristics

(Alzate-Marin et al. 2003; Souza et al. 2011). In Brazil, Ouro Negro is very **resistant to specific races** of ANT, ALS, and rust and exhibits excellent morpho-agronomic traits. Thus, this cultivar has been recommended by the National Agricultural Research System (SNPA) for cultivation in several Brazilian states (Alzate-Marin et al. 2004).

Previous studies have identified the disease-resistance genes *Co-10* (ANT resistance), *Phg-ON* (ALS resistance), and *Ur-14* (rust resistance) in Ouro Negro. These three genes are considered very important for common bean breeding programs in Brazil (Alzate-Marin et al. 2003; Souza et al. 2011). Ouro Negro confers resistance to *C. lindemuthianum* races 7, 9, 23, 55, 64, 67, 73, 79, 81, 83, 87, 89, 95, 102, 117, 119, 343, 453, 1033, 1545, and 1600 and to *P. griseola* races 31-31, 31-55, 63-39, 63-47, 31-23, 63-55, and 63-63 (Faleiro et al. 2001; Alzate-Marin et al. 2003; Gonçalves-Vidigal and Kelly 2006; Gonçalves-Vidigal et al. 2009; Ragagnin et al. 2009). Thus, Ouro Negro is an important source of disease resistance in the bean breeding projects of Brazil and has been used to develop ANT-resistant dry bean cultivars (Alzate-Marin et al. 2003; Souza et al. 2011).

The common bean ($2n = 2x = 22$) consensus map is organized into 11 chromosomes (Nodari et al. 1993; Freyre et al. 1998; Pedrosa Harand et al. 2008). Gepts (1999) and Kelly et al. (2003) reviewed the development of integrated consensus maps of the 11 chromosomes in both the common bean and cowpea (*Vigna unguiculata* L. Walp). These authors reported the map locations of major resistance genes to bean rust, ANT, common bacterial blight, and white mold in gene clusters on chromosomes Pv01, Pv04, Pv07, and Pv11 in the common bean. Clusters of *Co*-ANT and *Ur*-rust resistance genes have been reported on Pv01, Pv04, and Pv011 (Geffroy et al. 1999; Miklas et al. 2002; Kelly and Vallejo 2004; Miklas et al. 2006; Geffroy et al. 2009). Among these, the Pv04 cluster, in which the *Co-10* (ANT) resistance gene resides, contains a large number of genes that confer resistance to various other common bean pathogens. The ANT resistance genes included in the Pv04 cluster are *Co-y*, *Co-z*, and *Co-15* of Andean origin and *Co-R_{VI}*, *Co-3* (previously also *Co-9*), and *Co-10* of Mesoamerican origin (Adam-Blondon et al. 1994; Geffroy et al. 1999, 2008; Rodríguez-Suárez et al. 2007; David et al. 2008; Gonçalves et al. 2010; Campa et al. 2011). Recent evidence has confirmed that *Co-15* is also located on Pv04 (Sousa 2012). Moreover, López et al. (2003) identified clusters of resistance genes on Pv04 that are effective against several strains of *C. lindemuthianum*, *P. griseola*, and the bean golden yellow mosaic virus.

93 Similar gene clusters have been described in other species. The Pv04 cluster
94 containing the *Co-10* and *Phg-ON* genes has a homologous cluster in the soybean
95 genome (Ashfield et al. 2012). Moreover, McClean et al. (2010) demonstrated that the
96 Pv04 region was bounded by molecular markers g1375 and g2685, and the interval
97 from 14-77 cM was **homologous to regions on soybean chromosomes** Gm13 and Gm19.
98 Other examples of disease-resistance clusters in soybean include those identified by
99 Meksem et al. (1999), Kang et al. (2012), and McHale et al. (2012). The presence of
100 these clusters is widespread among higher plants, as they are also observed in members
101 of the grass family, such as sorghum (Martin et al. 2011).

102 Previous studies on Ouro Negro have revealed that the *Co-10* ANT-resistance gene
103 and *Ur-14* rust-resistance gene are positioned 12.3 cM apart on Pv04 (Corrêa et al.
104 2000); however, the relationship between the *Co-10* and *Phg-ON* ALS-resistance genes
105 in Ouro Negro has not been established. To date, there has been only one report of a
106 linkage between the *Co* and *Phg* genes. Gonçalves-Vidigal et al. (2011) described the
107 co-segregation of the ANT *Co-1⁴* and ALS *Phg-1* disease-resistant genes in the
108 common bean cultivar AND 277, in which the genes were located in a cluster of
109 resistance genes on Pv01.

110 The objectives of the present study were a) to investigate the linkage between the
111 *Co-10* (ANT) and *Phg-ON* (ALS) disease-resistance genes and the existence of a
112 possible gene cluster involving these genes, b) to verify the linkage between the newly
113 identified molecular marker g2303 and the *Co-10* and *Phg-ON* genes, and c) to examine
114 the value of the g2303 marker in comparison to the previously identified SF10
115 molecular marker (Corrêa et al. 2000) for the selection of common bean plants
116 containing the *Co-10* and *Phg-ON* genes.

117

118

119 **Materials and Methods**

120

121 *Genetic crosses and co-segregation analysis*

122

123 To conduct the co-segregation and linkage studies, two different crosses were
 124 performed with the Ouro Negro cultivar, which contains the *Co-10* and *Phg-ON*
 125 disease-resistance genes that confer resistance to races 7 and 73 of *C. lindemuthianum*
 126 and race 63-39 of *P. griseola*. In the first cross, Ouro Negro was crossed with the
 127 common bean cultivar Rudá (susceptible to races 73 of *C. lindemuthianum* and 63-39 of
 128 *P. griseola*). In the second cross, Ouro Negro was crossed with the common bean
 129 cultivar AND 277 (susceptible to races 7 of *C. lindemuthianum* and 63-39 of *P.*
 130 *griseola*). Both AND 277 and Rudá were used as the female parents. The F₁ seeds were
 131 sown in polyethylene vases (48 × 30 × 11 cm) containing a mixture of previously
 132 fertilized and sterilized substrate. The plant vases were kept in a greenhouse until the F₂
 133 seeds were produced. A total of 112 F₂ seeds derived from the Rudá × Ouro Negro cross
 134 were sown in plastic trays (50 × 30 × 9 cm) containing peat-based substrate. The
 135 seedlings were maintained in a greenhouse until the first trifoliolate leaves (stage V3;
 136 Gepts 1987) were fully expanded. At that time, the plants were inoculated with race 73
 137 of *C. lindemuthianum* and race 63-39 of *P. griseola*.

138 A set of 63 F₂ seeds from the AND 277 × Ouro Negro cross was multiplied in pots to
 139 obtain the F₃ generation plants. The F_{2:3} families, obtained by selfing individual F₂
 140 plants, were used to characterize the corresponding F₂ plants for resistance to race 7 of
 141 *C. lindemuthianum* and race 63-39 of *P. griseola*. The dominant violet flower trait,
 142 which was inherited from the resistant male Ouro Negro parent, was observed in the F₁
 143 plants, confirming that they were hybrids. The resistance genotype of each F₂ plant was
 144 inferred from the phenotypes of the corresponding F_{2:3} families.

145

146 *Inocula of Colletotrichum lindemuthianum and Pseudocercospora griseola*

147

148 Races 7 and 73 of *C. lindemuthianum* were obtained from the mycology collection of
 149 NUPAGRI (Nucleo de Pesquisa Aplicada a Agricultura, Universidade Estadual de
 150 Maringá, State of Parana, Brazil), and *P. griseola* race 63-39 was kindly provided by
 151 Dr. M. Goreti de Almeida Oliveira of the “Instituto de Biotecnologia Aplicada à
 152 Agropecuária” (Bioagro), Universidade Federal de Viçosa, State of Minas Gerais,

153 Brazil. The classification as races 7, 73, and 63-39 was confirmed by inoculating two
 154 separate sets of 12 common bean differential cultivars used to characterize the virulence
 155 spectra of the ANT and ALS pathogens (Pastor-Corrales 1991). The initial inocula of
 156 races 7 and 73 of *C. lindemuthianum* and race 63-39 of *P. griseola* were obtained from
 157 monosporic cultures. The subsequent inocula of the ANT races were produced on young
 158 green common bean pod medium incubated at 22°C for 14 days. The inoculum of race
 159 63-39 of ALS was first multiplied in Petri dishes containing 1-2 mL of a solution of 800
 160 mL of sterilized water, 200 mL of commercial tomato sauce, 15 g of agar, 4.5 g of
 161 calcium carbonate (CaCO₃), and 10 µg mL⁻¹ of streptomycin (Sanglard et al. 2009).
 162 Subsequent inocula of race 63-39 were produced in Petri dishes containing the tomato
 163 medium and maintained in a BOD incubator at 24°C for 15 days. Spore suspensions
 164 were adjusted to 1.2×10^6 and 1.2×10^4 conidia mL⁻¹ for *C. lindemuthianum* and *P.*
 165 *griseola*, respectively.

167 *Inoculation and evaluation of the Ouro Negro × Rudá F₂ population*

169 The parental, F₁ generation, and 112 F₂ plants of the Rudá × Ouro Negro cross were
 170 simultaneously inoculated with race 73 of *C. lindemuthianum* and race 63-39 of *P.*
 171 *griseola* according to Gonçalves-Vidigal et al. (2001). After the expansion of the first
 172 trifoliolate leaf, the right leaflet was inoculated with *C. lindemuthianum* and the left
 173 leaflet with *P. griseola*. Each pathogen was inoculated separately using small brushes
 174 (Tigre® model 266, number 14). After inoculation, the plants were placed in a mist
 175 chamber for 48 h at a temperature of 20 ± 2°C with light controlled at 12 h of daylight
 176 and 12 h of darkness (light intensity of 300 µmoles m⁻² s⁻¹ at a height of 1 m) and a
 177 relative humidity of > 95%. After the incubation period, the inoculated plants were
 178 transferred to open-air benches at a temperature of 22°C with artificial light (12 h of
 179 daylight at 25°C), where they remained for 7 days before visual symptom assessment.
 180 The visual assessments of ANT and ALS symptoms were performed using the severity
 181 scales proposed by Pastor-Corrales et al. (1995) and Inglis et al. (1988), respectively,
 182 with scores of 1-9. Plants with disease reaction scores between 1 and 3 were considered
 183 resistant, whereas plants with scores from 4 to 9 were considered susceptible.

186 *Evaluation of the population from the AND 277 × Ouro Negro cross*

187
 188 For the AND 277 × Ouro Negro cross, 12 plants from each of the 63 F_{2:3} families
 189 derived from the F₂ population were also inoculated with races 7 of *C. lindemuthianum*
 190 and 63-39 of *P. griseola*. The seedlings were grown under natural light in greenhouses
 191 supplemented with 400 W high-pressure sodium lamps providing a total light intensity
 192 of 115 $\mu\text{moles m}^{-2} \text{s}^{-1}$ for 7 to 10 days until they reached the first trifoliate leaf stage.
 193 Twenty parental plants (Ouro Negro and AND 277), 20 F₁ plants, and 12 plants from
 194 each of the 63 F_{2:3} families were separately inoculated with races 7 of *C.*
 195 *lindemuthianum* and 63-39 of *P. griseola*. A spore suspension containing 2.0×10^6
 196 spores mL⁻¹ of race 7 of *C. lindemuthianum* was inoculated using a DeVilbiss number
 197 15 atomizer powered by an electric air compressor (Schulz, SA, Joinville, Santa
 198 Catarina, Brazil). A similar procedure was employed for the inoculation with race 63-39
 199 of *P. griseola*. The procedures used after the bean plant inoculation with the ANT and
 200 ALS pathogens and for symptom evaluation were identical to those used for the Rudá ×
 201 Ouro Negro population.

202

203 *DNA extraction and bulked segregant analysis*

204

205 The DNA extraction process was performed according to Afanador et al. (1993) with
 206 some modifications; the DNA was extracted from the central leaflet from the first
 207 trifoliate leaf directly with 400 μL of CTAB extraction buffer. A total of 23
 208 molecular markers, all mapping to chromosome Pv04 (PhaseolusGenes database:
 209 <http://phaseolusgenes.bioinformatics.ucdavis.edu>), were chosen for testing. The markers
 210 included 10 Sequence Tagged Sites (STS) (McConnell et al. 2010), eight microsatellites
 211 (Blair et al. 2003; Gaitán-Solis et al. 2002), and five SCARs (Corrêa et al. 2000; Corrêa
 212 et al. 2001; Queiroz et al. 2004). For each F₂ plant from the Rudá × Ouro Negro cross
 213 inoculated simultaneously with races 73 of *C. lindemuthianum* and 63-39 of *P. griseola*,
 214 the total genomic DNA was isolated from the newly emerged first trifoliate leaflet. The
 215 same procedure was used to isolate DNA from the F₂ plants from the AND 277 × Ouro
 216 Negro cross. The resistant or susceptible phenotype of these F₂ plants was inferred from
 217 the phenotypes of the 63 F_{2:3} families that were separately inoculated with races 7 of *C.*
 218 *lindemuthianum* and 63-39 of *P. griseola*. Two contrasting DNA bulks were
 219 constructed by pooling equal volumes of fluorometrically standardized DNA from five
 220 F₂ plants that were homozygous RR for the resistant genotype and rr for the susceptible

genotype (Michelmore et al. 1991). The resistant and susceptible DNA pools were used for bulked-segregant analysis (Michelmore et al. 1991) for the identification of markers potentially linked to the *Co-10* ANT and *Phg-ON* ALS-resistance genes. All markers were tested on the parental plants and the resistant and susceptible bulks.

Of the tested STS and microsatellite markers, only the STS g2303 was polymorphic, with contrasting amplification patterns in the parental materials and the resistant and susceptible bulks and individuals from the bulks. The g2303 molecular marker was retained for subsequent studies. Among the tested SCARs, only SF10 was polymorphic; thus, SF10 was chosen for genotyping the F₂ population derived from the Rudá × Ouro Negro cross. In addition, the SF10 marker is linked to the *Ur-14* rust and *Co-10* ANT-resistance genes of Ouro Negro, and this marker has been mapped to Pv04 (Corrêa et al. 2000).

All amplification reactions were performed with a thermal cycler (MJ Research Inc., Waltham, MA). The polymerase chain reaction (PCR) program for g2303 consisted of 3 min at 95°C and 35 cycles of 30 s at 92°C, 1 min at 50°C, 60 s at 72°C, followed by a 5 min extension at 72°C and 4 min at 4°C. PCRs were performed in 25 µL total reaction volumes containing 40 ng total DNA; 0.2 mM each dNTP; standard *Taq* buffer containing 1.5 mM MgCl₂ and 0.2 µM forward primer and reverse primer; and 1 unit of *Taq* DNA polymerase. Following the addition of 2 µL loading buffer (30% glycerol and 0.25% bromophenol blue), the PCR products for g2303 were analyzed on 6% polyacrylamide gels stained with SYBR Safe (0.02%). The PCR for SF10 consisted of 3 min at 94°C, 35 cycles of 15 s at 94°C, 1 min at 65°C, and 90 s at 72°C, followed by a 7 min extension at 72°C and 4 min at 4°C. The PCR products from SF10 were visualized on agarose gels. The DNA bands were visualized under ultraviolet light, and digital images were recorded with an L-PIX Image EX model (Loccus Biotecnologia - Loccus do Brasil, Cotia, SP, Brazil).

Molecular mapping

The molecular markers g2303 and SF10 were both analyzed in the BAT93 × Jalo EEP558 (BJ: 71 lines; Freyre et al. 1998) recombinant inbred lines and in the F₂ populations from the AND 277 × Ouro Negro and Rudá × Ouro Negro crosses. The primer sequences for a segregation the g2303 marker were as follows:

'GGGGCGGAATCAGGTTACCA' (forward) and

255 'GGTTTAGGACAACATAATGAGAGTGATGTACCGTGT' (reverse) (McConnell et al.
 256 2010), as described in the PhaseolusGenes database
 257 (<http://phaseolusgenes.bioinformatics.ucdavis.edu/markers/?ALL=g2303&format.html>).
 258 The primer sequences for SF10 were 'GGAAGCTTGGTGAGCAAGGA' (forward) and
 259 'GGAAGCTTGGCTATGATGGT' (reverse), according to the BIC database
 260 (http://www.css.msu.edu/bic/PDF/SCAR_Markers_2010.pdf).

261

262 *Statistical analyses*

263

264 Segregation analyses of the disease reactions of 112 F₂ plants from the Rudá × Ouro
 265 Negro cross were performed using the χ^2 test according to a Mendelian segregation
 266 hypothesis of 3 R (resistant) to 1 rr (susceptible). Additionally, this test was performed
 267 with the data from the 63 F_{2:3} families from the AND 277 × Ouro Negro cross according
 268 to a segregation hypothesis of 1:2:1 (RR:Rr:rr). A goodness-of-fit test for a 1:1
 269 segregation ratio was performed for the segregation of the g2303 and SF10 markers in
 270 the BJ population. Linkage analyses were performed using the MAPMAKER/EXP 3.0
 271 (Lincoln and Lander 1993) computer software to estimate the genetic distances between
 272 the g2303 and SF10 markers and the *Co-10* and *Phg-ON* genes in the F_{2:3} families
 273 derived from the AND 277 × Ouro Negro cross and F₂ population from the Rudá ×
 274 Ouro Negro cross, respectively. A minimum likelihood of an odds ratio score of ≥ 3.0
 275 and a maximum distance of 30 cM were used to test the linkages between these
 276 markers, as described by Freyre et al. (1998). The cluster containing the *Co-10/Phg-ON*
 277 and g2303 and SF10 markers was localized on Pv04 according to the standardized
 278 common bean linkage map nomenclature (Pedrosa-Harand et al. 2008). The map was
 279 drawn using MapChart software (Voorrips 2002).

280

281 **Results**

283 *Interaction between the two pathogens used for bean plant co-inoculation*

285 To detect a possible interaction between the two pathogens used in this study, we
 286 inoculated Ouro Negro (resistant to races 73 and 63-39) and Rudá (susceptible to races
 287 73 and 63-39) singly with race 73 or race 63-39 of the ANT and ALS pathogens,
 288 respectively. We also inoculated these cultivars simultaneously with the same two
 289 races; the right leaflet was inoculated with *C. lindemuthianum*, and the left leaflet was
 290 inoculated with *P. griseola*. In both inoculation schemes, Ouro Negro was resistant and
 291 Rudá was susceptible to races 73 and 63-39. These results confirm previous
 292 observations by Gonçalves-Vidigal et al. (2012b). These authors inoculated singly the
 293 Ouro Negro (resistant to race 73 and susceptible to 63-23), Mexico 54 and Cornell 49-
 294 242 (susceptible to race 73 and resistant to race 63-23), and Rudá (susceptible to both
 295 races) cultivars with race 73 or 63-23 of the ANT and ALS pathogens. Simultaneous
 296 inoculation was also performed using the same cultivars with the 73 and 63-23 races. In
 297 both inoculation systems, Ouro Negro was resistant to race 73 and susceptible to 63-23;
 298 Mexico 54 and Cornell 49-242 were susceptible to race 73 and resistant to race 63-23,
 299 and Rudá was susceptible to both races. The results indicated that there was no
 300 interaction between these pathogens when inoculated simultaneously on the same bean
 301 plants.

303 *Genetic resistance and co-segregation of the Co-10 and Phg-ON genes*

305 A total of 112 F₂ plants derived from the Rudá × Ouro Negro cross and 20 plants each
 306 from the parental and F₁ generation were inoculated simultaneously with race 73 of *C.*
 307 *lindemuthianum* and race 63-39 of *P. griseola*. Nine days after the inoculation of the
 308 plants with both pathogens, a segregation was observed in the F₂ population, with 84
 309 plants resistant and 28 susceptible to *C. lindemuthianum* ($P = 1.0$) and 83 plants
 310 resistant and 29 susceptible ($P = 0.83$) to *P. griseola* (Table 1). Except for one plant, the
 311 112 F₂ plants from the Rudá × Ouro Negro cross exhibited identical resistant and
 312 susceptible phenotypes; that is, all but one of the 112 F₂ plants that were resistant to the
 313 ANT pathogen were also resistant to the ALS pathogen. Similarly, plants that were
 314 susceptible to the ANT pathogen were also susceptible to the ALS pathogen.

315

316 **Insert Table 1**

317

318 The segregation observed in the 63 F_{2:3} families derived from the AND 277 × Ouro
 319 Negro cross inoculated separately with races 7 of *C. lindemuthianum* and 63-39 of *P.*
 320 *griseola* exhibited a similar co-segregation of resistance/susceptibility to both pathogens
 321 (Table 2). All plants that were resistant to ANT were also resistant to the ALS pathogen,
 322 and the plants that were susceptible to ANT were also susceptible to ALS. Thus, no
 323 recombinants were observed in these F_{2:3} families, suggesting that the *Co-10* and *Phg-*
 324 *ON* genes are very tightly linked. The 63 F_{2:3} families segregated into classes as follows:
 325 14RR:34Rr:15rr ($P = 0.81$). These results revealed a co-segregation between the *Co-10*
 326 and *Phg-ON* genes that fit a 1R:2Rr:1S ratio for a single dominant gene.

327 As noted earlier, except for one plant, the 112 F₂ plants from the Rudá × Ouro Negro
 328 cross also exhibited identical resistant or susceptible phenotypes. Thus, the segregation
 329 results from both the Rudá × Ouro Negro cross and from the AND 277 × Ouro Negro
 330 cross suggest that the *Co-10* and *Phg-ON* genes are very tightly linked. These
 331 segregation patterns fit the model of a 3R:1S ratio of monogenic, dominant resistance.
 332 Therefore, it was concluded that the resistance to races 7 and 73 of *C. lindemuthianum*
 333 and 63-39 of *P. griseola* in the Mesoamerican common bean cultivar Ouro Negro is
 334 conferred by the tightly linked *Co-10* and *Phg-ON* genes. Other researchers have
 335 previously conducted separate ANT and ALS inheritance studies and observed that
 336 Ouro Negro exhibited dominant and monogenic inheritance of resistance to each of the
 337 pathogens causing these diseases (Corrêa et al. 2001; Alzate-Marin et al. 2003).

338

339 **Insert Table 2**

340

341 *Marker analysis and mapping of the Co-10 and Phg-ON genes*

342

343 In order to identify a molecular marker linked to the *Co-10* and *Phg-ON* resistance
 344 genes and determine the chromosomal location of these genes, 23 molecular markers
 345 previously mapped on Pv04 were evaluated using bulked segregant analysis
 346 (Michelmore et al. 1991). Of the 23 tested molecular markers, only g2303 and SF10
 347 exhibited polymorphisms characterized by contrasting amplification patterns in the
 348 parental materials and the resistant versus susceptible bulks or individuals and were

chosen for further studies. The SF10 marker amplified a band of 1072 bp in the genomic region of interest in the resistant Ouro Negro parental plants and the resistant bulk. The g2303 marker amplified a 350 bp fragment in the resistant parental Ouro Negro plants and in all resistant F₂ plants from the Rudá × Ouro Negro and AND 277 × Ouro Negro crosses. The presence of the 350 bp amplicon of g2303 in Ouro Negro (*Co-10/Phg-ON*), Mexico 222 (*Co-3*), and BAT93 (*Co-3³*) is presented in Figure 1; conversely, this marker was absent in the Corinthiano (*Co-15*) and Rudá cultivars.

As illustrated in Table 3, the co-segregation of resistance versus susceptibility to ALS and ANT and the SF10 marker in the F₂ population from the Rudá × Ouro Negro cross revealed a segregation pattern of 92(+):20(-), indicating that SF10 is linked to the *Co-10* and *Phg-ON* loci at a distance of 7.8 cM.

Insert Table 3 and Figure 1

The results of the co-segregation analyses of the F_{2:3} families from the AND 277 × Ouro Negro cross inoculated separately with ANT and ALS and the molecular analysis using the g2303 marker are depicted in Table 4. The genetic linkage analysis resulted in a segregation of 48(+):15(-), indicating a good fit to the expected ratio of 3R:1S ($P = 0.83$). All 48 resistant plants possessed the g2303 molecular marker, and the marker was absent in the 15 susceptible plants. The lack of recombinants among the *Co-10* and *Phg-ON* genes and the g2303 marker indicated that g2303 is tightly linked to these genes and can be used to map the *Co-10* and *Phg-ON* resistance genes of Ouro Negro.

Figure 2 presents the distances and locations of the *Co-10* (ANT resistance) and *Phg-ON* (ALS resistance) genes and the g2303 and SF10 markers on Pv04. The linkage analysis revealed that the g2303 marker was linked to the ANT and ALS genes at a distance of 0.0 cM on Pv04 and that the SF10 marker was linked at a distance of 7.8 cM.

Insert Table 4 and Figure 2

Marker g2303, which was previously mapped to chromosome Pv04 (McConnell et al. 2010), was tested in the BAT93/Jalo EEP558 (BJ) RI population, resulting in a segregation of 36(+):37(-) ($\chi^2 = 0.013$; $P = 0.90$), for a good fit to a 1R:1S ratio. In addition, the SF10 marker, positioned on chromosome Pv04 (Corrêa et al. 2000), was

383 also assessed in the same mapping population, yielding a ratio of 38(+):35 (-) ($\chi^2 =$
384 0.12; $P = 0.72$) and tight linkage with g2303 (Figure 2).
385

Discussion

In this study, to better understand the existing association between the *Co-10* and *Phg-ON* genes and the trait of simultaneous resistance to two major common bean pathogens, *C. lindemuthianum* and *P. griseola*, we performed co-segregation and linkage analyses of the F₂ population from the Rudá × Ouro Negro cross and the F_{2:3} families from the AND 277 × Ouro Negro cross.

The co-segregation between the *Co-10* and *Phg-ON* resistance genes for the ANT and ALS pathogens, respectively, was detected when the F₂ population from the Rudá (S) × Ouro Negro (R) cross was simultaneously inoculated with both pathogens. We observed only one recombinant in this F₂ population, suggesting a tight linkage (0.9 cM) between the *Co-10* ANT and *Phg-ON* ALS-resistance genes in Ouro Negro. These results also indicated that the resistance of Ouro Negro to race 73 of *C. lindemuthianum* and race 63-39 of *P. griseola* was linked to the g2303 and SF10 molecular markers. As expected, the genotyping of the SF10 marker in this F₂ population revealed that it is linked to the *Co-10* and *Phg-ON* loci at a distance of 7.8 cM. A previous study reported that the SF10 marker was linked to the *Ur-14* rust and *Co-10* ANT-resistance genes on chromosome Pv04 at a distance of 6.0 ± 1.3 cM (Corrêa et al. 2000).

Additionally, to obtain more precise phenotypic results, we evaluated 63 F_{2:3} families from the AND 277 (S) × Ouro Negro (R) cross inoculated separately with race 7 of *C. lindemuthianum* and race 63-39 of *P. griseola*. Each of the 63 F_{2:3} families exhibited identical responses to both pathogens. The data for the segregation of these F_{2:3} families revealed a tight linkage between the *Co-10* and *Phg-ON* genes. The segregation obtained for both pathogens fit the expected ratio of 1RR:2Rr:1rr. No recombinants were observed during the evaluation of the F_{2:3} families, revealing a tight linkage of the two genes.

The molecular analysis of F₂ population corresponding to the above mentioned F_{2:3} families revealed close linkage between the resistance genes and g2303 maker (0.0 cM). To date, SF10 is the only mapped marker linked to Ouro Negro resistance genes (Corrêa et al. 2000). However, we identified g2303 as a molecular marker that is much more tightly linked to the *Co-10* ANT and *Phg-ON* ALS disease-resistance genes.

A previous study involving ANT and ALS established that the ANT *Co-1⁴* and ALS *Phg-1* resistance genes, present in the AND 277 cultivar, co-segregated to form a cluster of resistance genes on Pv01 (Gonçalves-Vidigal et al. 2011). In the course of the present

study, we identified a tight linkage between the resistance genes *Co-10* and *Phg-ON*. These genes are also closely linked to a third gene, the *Ur-14* rust-resistance gene (Corrêa et al. 2000), suggesting the existence of a resistance gene cluster on chromosome Pv04. Prior studies have suggested that the majority of R (resistance) genes reside in clusters, and the frequency of recombination between clustered genes can vary strikingly, even within a single cluster (McDowell and Simon 2006). David et al. (2008) confirmed that the B4 (Pv04) R gene cluster is very large, spanning several megabases. Moreover, Oblessuc et al. (2012) reported the existence of different ANT-resistance genes clustered in the bean genome. In addition, David et al. (2008) observed that this Pv04 cluster contained genes conferring resistance not only to *C. lindemuthianum* but also to other pathogens, such as *Uromyces appendiculatus* (rust) and *Pseudomonas syringae* pv. *phaseolicola* (halo blight).

Geffroy et al. (1999, 2000) and Méndez-Vigo et al. (2005) were among the first to describe the *Co-3* gene cluster, which confers specific resistance to different races of *C. lindemuthianum* in the Mesoamerican cultivars Mexico 222 and BAT93. According to Campa et al. (2011), the gene present in the Andean cultivar Michigan Dark Red Kidney confers resistance to race 1545 of *C. lindemuthianum* and is located within the *Co-3* cluster on Pv04. The *Co-10/Phg-ON* resistance cluster present in Ouro Negro may correspond to the ANT-resistance cluster *Co-3*, as both gene clusters are very closely linked to the molecular marker g2303 on Pv04. However, previously conducted tests in an F₂ population from the Ouro Negro (*Co-10*) × Mexico 222 (*Co-3*) cross, inoculated with race 23 of *C. lindemuthianum*, revealed a lack of allelism (Alzate-Marin et al. 2003). In addition, the resistance spectrum of *Co-10* to 21 races of *C. lindemuthianum* shows only 28.5% similarity with the resistance of *Co-3*.

To verify the independence of the newly identified gene cluster *Co-10/Phg-ON*, electrophoretic analysis of the amplification products was conducted to test the association of the g2303 marker with the *Co-10/Phg-ON* and *Co-39* gene clusters and the *Co-15* resistance gene. The g2303 marker was assessed in the following cultivars: Corinthiano (*Co-15*), Ouro Negro (*Co-10/Phg-ON*), BAT93 (*Co-3³*), Mexico 222 (*Co-3*), and Rudá. These results revealed an association between g2303 and the *Co-10/Phg-ON* and *Co-3* gene clusters but not with the *Co-15* gene. The *Co-15* gene, present in Corinthiano, is linked to g2685¹⁵⁰, a marker that also maps on Pv04 (Sousa, 2012). A previous study conducted by McConnell et al. (2010) determined that the two markers g2303³⁵⁰ and g2685¹⁵⁰ are unlinked (58 cM) on chromosome Pv04. The physical

position in bp on chromosome 04 also confirms a loose association between the two markers as g2303 maps at position 3,356,300 bp and g2685 at position 9,078,200 bp (out of a total chromosome length of 45,960,019 bp; PhaseolusGenes). Similarly, allelism tests conducted by Gonçalves et al. (2010) confirmed that *Co-15* is distinct from the *Co-3*, *Co-3³*, and *Co-10* resistance genes. These data suggest the presence of a large gene cluster containing the resistance genes *Co-3*, *Co-9* (now re-labeled *Co-3³*), *Co-10*, and *Phg-ON*. An independent locus, *Co-15*, is placed on a distinct chromosome region of Pv04.

The main objective of this study was to verify the presence of a resistance gene cluster in Ouro Negro that not only confers resistance to ANT but also to ALS, regardless of how it should be properly named. In addition, a much more effective molecular marker, g2303, was found linked to the above mentioned gene clusters (*Co-3/Co-10/Phg-ON*) and it should be recommended to breeding programs. Whether *Co-10* should be considered an additional allele of the *Co-3* locus (*Co-3⁴*?) and *Phg-ON* should be re-labeled as *Phg-3*, should be decided by the Genetics Committee of the Bean Improvement Cooperative.

The results of this study will be very useful for breeding programs wishing to develop bean cultivars with combined ANT and ALS resistance using marker-assisted selection. The g2303 marker, in particular, will reduce the time and cost of pyramiding the *Co-10* and *Phg-ON* genes into commercial common bean cultivars.

Acknowledgements

This research was financially supported by the National Council for Scientific and Technological Development (CNPq). Anelise S. Cruz was supported by a scholarship from Capes. We would like to thank Andressa Gonçalves Vidigal for additional review of the manuscript. M.C. Gonçalves-Vidigal and P.S. Vidigal Filho are grateful for grants from CNPq.

References

- Adam-Blondon AF, Seignac M, Dron M, Bannerot H (1994) A genetic map of common bean to localize specific resistance genes against anthracnose. Genome 37:915-924

- 486 Afanador LK, Haley SD, Kelly JD (1993) Adoption of a ‘mini-prep’ DNA extraction
487 protocol for RAPD marker analysis in common bean (*Phaseolus vulgaris* L.). Ann
488 Rep Bean Improv Coop 36:10–11
- 489 Alzate-Marin AL, Costa MR, Arruda KM, Barros EG, Moreira MA (2003)
490 Characterization of the anthracnose resistance gene present in Ouro Negro
491 (Honduras 35) common bean cultivar. Euphytica 133:165-169
- 492 Alzate-Marin AL, Souza TLPO, Ragagnin VA, Moreira MA, Barros EG (2004)
493 Allelism tests between the rust resistance genes present in common bean cultivar
494 Ouro Negro and genes *Ur-5* and *Ur-11*. J Phytopathology 152: 60-64
- 495 Ashfield T, Egan AN, Pfeil BE, Chen NWG, Podicheti R, Ratnaparkhe MB, Ameline-
496 Torregrosa C, Denny R, Cannon S, Doyle JJ, Geffroy V, Roe BA, Saghai Maroof
497 MA, Young ND, Innes RW (2012) Evolution of a complex disease resistance gene
498 cluster in diploid *Phaseolus* and tetraploid *Glycine*. Plant Physiol 159:336-354
- 499 Blair MW, Pedraza F, Buendia HF, Gaitán-Solís E, Beebe SE, Gepts P, Tohme J (2003)
500 Development of a genome-wide anchored microsatellite map for common bean
501 (*Phaseolus vulgaris* L.). Theor Appl Genet 107:1362–1374
- 502 Broughton WJ, Hernandez G, Blair M, Beebe S, Gepts P, Vanderleyden J (2003) Beans
503 (*Phaseolus* spp.) - model food legumes. Plant Soil 252:55-128
- 504 Caixeta ET, Borém A, Niestche S, Moreira MA, Barros EG (2003) Inheritance of
505 angular leaf spot resistance in common bean line BAT 332 and identification of
506 RAPD markers linked to the resistance gene. Euphytica 134:297-303
- 507 Caixeta ET, Borém A, Alzate-Marin AL, Fagundes S, Morais SMG, Barros EG,
508 Moreira MA (2005) Allelic relationships for genes that confer resistance to
509 angular leaf spot in common bean. Euphytica 145:237-245
- 510 Campa A, Giraldez R, Ferreira JJ (2011) Genetic analysis of the resistance to eight
511 anthracnose races in the common bean differential cultivar Kaboon.
512 Phytopathology 101:757-764

- 513 Corrêa RX, Costa MR, Good-God PI, Ragagnin VA, Faleiro FG, Moreira MA, Barros
514 EG (2000) Sequence characterized amplified regions linked to rust resistance
515 genes in the common bean. *Crop Sci* 40:804-807
- 516 Corrêa RX, Good-God PI, Oliveira MLP, Nietsche S, Moreira MA, Barros EG (2001)
517 Herança da resistência à mancha-angular do feijoeiro e identificação de
518 marcadores moleculares flanqueando o loco de resistência. *Fitopatol Bras* 26:27-
519 32
- 520 Correa-Victoria FJ, Pastor-Corrales MA, Saettler AW (1989) Angular leaf spot. In:
521 Schwartz HF, Pastor-Corrales MA (eds). *Bean production problems in the tropics*,
522 2nd edn. CIAT, Cali, Colombia, pp 59-76
- 523 Crous PW, Liebenberg MM, Braun U, Groenewald JZ (2006) Re-evaluating the
524 taxonomic status of *Phaeoisariopsis griseola*, the causal agent of angular leaf spot
525 of bean. *Stud Mycol* 55:163-173
- 526 David P, Sévignac M, Thareau V, Catillon Y, Kami J, Gepts P, Langin T, Geffroy V
527 (2008) BAC end sequences corresponding to the B4 resistance gene cluster in
528 common bean: a resource for markers and synteny analyses. *Mol Genet Genomics*
529 280:521-533
- 530 Faleiro FG, Nietsche S, Ragagnin VA, Borém A, Moreira MA, Barros EG (2001)
531 Resistance of common bean rust and angular leaf spot under greenhouse
532 conditions. *Fitopatol Bras* 26: 86-89
- 533 Freyre R, Skrock PW, Geffroy V, Adam-Blondon AF, Shirmohamadali A, Johnson
534 WC, Llaca V, Nodari RO, Pereira PA, Tsai SM, Tohme J, Dron M, Nienhuis J,
535 Vallejo CE, Gepts P (1998) Towards an integrated linkage map of common
536 bean.4. Development of a core linkage map and alignment of RFLP maps. *Theor*
537 *Appl Genet* 97:847-856
- 538 Gaitán-Solís E, Duque MC, Edwards KJ, Tohme J (2002) Microsatellite repeats in
539 common bean (*Phaseolus vulgaris*): isolation, characterization and cross-species
540 amplification in *Phaseolus* ssp. *Crop Sci* 42:2128-2136

- 541 Geffroy V, Sicard D, Oliveira JCF, Sévignac M, Cohen S, Gepts P, Neema C, Langin T,
 542 Dron M (1999) Identification of an ancestral resistance gene cluster involved in
 543 the coevolution process between *Phaseolus vulgaris* and its fungal pathogen
 544 *Colletotrichum lindemuthianum*. Mol Plant Microbe Interact 12:774-784
- 545 Geffroy V, Sévignac M, De Oliveira JCF, Fouilloux G, Skroch P (2000) Inheritance of
 546 partial resistance against *Colletotrichum lindemuthianum* in *Phaseolus vulgaris*
 547 and co-localization of quantitative trait loci with genes involved in specific
 548 resistance. Mol Plant Microbe Interact 13: 287-296
- 549 Geffroy V, Sévignac M, Billant P, Dron M, Langin T (2008) Resistance to
 550 *Colletotrichum lindemuthianum* in *Phaseolus vulgaris*: a case study for mapping
 551 two independent genes. Theor Appl Genet 116:407-415
- 552 Geffroy V, Macadre C, David P, Pedrosa-Harand A, Sévignac M, Dauga C, Langin T
 553 (2009) Molecular analysis of a large subtelomeric nucleotide-binding-site-leucine-
 554 rich-repeat family in two representative genotypes of the major gene pools of
 555 *Phaseolus vulgaris*. Genetics 181:405-419
- 556 Gepts P (1987) Characterizing plant phenology. In: Wisiol K, Hesketh J (eds) Plant
 557 growth modeling for resource management. CRC Press, Boca Raton, FL, pp 3-24
- 558 Gepts P (1999) Development of an integrated genetic linkage map in common bean
 559 (*Phaseolus vulgaris* L.) and its use. In: Singh S (ed) Bean breeding for the 21st
 560 century. Kluwer, Dordrecht, the Netherlands, pp 53-91, 389-400
- 561 Gepts P, Aragão F, de Barros E, Blair MW, Brondani R, Broughton W, Galasso I,
 562 Hernández G, Kami J, Lariguet P, McClean P, Melotto M, Miklas P, Pauls P,
 563 Pedrosa-Harand A, Porch T, Sánchez F, Sparvoli F, Yu K (2008) Genomics of
 564 *Phaseolus* beans, a major source of dietary protein and micronutrients in the
 565 tropics. In: Moore PH, Ming R (eds). Genomics of tropical crop plants. Springer,
 566 pp 113-143
- 567 Gonçalves AMO, Gonçalves-Vidigal MC, Vidigal Filho PS, Poletine JP, Lacanallo GF,
 568 Coimbra GK (2010) Characterization of the Anthracnose Resistance Gene in

- 569 Andean Common Bean Corinthiano Cultivar. Ann Rep Bean Improv Coop
570 53:220-221
- 571 Gonçalves-Vidigal MC, Sakiyama NS, Vidigal Filho PS, Amaral JR AT, Poletine JP,
572 Oliveira VR (2001) Resistance of common bean cultivar AB 136 races 31 and 69
573 of *Colletotrichum lindemuthianum*: the *Co-6* locus. Crop Breed Appl Biotech
574 1:99-104
- 575 Gonçalves-Vidigal MC, Kelly JD (2006) Inheritance of anthracnose resistance in the
576 common bean cultivar Widusa. Euphytica 151:411-419
- 577 Gonçalves-Vidigal MC, Lacanallo, GF, Vidigal Filho PS (2008) A new gene conferring
578 resistance to anthracnose in Andean common bean (*Phaseolus vulgaris* L.)
579 cultivar 'Jalo Vermelho'. Plant Breed 127:592-596
- 580 Gonçalves-Vidigal MC, Vidigal Filho PS, Medeiros AF, Pastor-Corrales MA (2009)
581 Common bean landrace Jalo Listras Pretas is the source of a new Andean
582 anthracnose resistance gene. Crop Sci 49:133-138
- 583 Gonçalves-Vidigal MC, Cruz AS, Garcia A, Kami J, Vidigal Filho PS, Sousa LL,
584 McClean P, Gepts P, Pastor-Corrales MA (2011) Linkage mapping of the *Phg-1*
585 and *Co-1⁴* genes for resistance to angular leaf spot and anthracnose in the common
586 bean cultivar AND 277. Theor Appl Genet 122:893-903
- 587 Gonçalves-Vidigal MC, Meirelles AC, Poletine JP, Sousa LL, Cruz AS, Nunes MP,
588 Lacanallo GF, Vidigal Filho PS (2012a) Genetic analysis of anthracnose resistance
589 in 'Pitanga' dry bean cultivar. Plant Breed 131:423-429
- 590 Gonçalves-Vidigal MC, Vidigal Filho PS, Nogueira LM, Reck SAC, Oliveira MB,
591 Motomiya WR, Lacanallo GF, Cruz AS, Grunvald Ak, Nunes MPBP, Sousa LL,
592 Pastor-Corrales MA (2012b) Reaction of common bean cultivars inoculated
593 simultaneously with *Colletotrichum lindemuthianum* and *Pseudocercospora*
594 *griseola*. Ann Rep Bean Improv Coop 55: 97-98
- 595 Inglis DA, Hagedorn J, Rand RE (1988) Use of dry inoculum to evaluate beans for
596 resistance to anthracnose and angular leaf spot. Plant Dis 72:771-774

- 597 Kang Y, Kim K, Shim S, Yoon M, Sun S, Kim M, Van K, Lee S-H (2012) Genome-
598 wide mapping of NBS-LRR genes and their association with disease resistance in
599 soybean. *BMC Plant Biology* 12:139
- 600 Kelly JD, Gepts P, Miklas PN, Coyne DP (2003) Tagging and mapping of genes and
601 QTL and molecular marker-assisted selection for traits of economic importance in
602 bean and cowpea. *Field Crops Res* 82:135-154
- 603 Kelly JD, Vallejo VA (2004) A comprehensive review of the major genes conditioning
604 resistance to anthracnose in common bean. *HortScience* 39:1196-1207
- 605 Lincoln SE, Lander SL (1993) Mapmaker/exp 3.0 and Mapmaker/QTL 1.1. Whitehead
606 Inst. of Med. Res. Tech Report, Cambridge, MA
- 607 Lopez CE, Acosta IF, Jara C, Pedraza P, Gaitan-Solis E, Gallego G, Beebe S, Thome J
608 (2003) Identifying resistance gene analogs associated with resistances to different
609 pathogens in common bean. *Phytopathology* 93:88-95
- 610 Mahuku GS, Henriquez MA, Montoya C, Jara C, Teran H, Beebe S (2011) Inheritance
611 and development of molecular markers linked to angular leaf spot resistance genes
612 in the common bean accession G10909. *Mol Breed* 28: 57-71
- 613 Mahuku GS, Iglesias AM, Jara C (2009) Genetics of angular leaf spot resistance in the
614 Andean common bean accession G5686 and identification of markers linked to the
615 resistance genes. *Euphytica* 167: 381-396.
- 616 Martin T, Biruma M, Fridborg I, Okori P, Dixelius C (2011) A highly conserved NB-
617 LRR encoding gene cluster effective against *Setosphaeria turcica* in sorghum.
618 *BMC Plant Biol* 11:151
- 619 McClean P, Mamidi S, McConnell M, Chikara S, Lee R (2010) Syntenic mapping
620 between common bean and soybean reveals extensive blocks of shared loci. *BMC*
621 *Genomics* 11:184
- 622 McConnell M, Mamidi S, Lee R, Chikara S, Rossi M, Papa R, McClean P (2010)
623 Syntenic relationships among legumes revealed using a gene-based genetic

- linkage map of common bean (*Phaseolus vulgaris* L.) Theor Appl Genet 121: 1103-1116
- McDowell JM, Simon SA (2006) Recent insights into R gene evolution. Mol Plant Pathol 5: 437-448
- McHale LK, Haun WJ, Xu WW, Bhaskar PB, Anderson JE, Hyten DL, Gerhardt DJ, Jeddeloh JA, Stupar RM (2012) Structural variants in the soybean genome localize to clusters of biotic stress-response genes. Plant Physiol 159:1295-1308
- Meksem K, Doubler TW, Chanchaoenchai K, Nijti N, Chang SJ, Arelli APR, Cregan PE, Gray LE, Gibson PT, Lightfoot DA (1999) Clustering among loci underlying soybean resistance to *Fusarium solani*, SDS and SCN in near-isogenic lines. Theor Appl Genet 99:1131-1142
- Méndez-Vigo B, Rodríguez-Suárez C, Pañeda A, Ferreira J J, Giraldez R (2005) Molecular markers and allelic relationships of anthracnose resistance gene cluster B4 in common bean. Euphytica 141:237-245
- Michelmore RW, Paran I, Kesseli RV (1991) Identification of markers linked to disease resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions using segregating populations. Proc Natl Acad Sci USA 88:9828-9832
- Miklas PN, Pastor-Corrales MA, Jung G, Coyne DP, Kelly JD, McClean PE, Gepts P (2002) Comprehensive linkage map of bean rust resistance genes. Ann Rep Bean Improv Coop 45:125-129
- Miklas PN, Hu J, Grunwald NJ, Larsen KM (2006) Potential application of TRAP (Targeted Region Amplified Polymorphism) markers of mapping and tagging disease resistance traits in common bean. Crop Sci 46:910-916
- Nodari RO, Tsai SM, Guzmán P, Gilbertson RL, Gepts P (1993) Toward an integrated linkage map of common bean. III. Mapping genetic factors controlling host-bacteria interactions. Genetics 134:341-350

- 651 Oblessuc PR, Baroni RM, Garcia AAF, Chioratto AF, Carbonell SAM, Camargo LEA,
652 Benchimol LL (2012) Mapping of angular leaf spot resistance QTL in common
653 bean (*Phaseolus vulgaris* L.) under different environments. BMC Genetics 13: 50
- 654 Pastor-Corrales MA, Tu JC (1989) Anthracnose. In: Schwartz HF, Pastor-Corrales MA
655 (eds) Bean production problems in the tropics. CIAT, Cali, Colombia, pp 77-104
- 656 Pastor-Corrales MA (1991). Estandarización de variedades diferenciales y de
657 designación de razas de *Colletotrichum lindemuthianum*. Phytopathology 81:694.
- 658 Pastor-Corrales MA, Otoyá MM, Molina A, Singh SP (1995) Resistance to
659 *Colletotrichum lindemuthianum* isolates from Middle America and Andean South
660 America in different common bean races. Plant Dis 79:63-67
- 661 Pedrosa-Harand A, Porch T, Gepts P (2008) Standard nomenclature for common bean
662 chromosomes and linkage groups. Ann Rep Bean Improv Coop 51:106–107
- 663 Queiroz VT, Sousa CS, Costa MR, Sanglard DA, Arruda KMA, Souza TLPO,
664 Ragagnin VA, Barros EG, Moreira MA (2004) Development of SCAR markers
665 linked to common bean angular leaf spot resistance genes. Ann Rep Bean Improv
666 Coop 47:237-238
- 667 Ragagnin VA, Souza TLPO, Sanglard DA, Arruda KMA, Costa MR, Alzate-Marin AL,
668 Carneiro JES, Moreira MA, Barros G (2009) Development and agronomic
669 performance of common bean lines simultaneously resistant to anthracnose,
670 angular leaf spot and rust. Plant Breeding 128:156-163
- 671 Rodríguez-Suárez C, Méndez-Vigo B, Pañeda A, Ferreira JJ, Giraldez R (2007) A
672 genetic linkage map of *Phaseolus vulgaris* L. and localization of genes for specific
673 resistance to six races of anthracnose (*Colletotrichum lindemuthianum*). Theor
674 Appl Genet 114:713-722
- 675 Sanglard DA, Balbi BP, de Barros EG (2009) An efficient protocol for isolation,
676 sporulation and maintenance of *Pseudocercospora griseola*. Ann Rep Bean
677 Improv Coop 52:62-63

- 678 Schwartz HF, Pastor-Corrales MA (1989) Bean production problems in the tropics.
679 Cali: CIAT, p.105-157
- 680 Singh SP, Schwartz F (2010) Breeding common bean for resistance to diseases: a
681 review. Crop Sci 50:2199-2223
- 682 Sousa LL (2012) Molecular mapping of the alleles *Co-5²* and *Co-15* of resistance to
683 *Colletotrichum lindemuthianum* in common bean. PhD Thesis (Genetic and Plant
684 Breeding). Maringá: Universidade Estadual de Maringá, 2012. 93p.
- 685 Souza TLPO, Dessaune SN, Sanglard DA, Moreira MA, Barros EG (2011)
686 Characterization of the rust resistance gene present in the common bean cultivar
687 Ouro Negro, the main rust resistance source used in Brazil. Plant Pathol 60:839-
688 845
- 689 Voorrips RE (2002) Mapchart: software for the graphical presentation of linkage maps
690 and QTLs. J Hered 93:77-78

1

Table 1 Segregation for resistance to races 73 of *Colletotrichum lindemuthianum* and 63-39 of *Pseudocercospora griseola* in common bean F₂ plants from the Rudá × Ouro Negro cross

Parental cross	Generation	Observed	Expected	χ^2	<i>P</i> value (1 df)
		Ratio (R:S) ^a	Ratio (3R:1S)		

Race 73 of <i>Colletotrichum lindemuthianum</i>					
Ouro Negro	RP ^b	20:0			
Rudá	SP ^c	0:20			
Rudá × Ouro Negro	F ₁	20:0			
Rudá × Ouro Negro	F ₂	84:28	84:28	0.00	1.0

Race 63-39 of <i>Pseudocercospora griseola</i>					
Ouro Negro	RP ^b	20:0			
Rudá	SP ^c	0:20			
Rudá × Ouro Negro	F ₁	20:0			
Rudá × Ouro Negro	F ₂	83:29	84:28	0.04	0.83

^aR = Resistant; S = Susceptible; RP^b = Resistant Parent; SP^c = Susceptible Parent.

2

3

1

Table 2 Segregation for resistance to races 7 of *Colletotrichum lindemuthianum* and 63-39 of *Pseudocercospora griseola* in 63 F_{2:3} plants from the AND 277 × Ouro Negro cross

Parental cross	Generation n	Observed Ratio (RR:RS:SS) ^a	Expected Ratio (1RR:2RS:1SS)	χ^2	<i>P</i> value (2 df)
Race 7 of <i>Colletotrichum lindemuthianum</i> – the ANT pathogen of common bean					
Ouro Negro	RP ^b	20:0:0			
AND 277	SP	0:0:20			
AND 277 × Ouro Negro	F ₁	20:0			
AND 277 × Ouro Negro	F _{2:3}	14:34:15	15.75:31.5: 15.75	0.43	0.81
Race 63-39 of <i>Pseudocercospora griseola</i> – the ALS pathogen of common bean					
Ouro Negro	RP	20:0:0			
AND 277	SP	0:0:20			
AND 277 × Ouro Negro	F ₁	20:0			
AND 277 × Ouro Negro	F _{2:3}	14:34:15	15.75: 31.5: 15.75	0.43	0.81

^a F_{2:3} families were classified as having all individuals resistant (RR), individuals resistant and susceptible (RS), and all individuals susceptible (SS);

^b RP = Resistant parent; SP = Susceptible parent.

2

Table 3 Reaction of F₂ plants from the Rudá × Ouro Negro common bean cultivar cross inoculated with races 73 of *Colletotrichum lindemuthianum* and 63-39 of *Pseudocercospora griseola* and the presence ^f(+) or absence (-) of the SF10 molecular marker

Generation	Observed numbers of plants resistant to races 73 and 63-39 ^a						Expected ratio 3:0:0:1 ^d	χ^2	P value (3 df)	Linkage distance (cM) ^e <i>Co-10/Phg-ON</i>
	R ^{ANT b} R ^{ALSc}	R ^{ANT b}	S ^{ALSc}	S ^{ANT b}	R ^{ALSc}	S ^{ANT b}				
F ₂	83	1		0		28	84:0:0:28	0.023	0.92	0.9
SF10	92 (+)	0		0		20 (-)	84:0:0:28	3.047	0.09	7.8

^a R = Resistant plants; S = Susceptible plants;

^b ANT = Anthracnose;

^c ALS = Angular leaf spot;

^d Two dominant genes in absolute linkage;

^e Linkage distance between the *Co-10/Phg-ON* genes indicates one recombinant was observed for the reaction to both pathogens;

^f Marker present (+); absent (-).

Table 4 Reaction of F_{2:3} plants from the AND 277 × Ouro Negro cross of common bean cultivars inoculated with races 7 of *Colletotrichum lindemuthianum* and 63-39 of *Pseudocercospora griseola* and the presence (+) or absence (-)^d of the g2303 molecular marker

Generation	Observed numbers of plants				Expected ratio ^b	χ^2	P value (3 df)	Linkage distance (cM) ^c
	Observed ratio to race 7		Observed ratio to race 63-39					
	1RR:RS:SS ^a		RR:RS:SS ^a					
F _{2:3}	14:34:15		14:34:15		15.75:31.5: 5.75	0.43	0.81	
(+) Presence and (-) absence of marker in F ₂ plants	48 (+)	0	0	15 (-)	47.25: 0: 0:15.75	0.05	0.83	0.0

^a R = Resistant plants; S = Susceptible plants;

^b Two dominant genes in linkage;

^c Linkage distance between the g2303 marker and the *Co-10/Phg-ON* genes indicates no recombinants were observed.

^d Marker present (+); absent (-).

Figure legends

Fig. 1 Electrophoretic analysis of the amplification products from the g2303 marker. Lanes: L, 100 bp ladder; 1, Corinthiano; 2, Ouro Negro; 3, BAT93; 4, Mexico 222; 5, Rudá. The arrow indicates the DNA band of 350 bp linked to the resistance cluster *Co-10/Phg-ON*

Fig. 2 Genetic distances and locations of the *Co-10* gene for resistance to common bean ANT, the *Phg-ON* gene for resistance to ALS, and the molecular markers g2303 and SF10 on linkage group Pv04 of *Phaseolus vulgaris* L. using the populations from the AND 277 × Ouro Negro and Rudá × Ouro Negro crosses. The map was drawn with MapChart (Voorrips 2002)

Figure 1

Figure 2