Lawrence Berkeley National Laboratory

Recent Work

Title

INSTABILITY OF A POSITIVE COLUMN IN A MAGNETIC FIELD

Permalink

<https://escholarship.org/uc/item/9kv4j844>

Authors

Allen, T.K.

Paulikas, George A.

Pyle, Robert V.

Publication Date

1960-09-14

UCRL

9399

UNIVERSITY OF
CALIFORNIA

Ernest O. Lawrence

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

**INSTABILITY OF A POSITIVE COLUMN IN A
MAGNETIC FIELD**

T. K. Allen, George A. Paulikas, and Robert V. Pyle

September 14, 1960

INSTABILITY OF A POSITIVE COLUMN IN A MAGNETIC FIELD*

T. K. Allen,[†] George A. Paulikas,[‡] and Robert V. Pyle

Lawrence Radiation Laboratory
University of California
Berkeley, California

September 14, 1960

The particle and energy losses from a long positive column were shown by Lehnert¹ to decrease with increasing magnetic field, as predicted, but only for fields less than a critical field, B_c . Above B_c the losses increased, leading to speculation about the origin of the "enhanced diffusion". Further measurements by Lehnert and Hoh,² and at this laboratory, extended Lehnert's measurements to other gases and discharge tube parameters; in particular, it was shown³ that at B_c the discharges lose their azimuthal symmetry and appear as constricted, rotating, luminous helices (Fig. 1).

Recently, Hoh⁴ and Kadomtsev and Nedospasov⁵ have considered the origin of the instability from two different standpoints. Hoh made use of a criterion for the stability of a wall sheath and showed that it is no longer satisfied when the magnetic field exceeds a critical value. The manner in which the failure of the sheath-stability criterion affects the discharge was not discussed. Kadomtsev and Nedospasov, on the other hand, show that perturbations of the discharge column can grow when diffusion to the walls can no longer overcome the effects of $\mathbf{j} \times \mathbf{B}$ forces that tend to increase the perturbations. $\mathbf{j}$ can not be zero, but the value of B_c obtained does not

---

*Work was performed under the auspices of the U. S. Atomic Energy Commission.

[†]AERE, Harwell

[‡]National Science Foundation Fellow

depend on the magnitude of the current density. In addition to predicting B_c , their theory predicts the frequency, wavelength, and growth rate of the oscillation.

In our experiment, discharge tubes 250 cm long were placed at the axis of a solenoid which could be adjusted in length up to 200 cm. As in Refs. 1 and 2, the discharges were run at constant current, and the axial field was used as a measure of particle losses. These fields were obtained from a number of probes. The time and space variations of the visible light from the discharge were photographically recorded from a rotating mirror and a series of well-collimated photomultipliers. From the measurements it was possible to obtain data on the critical magnetic fields and the oscillation frequencies, wavelengths, and growth rates for H_2 , D_2 , He, Ne, and Ar in tubes with radii of 0.9, 0.95, 1.27, and 2.75 cm.

Luminous helices of the clarity shown in Fig. 1 were observed only for special combinations of gas, pressure, and tube radius. In general, the appearance was more chaotic, often appearing as moving striations superimposed on the spirals. The pitches of the spirals were such that $\underline{j} \times \underline{B}$ was directed toward the tube walls.

Some of the experimental B_c vs p data are shown in Figs. 2 and 3, together with calculations from the formulae⁶ of Refs. 4 and 5. Numerical data were taken from the compilation by Brown.⁷ Considerable uncertainty exists as to the actual conditions in a positive column, e. g. both atomic and molecular ions are known to be present. The predictions of the sheath theory are not particularly sensitive to temperature, but depend critically on the mobilities; we show calculated curves for two ionic species to illustrate the latter point. On the other hand, the results of the current-channel instability theory depend much more sensitively on temperature than on mobility, B_c varying roughly as T_e for a given pressure. The moving

striations that were generally present in our discharges may appreciably affect the electron energy distributions, although the B_c vs p curves go smoothly through transitions from unstriated to striated regimes. Similar uncertainties exist for the H_2 , D_2 , and A discharges.

Where it was possible to obtain oscillation frequencies and wavelengths from the experimental results, these quantities were in good agreement with values obtained from the formulae in Ref. 5. For example, helium at 0.3 mm Hg pressure in a 2.75 cm radius tube gave an angular frequency of 5×10^4 radians/sec and a wavelength of 73 cm, the calculated values being 4.3×10^4 and 63, respectively.⁸ In addition, the frequencies increased with decreasing radius and decreasing pressure, as predicted in Eq. (22) of Ref. 5. Growth rates were difficult to determine experimentally, but showed agreement in order of magnitude.

We conclude that with the proper choice of parameters either theory can approximate the experimental B_c vs p curves, although the shapes of the latter are perhaps most like the results of Kadomtsev and Nedospasov. Their theory also yields perturbation frequencies and wavelengths in good agreement with experiment.

ACKNOWLEDGMENTS

We wish to thank C. M. Van Atta for supporting this research,

J. Warren Stearns for helping with many of the measurements, Y. T. Fung,

L. S. Hall, and W. B. Kunkel for discussions of the theory and experiment,

H. S. Powell for the glass blowing, and many members of William R. Baker's

Sherwood research group for aid in construction of the apparatus.

References

1. B. Lehnert, Proceedings of the Second International Conference on the Peaceful Uses of Atomic Energy (United Nations, Geneva 1958), Vol. 32, p. 349.
2. B. Lehnert and F. C. Hoh, Phys. of Fluids 3, 600 (1960).
3. T. K. Allen, G. A. Paulikas, and R. V. Pyle, Bull. Am. Phys. Soc. II, 5, No. 2, 314 (1960).
4. F. C. Hoh, Phys. Rev. Letters 4, 559 (1960).
5. B. B. Kadomtsev and A. V. Nedospasov, J. Nuclear Energy, Part C, 1960 (to be published). We are grateful to the authors for a pre-publication copy of this paper.
6. Equation (4) of Ref. 4 has apparently been misprinted; the second quantity in brackets should read $(g' p_0 \lambda / R - p_0^2)^{1/2}$.
7. S. C. Brown, Basic Data of Plasma Physics, (Wiley and Sons, Inc., New York, 1959).
8. The experimental frequencies and wavelengths were obtained from the fully developed, steady state helix. It is not obvious that they should be the same as the quantities calculated from the small amplitude theory. The amplitude dependence might be examined experimentally by using pulsed magnetic fields.

Figure Captions

Fig. 1. The critical magnetic field, B_c , vs gas pressure.

Δ : $R = 1.27$ cm, $I = 400$ ma; $\triangle$: $R = 0.9$ cm, $I = 200$ ma.

(a) $R = 1.27$ cm and (b) $R = 0.9$ cm, calculated from Kadomtsev's theory. (c) $R = 1.27$ cm and (d) $R = 0.9$ cm, calculated from Hoh's theory, assuming only He_2^+ ions present. (e) $R = 1.27$ and (f) $R = 0.9$ cm, from Hoh's theory, assuming only He^+ ions present.

Fig. 2. The critical magnetic field, B_c , vs gas pressure.

Δ : $R = 1.27$ cm, $I = 400$ ma; $\triangle$: $R = 0.9$ cm, $I = 200$ ma.

(a) $R = 1.27$ cm and (b) $R = 0.9$ cm, calculated from Kadomtsev's theory. (c) $R = 1.27$ cm and (d) $R = 0.9$ cm, calculated from Hoh's theory, assuming only Ne_2^+ ions present. (e) $R = 1.27$ cm and (f) $R = 0.9$ cm, from Hoh's theory, assuming only Ne^+ ions present.

Fig. 3. 90 deg stereo streak photograph of helical constriction in helium.

$R = 2.75$ cm, $p = 0.23$ mm Hg, $B = 710$ gauss. Traces show variation of light intensity with time and radius as seen through a slit perpendicular to axis of discharge tube. Note that this is not a photograph of the helix; the helicity has been established by wavelength measurements with photomultipliers.

Fig. 1

Fig. 2

Fig. 3