

Lawrence Berkeley National Laboratory

Recent Work

Title

UNIFIED GAUGE THEORIES OF HADRONS AND LEPTONS

Permalink

<https://escholarship.org/uc/item/9m69f093>

Authors

Bars, I.
Halpern, M.B.
Yoshimura, M.

Publication Date

1972-10-01

Submitted to
Physical Review D

RECEIVED
LAWRENCE
RADIATION LABORATORY

LBL-1339
Preprint 61

DEC 18 1972

LIBRARY AND
DOCUMENTS SECTION

UNIFIED GAUGE THEORIES OF HADRONS AND LEPTONS

I. Bars, M. B. Halpern, and M. Yoshimura

October 13, 1972

For Reference

Not to be taken from this room



Prepared for the U.S. Atomic Energy
Commission under Contract W-7405-ENG-48

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

moves further toward a unified gauge theory of particle forces. Such a unified model of hadrons and leptons has now been briefly presented in the literature.⁸ It is the purpose of this paper to discuss the model in some depth.

For strong interactions a Lagrangian is not very useful from the practical point of view: Such a Lagrangian can at best be used to describe low energy hadronic data, but we feel it will be extremely illuminating as a guide to a better understanding of hadron dynamics, and to the interplay of strong, weak, and electromagnetic interactions. For example, strictly from the hadronic viewpoint, such a model suggests that it will be useful to consider a hadron dynamics in which the strong vector mesons, at some intermediate stage in the calculation, have zero mass. (We remind the reader that this is indeed exactly what is happening in dual models at the moment.) It has been shown that there is an intimate connection between dual models and gauge theories.⁹ Now, the search is beginning for a dual Higgs-Kibble mechanism to raise the rho mass. We believe there is an intimate connection between our hadron model and the future spontaneously broken dual model, with internal symmetries, and hope our efforts may serve as a guide in the duality situation.

Further, as we shall see below, the presence of the leptons does dictate in a certain way the structure of hadronic symmetry breaking. Thus, a full understanding of strong interactions seems to require the simultaneous understanding of weak and electromagnetic forces. Intimate connections between strong, weak, and electromagnetic forces, such as shown in our model, will, we believe, be of much more than passing interest in future theory and experiment.

Our goal in this paper is then a unified renormalizable gauge theory of strong, weak, and electromagnetic forces. Our approach is based on the following reasoning: It is the hadrons whose symmetries are "known"--not the leptons. This is reflected by the plethora of lepton models, but only one $SU(3) \otimes SU(3)$ hadron model.⁷ For this reason we start firmly with the $SU(3) \otimes SU(3)$ hadrons, and search through the various lepton universes for one which "fits."

In this, we are extremely encouraged by the structure of the hadron world. As shown in Ref. 7, the symmetric hadron theory necessarily begins (before spontaneous symmetry breakdown) with a local $SU(3) \otimes SU(3)$ and an extra "global" group at least as large. The final symmetry is the product group. With Bardakci and Halpern, we thus interpret the hadrons as "welcoming" a lepton theory as a local subgroup of its "extra" group. In this paper, the extra group will be called "primed" or "leptonic."

The program is orderly. We study embedding the leptons in progressively larger "primed" groups of the hadron model. Taking $SU(3)' \otimes SU(3)'$ for the "primed" group leads to trouble with strangeness-changing processes, but when we go to $SU(4)' \otimes SU(4)'$, everything falls in place beautifully.

In our search through lepton universes, we first set ourselves the following additional boundary conditions:

- (1) We require the leptons to allow a $(3, \bar{3}) + (\bar{3}, 3)$ symmetry-breaking mechanism for hadrons.
- (2) In keeping with having only an $SU(3) \otimes SU(3)$ hadron world (3 quarks), we want the lepton model to contain only the known leptons. Requirement (2) limits us to Weinberg's theory, and is relaxed later.

It is worth remarking here that although some other lepton models can "fit" our hadrons, none is as natural as Weinberg's.

In any case, of course, some extra (heavy) quarks and leptons are required to cancel anomalies. Our models lead uniquely to an anomaly-removal scheme which, for hadrons is very much in the spirit of dual models: In particular, we find removal of anomalies and proper rate for $\pi_0 \rightarrow 2\gamma$ implies the existence of a heavy pion.

The plan of this paper is as follows. Section II contains a general formulation of gauge theories. In Sec. III we reformulate Weinberg's theory in a suggestive notation, involving a new classification of the leptons. Section IV is a review of the $U(3) \otimes U(3)$ hadron theory; we include here a discussion of the hadronic currents. In Sec. V, we discuss a physical induction of the lepton world from this hadron world. Section VI contains the model itself, details, and possible alternatives. There are two appendices. Appendix A discusses the spontaneous breakdown in the somewhat involved system of scalar mesons. Appendix B mentions the alternative but unsuccessful attempt to embed the leptons in $SU(3)' \otimes SU(3)'$.

II. GENERAL GAUGE FORMALISM

In order to present our analysis in an organized way, we outline here an operational approach, independent of representations, for writing a gauge invariant Lagrangian. We will always follow just three steps in each model we consider in this paper:

Step 1: Classify the particles in the theory with an appropriate group.

Step 2: Write a gauge invariant Lagrangian with dimension less than or equal to 4.

Step 3: Break the symmetry spontaneously, guided by physical arguments.

We emphasize here that the requirements of gauge invariance and dimension $d \leq 4$ are so restrictive that the physical content of the theory is essentially determined by the classification of the particles. Therefore, Step 1 contains the most important ingredients in building a model.

Step 1: We assume we have chosen a group whose generators are denoted by F_α , $\alpha = 1 \dots n$. The operator which generates local transformations is $U(x) = \exp i\{F_\alpha \omega_\alpha(x)\}$. The transformation properties of the particles are determined by the linear representation to which they have been assigned (nonlinear representations are excluded from our analysis, because of the criterion $d \leq 4$ for renormalizability). Thus, denoting the particles as $\phi_i(x)$, we have

$$\phi_i(x) \rightarrow u \phi_i(x) u^{-1} = S_{ij}(x) \phi_j(x) \tag{2.1}$$

0.000.00390000

where $S_{ij}(x)$ defines the representation. For example,

(a) If the group is $SU(2)$ with generators \vec{T} and $\phi_i = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}$

is a doublet, then

$$\phi \rightarrow \mathcal{U} \phi \mathcal{U}^{-1} = e^{+i \frac{\vec{T} \cdot \vec{\omega}(x)}{2}} \phi, \quad \vec{T} = \text{Pauli matrices.} \quad (2.2)$$

(b) If the group is $SU(3)_L \otimes SU(3)_R$ with generators F_L^α, F_R^α and $\phi_{ij}(x)$ is a 3×3 matrix in the $(3, \bar{3})$ representation, then

$$\phi \rightarrow \mathcal{U} \phi \mathcal{U}^{-1} = e^{i \frac{\lambda}{2} \cdot \omega_L(x)} \phi e^{-i \frac{\lambda}{2} \cdot \omega_R(x)} \quad (2.3)$$

where λ_α are the usual 3×3 $SU(3)$ matrices. The infinitesimal form of the transformation equation defines the commutators of the generators with the fields. In examples (a) and (b) above we get respectively,

$$(a) \quad [T_i, \phi] = \frac{\tau_i}{2} \phi \quad (2.4)$$

$$(b) \quad [F_L^\alpha, \phi] = \frac{\lambda^\alpha}{2} \phi, \quad [F_R^\alpha, \phi] = -\phi \frac{\lambda^\alpha}{2}. \quad (2.5)$$

Step 2: The derivative $\partial_\mu \phi = -i[P_\mu, \phi]$ ($P_\mu =$ momentum op.) does not transform covariantly when ϕ is replaced by $\mathcal{U}(x) \phi \mathcal{U}^{-1}(x)$. To define a covariant derivative independent of representation, we first define a covariant momentum operator \mathcal{P}_μ by introducing as many vector gauge bosons $V_\mu^\alpha(x)$ as there are generators,

$$\mathcal{P}_\mu = P_\mu + g V_\mu \cdot F \quad (2.6)$$

(V_μ^α are considered as c-numbers with respect to F^α in the following formal manipulations.) We demand that under $\mathcal{U}(x)$, \mathcal{P}_μ transforms covariantly

$$\mathcal{P}_\mu \rightarrow \mathcal{U} \mathcal{P}_\mu \mathcal{U}^{-1} = P_\mu + \mathcal{U}(g V_\mu \cdot F + i \partial_\mu) \mathcal{U}^{-1} \quad (2.7)$$

where the rhs is found by calculating $\mathcal{U} P_\mu \mathcal{U}^{-1} = P_\mu + \mathcal{U}[P_\mu, \mathcal{U}^{-1}] = P_\mu + i \mathcal{U} \partial_\mu \mathcal{U}^{-1}$. Here we have assumed that $[P_\mu, F^\alpha] = 0$, which means that F^α are internal symmetry generators. If we allow the more general case of $[P_\mu, F^\alpha] \neq 0$, like for example, F^α being generators of Lorentz transformations, or dilatations, etc. then we have to consider general relativity in curved space. This displays the known close relationship that exists between general relativity and the Yang-Mills approach.¹⁰ Equation (2.7) induces a transformation on $V_\mu^\alpha(x)$:

$$V_\mu \cdot F \rightarrow V_\mu' \cdot F = \mathcal{U}(V_\mu \cdot F + \frac{i}{g} \partial_\mu) \mathcal{U}^{-1}. \quad (2.8)$$

With these properties we can see that the covariant derivative is

$$\nabla_\mu \phi = -i[\mathcal{P}_\mu, \phi] = \partial_\mu \phi - ig V_\mu^\alpha [F^\alpha, \phi] \quad (2.9)$$

where the commutator $[F^\alpha, \phi]$ is specified in Step 1 and depends on representation. Indeed under a simultaneous transformation of ϕ and V_μ^α we get

$$\nabla_\mu \phi \rightarrow -i[\mathcal{U} \mathcal{P}_\mu \mathcal{U}^{-1}, \mathcal{U} \phi \mathcal{U}^{-1}] = \mathcal{U}(\nabla_\mu \phi) \mathcal{U}^{-1}. \quad (2.10)$$

Thus $\nabla_\mu \phi$ transforms covariantly (like ϕ). The covariant derivatives for V_μ^α can be found from the commutator

$$[P_\mu, P_\nu] \equiv ig F_{\mu\nu}^\alpha \cdot F^\alpha. \quad (2.11)$$

Since the lhs is covariant, so is the rhs. We get

$$F_{\mu\nu}^\alpha = \partial_\mu V_\nu^\alpha - \partial_\nu V_\mu^\alpha + g f^{\alpha\beta\gamma} V_\mu^\beta V_\nu^\gamma \quad (2.12)$$

where $f^{\alpha\beta\gamma}$ are the structure constants of the group under consideration.

Using only covariant derivatives we can now write an invariant Lagrangian as if we had global invariance, as usual.¹¹ Mass terms for V_μ^α should be omitted since they are not invariant. For renormalizability we should also require that any term in the Lagrangian has dimension $d \leq 4$.

Step 3: The gauge particles acquire mass through the Higgs-Kibble mechanism.¹ The local gauge symmetry is broken spontaneously by introducing a set of scalar mesons which acquire nonvanishing vacuum expectation value. A counting argument due to Kibble¹ shows that, when one considers this scalar meson system alone, the number of massless Goldstone bosons generated by the spontaneous breakdown is equal to the difference of the number of global symmetries existing within the scalar system before and after spontaneous breakdown. In simple¹² physically reasonable models, these Goldstone bosons are completely eliminated from the Lagrangian by a gauge transformation and they become the longitudinal components of the vector gauge bosons which acquire mass. Thus, the scalar mesons must be assigned to a

representation such as to generate, through spontaneous breakdown, the same number of Goldstone bosons as the number of gauge particles that are desired to be raised in mass.

The restrictive power of the procedure is self evident. The model is essentially completed in the first step, simply by the classification of the particles. The form of the vacuum expectation value is further restricted by physical requirements such as the existence of a massless and universal photon, masses of fermions, masses of gauge mesons (if known), physical values of coupling constants etc. This procedure also produces many relations between (bare) masses and coupling constants, which are adjusted to best fit experimental data (say to zeroth order). As a result, few possible classifications of particles are capable of yielding a viable theory. If in addition we restrict ourselves to as small a group as possible which can describe all possible interactions, and fit the data as close as possible to first order, then the form of the theory that one can write is extremely limited. This will be illustrated in the following sections.

7 0 0 0 6 9 0 0 0

III. WEINBERG'S THEORY EMBEDDED IN $SU(2)' \otimes SU(2)'$

In this section we present Weinberg's theory as an example of the procedure outlined in the preceding section. We classify the leptons with $SU(2)'_L \otimes SU(2)'_R$ with generators \vec{F}'_L, \vec{F}'_R among which only \vec{F}'_L, F'_{3R} [corresponding to a subgroup $SU(2)' \otimes U(1)$] generate local transformations; the other generators correspond only to global transformations. We are denoting our generators with a prime for notational convenience which will become clear later. This formalism, as shown below, suggests that the electronic and muonic systems form a (badly) broken $SU(2)'_L \otimes SU(2)'_R$ multiplet. The classification is such that it generates exactly Weinberg's theory for leptons. Thus, it appears that as long as we consider only the leptonic system without any reference to the hadrons this classification is equivalent to Weinberg's. However, it will be shown that it suggests a natural extension to the hadrons (not implied by Weinberg) which will be crucial in the building of a unified theory of strong, weak, and electromagnetic interactions,⁸ such that the group associated with strong interactions is $SU(3) \otimes SU(3)$.

The known leptons are embedded within $(2, \bar{2})$ and $(1, 3)$ representations of $SU(2)'_L \otimes SU(2)'_R$; ($\nu_R = \nu_\mu^C$, D = doublet, S = singlet)

$$\psi_D = \begin{pmatrix} \nu_L & \mu_R^+ \\ e_L^- & -\nu_R \end{pmatrix} \quad \psi_S = \begin{pmatrix} 0 & \mu_L^+ \\ e_R^- & 0 \end{pmatrix} \quad (3.1)$$

with the following transformation properties under $\mathcal{U}(x)$

$$\mathcal{U}(x) = \exp i[\vec{F}'_L \cdot \vec{\alpha}'_L(x) + F'_{3R} \alpha'_{3R}(x)]: \mathcal{U} \psi_D \mathcal{U}^{-1} = S'_L \psi_D S'^{-1}_R,$$

$$\mathcal{U} \psi_S \mathcal{U}^{-1} = S'_R \psi_S S'^{-1}_R, \quad \text{where } S'_L(x) = \exp i \frac{\vec{\tau}}{2} \cdot \vec{\alpha}'_L(x),$$

$$S'_R(x) = \exp i \frac{\tau_3}{2} \alpha'_{3R}(x). \quad \text{The commutators of the generators with the}$$

fields are formally defined by the infinitesimal form of the

$$\text{transformation equations. Thus } [\vec{F}'_L, \psi_D] = \frac{\vec{\tau}}{2} \psi_D, \quad [F'_{3R}, \psi_D] = -\psi_D \frac{\tau_3}{2}$$

$$[F'_{3R}, \psi_S] = [\frac{\tau_3}{2}, \psi_S]. \quad \text{The electric charge is given as } Q = F'_{3L} + F'_{3R}$$

which identifies the weak hypercharge $Y_W = F'_{3R}$. It can be checked

that this charge assigns the correct charges to each field by

commuting it with ψ_D and ψ_S . Electronic and muonic-type leptons are

not mixed by any local transformation, [only F'_{3R} is included in our

local group, while F'_{1R} and F'_{2R} are excluded]. It is due to this

choice of a local group that $SU(2)'_R$ is broken and thus, electron and

muon type leptons are distinguished by their $SU(2)'_R$ quantum numbers.

We remark that the doublet $\hat{\psi}_\mu = \begin{pmatrix} \mu_R^+ \\ -\nu_R \end{pmatrix}$ is the "G-parity" conjugate

of $\psi_\mu = \begin{pmatrix} \nu_{\mu L} \\ -\mu_L^- \end{pmatrix}$, i.e., $\hat{\psi}_\mu = i\tau_2 \psi_\mu^C$. As is well known, under $SU(2)$

both ψ_μ and $\hat{\psi}_\mu$ transform in exactly the same way: That is, if

$\psi_\mu \rightarrow S'_L \psi_\mu$ (like electronic doublet) then also $\hat{\psi}_\mu \rightarrow S'_L \hat{\psi}_\mu$. Therefore,

our local group is equivalent to Weinberg's when we consider only the

leptons.

To make our classification more transparent, we remark that

this is not the only possible 2×2 matrix classification of

leptons. Another one given by Gursey and Feinberg¹³ [which can also

be fitted to give Weinberg's theory of leptons] is very close to

Weinberg's formulation

where $\tan \phi = g'/g$. Thus, we can read off Weinberg's photon and electric charge as the coefficient of the charge operator;

$$A_\mu = \sin \phi W_{3\mu} + \cos \phi B_\mu$$

$$e = g \sin \phi = \frac{gg'}{(g^2 + g'^2)^{\frac{1}{2}}}$$
(3.9)

We emphasize that we have found the photon before spontaneous breakdown. This is because we knew a priori the form of the charge operator by making sure it assigned the correct physical charges to the particles in the theory. In fact, of course, it is quite general that specification of Q defines the photon and charges independent of spontaneous breakdown. In our later analysis we found it very convenient to follow this procedure, because it could show a priori whether a certain classification of particles involving both leptons and hadrons could give a massless, universal photon or not.

The spontaneous breakdown should be arranged such that the photon remains massless, while W_1, W_2 and $Z = \cos \phi W_3 - \sin \phi B$ become massive. For a massless photon we must demand

$$[Q, \phi]_{\phi = \langle \phi \rangle} = 0$$
(3.10)

Thus, by taking $\langle \phi \rangle = \lambda \frac{\tau_0}{2}$, i.e. $\langle \phi_0 \rangle = \lambda$ we can give masses to the gauge bosons as well as the electron and muon. It is more convenient to use the covariant derivatives obtained with \mathcal{P}_μ of eq. (3.8). With Weinberg we obtain

$$m_W = \frac{1}{2} g \lambda, \quad m_Z = \frac{1}{2} (g^2 + g'^2)^{\frac{1}{2}} \lambda, \quad G = \frac{2}{\lambda} \begin{pmatrix} m_e \\ m_\mu \end{pmatrix}$$

$$\lambda^2 = \frac{1}{\sqrt{2} G_W}$$
(3.4)

With the single proviso below, the structure of our classification is just that of Weinberg.

The form of G emphasizes the point suggested earlier that the source for the difference between the electron and the muon might be the breaking of $SU(2)'_R$. Furthermore, there is one amusing philosophical consequence of our representation. In Weinberg's original classification, the universality of electromagnetic charge is fixed by hand. However, by imagining that the leptons belong to a badly broken $SU(2)'_L \otimes SU(2)'_R$, universality of the electric charge is automatically obtained due to the construction of Q as a generator belonging to a nonabelian group.

The present $SU(2)'_L \otimes SU(2)'_R$ classification of the leptons has been our starting point for a search through schemes to unify strong, weak, and electromagnetic interactions [such that strongly interacting particles are classified with $SU(3)'_L \otimes SU(3)'_R$]. We shall see that the introduction of the Cabibbo angle resulting in unwanted $\Delta S = 1$ neutral currents, will suggest embedding the above classification in progressively larger matrices, finally resulting in the scheme of Sec. VI.

and $F_{\mu\nu}^{L\alpha}, F_{\mu\nu}^{R\alpha}$ obtained from Eq. (2.12) in terms of $V_{\mu}^{\alpha L}$ and $V_{\mu}^{\alpha R}$.
The gauge invariant Lagrangian with dimension $d \leq 4$ is

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} \text{Tr}(F_{\mu\nu}^L F_L^{\mu\nu} + F_{\mu\nu}^R F_R^{\mu\nu}) \\ & -\frac{1}{2} \text{Tr}[(\nabla_{\mu} M_L)^{\dagger} \nabla_{\mu} M_L + (\nabla_{\mu} M_R)^{\dagger} \nabla_{\mu} M_R] + m_0^2 \text{Tr}(M_L^{\dagger} M_L + M_R^{\dagger} M_R) \\ & + h_1 [(\text{Tr}(M_L^{\dagger} M_L))^2 + (\text{Tr}(M_R^{\dagger} M_R))^2] + h_2 \text{Tr}[(M_L^{\dagger} M_L)^2 + (M_R^{\dagger} M_R)^2] \end{aligned} \quad (4.7)$$

where we have written $F_{\mu\nu}^L = \sum_0^8 \frac{\lambda^{\alpha}}{2} F_{\mu\nu}^{\alpha L}$ etc.

The gauge symmetry is broken spontaneously by taking

$$\langle M_L \rangle = \langle M_R \rangle = \kappa 1. \quad (4.8)$$

There are 18 massless Goldstone bosons which are identified as $M_L^{\dagger} - M_L$ and $M_R^{\dagger} - M_R$, and which are eliminated by using the 18 degrees of freedom generated by F_L^{α} and F_R^{α} . The Goldstone bosons become the longitudinal degrees of freedom of the massive vector and axial vector mesons with masses

$$m_V^2 = m_A^2 = f^2 \kappa^2. \quad (4.9)$$

The remaining scalar particles are the hermitian part of M_L and M_R and have arbitrary $SU(3) \otimes SU(3)$ invariant masses. The final Lagrangian then is obtained from (4.7) by replacing $M_L \rightarrow M_L + \kappa 1$ and $M_R \rightarrow M_R + \kappa 1$, where now M_L and M_R are hermitian matrices.

As observed by Bardakci and Halpern, this final Lagrangian is invariant under a global final group $U(3)_L \otimes U(3)_R$ generated by $(F_L + F'_L)_{\alpha}$ and $(F_R + F'_R)_{\alpha}$.

Hadron Currents

Here we also discuss the structure of the currents associated with the final (product) group. As in massive Yang-Mills theories (in general), we distinguish two kinds of currents, both conserved but with equal charges. The first is the usual Noether current(s) $[J_N^{\mu}]$ generated by the transformation

$$V_{\mu} \rightarrow S V_{\mu} S^{-1}, \quad M + \kappa \rightarrow S(M + \kappa)S^{-1}, \quad (4.10)$$

(left and right); the second kind $[J^{\mu}]$ is associated with the transformation

$$V_{\mu} \rightarrow S(V_{\mu} + \frac{i}{f} \partial_{\mu})S^{-1}, \quad (M + \kappa) \rightarrow S(M + \kappa)S^{-1} \quad (4.11)$$

(left and right). In an ordinary massive Yang-Mills model, J^{μ} is proportional to the vector meson field. In our case, because masses arise spontaneously, we will obtain a modified field-current identity-- however, as is generally true, we maintain

$$(+f^{-1})\partial_{\mu} F^{\mu\nu} + J^{\nu} = J_N^{\nu}. \quad (4.12)$$

As seen below, weak interactions and electromagnetism do in fact couple to J^{ν} rather than J_N^{ν} , so we give next some more of its structure. The variation (4.11) gives $[S \approx 1 + i\alpha(x)]$

$$\delta \mathcal{L} = \text{Tr}[J^\mu(x) \partial_\mu \alpha(x)]$$

$$J_\mu = + \frac{i}{2}[M, \partial_\mu M] + f(M + \kappa) V_\mu(M + \kappa)$$

in the unitary gauge. We notice that, in the absence of the M 's we have the usual field-current identity.

The algebra of the J_μ is found in the usual manner, using (4.12). The results are almost those of field-algebra, with the exception that the usual c-number Schwinger term C in the space-time algebra becomes an operator. Where algebra of fields has $\delta_{\alpha\beta} C$,

$C = \frac{m_0^2}{f^2}$, we obtain

$$c_{\alpha\beta}(\text{operator}) = (M + \kappa)_{\alpha\beta}^2 = \frac{m_0^2}{f^2} \delta_{\alpha\beta} + (\text{operator terms})_{\alpha\beta} \quad (4.13)$$

The algebra including time-derivatives of currents is more complicated and will be discussed elsewhere.¹⁴

In the presence of additional hadrons, such as quarks, pions, etc., J^μ does not change, while, J_N^μ acquires extra terms involving the additional fields.

V. UNIFICATION OF HADRONIC AND LEPTONIC GAUGE THEORIES

A. Extension of Hadron Theory and General Considerations

As already suggested, the path we will explore for the unification of hadronic and leptonic models is the freedom of making local a certain subgroup of the "primed" group, and classifying the leptons with it. The following picture emerges: Hadrons are classified and transform only with the unprimed $U(3)_L \otimes U(3)_R$ chiral local group, while leptons are classified and transform with only the primed group. Gauge invariance does not allow any direct lepton-hadron interactions before spontaneous breakdown. The only fields that transform with both groups are M_L and M_R and they couple to both strong and weak gauge bosons. Thus, M_L and M_R play the role of a "bridge" between hadrons and leptons. Before spontaneous breakdown all semi-leptonic and nonleptonic weak interactions occur through intermediate M_L and M_R loops. After spontaneous breakdown however, we generate direct mixings between strong and weak gauge bosons, so that at low energies semi-leptonic and nonleptonic weak interactions occur through vector meson dominance.

For hadrons, we consider $U(3)_L \otimes U(3)_R$ chiral theory¹⁵ which includes usual quarks, $(3, \bar{3})$ scalar and pseudoscalar mesons ($\Sigma = \sigma + i\pi$), as well as vector and axial-vector mesons. These fields transform only with the unprimed generators F_L^α, F_R^α of Section IV:

$$q_L \rightarrow S_L q_L; \quad q_R \rightarrow S_R q_R; \quad \Sigma \rightarrow S_L \Sigma S_R^{-1} \quad (5.1)$$

$$V_L^\mu \rightarrow S_L (V_L^\mu + i\partial^\mu) S_L; \quad V_R^\mu \rightarrow S_R (V_R^\mu + i\partial^\mu) S_R^{-1} .$$

In this model, gauge invariance does not allow mass terms for V_L , V_R , and q . Masses for these fields can only be generated through spontaneous breakdown. For quarks we need a term in the Lagrangian of the type $(\bar{q}_L \Sigma q_R + \text{h.c.})$ (this is another reason for introducing Σ), and to generate masses for all vector and axial-vector mesons, we have to introduce the Bardakci-Halpern scalars,⁷ M_L and M_R of Sec. IV. Notice that we cannot break $SU(3)_L \otimes SU(3)_R$ in the usual way, by adding a linear term in Σ , like $\text{Tr}(f \Sigma)$, (f is a numerical matrix). This would spoil the gauge invariance (and hence renormalizability). Such a linear term must be induced only through spontaneous breakdown from a gauge invariant term. If only a hadron theory is desired, such a term is easy to find: $\text{Tr}(G M_L^+ \Sigma M_R) + \text{h.c.}$ [G is a numerical "insertion" of form $(^a a_b)$]. In the presence of leptons the term is a bit harder, but we shall find later just such a term which, in the limit of no weak or electromagnetic interactions reduces to just the above hadronic term.

Again, if only a hadron theory is desired, M_L and M_R may be taken 3×3 . In the presence of leptons however, we will need to take them as 3×4 matrices (to eliminate neutral strangeness changing currents). In general of course, we can enlarge to $3 \times n$, $n \geq 3$, thus enlarging the "primed" group to $U(n)' \otimes U(n)'$. No Goldstone bosons will couple as long as the "new" columns do not develop any vacuum expectation values (the extra global symmetries associated with the extra columns should be broken by hand). Thus

$$\langle M_L \rangle = \langle M_R \rangle = \kappa = \begin{pmatrix} \kappa_1 & 0 & 0 & 0 & 0 \\ 0 & \kappa_1 & 0 & 0 \dots 0 \\ 0 & 0 & \kappa_2 & 0 & 0 \end{pmatrix}. \quad (5.2)$$

We find that the smallest n we need is $n = 4$.

This, we find below, eliminates all neutral strangeness changing processes to first order.

B. "Induction" of Leptonic Structure from Hadrons

Among the constraints on the unification, two are particularly worth focussing on. These are (a) proper introduction of Cabibbo angle and (b) having a universal, massless photon. These play particularly crucial roles in the choice of a local group to classify the leptons, as a subgroup of the "primed" group (as well as its representations).

We first consider the photon, whose structure is closely related to the construction of the charge operator Q . In the type of theory we want to propose Q will be constructed from the unprimed as well as the primed generators. The unprimed part, which will assign the correct charges to quarks, Σ , V_μ , and A_μ , is the usual $SU(3)_L \otimes SU(3)_R$ charge operator, namely $(F_{3L} + F_{3R}) + \frac{1}{\sqrt{3}}(F_{8L} + F_{8R})$. The primed part of the charge operator should first of all give the correct charges for the leptons. We assume that we have chosen $SU(2)'_L \otimes Y'$ as a subgroup of a $U(n)'_L \otimes U(n)'_R$ group. We choose $SU(2)'_L \otimes Y'$ both because it is the natural group of the known leptons, and because, as it will turn out, our hadrons will not connect to any smaller leptonic group. Of course we will search for the smallest n compatible with data.

Thus, the total charge operator is

$$Q = F_{3L} + F_{3R} + \frac{1}{\sqrt{3}}(F_{8L} + F_{8R}) + F'_{3L} + Y'. \quad (5.3)$$

to ψ_D and ψ_S only by F'_{3R} , just as in Sec. III. Thus the charge operator that we have just chosen also assigns the correct charges to the leptons. This is why we think Weinberg's leptons are much more suggestive if classified as in Eq. (3.1) [rather than as in Eq. (3.2)].

Thus (as mentioned above) the presence of the hadrons in a sense distinguishes the electron and muon type leptons by their $SU(2)'_R$ quantum numbers, and gives a zero charge neutrino.

As already mentioned and as shown in Appendix A

$U(3)'_L \otimes U(3)'_R$ will turn out unsatisfactory. The next most economic scheme is an embedding of $SU(2)'_L \otimes Y'$ in $U(4)'_L \otimes U(4)'_R$. Out of the generators $F'^{L,R}_\alpha$, $\alpha = 0, \dots, 15$ we choose $F'^L_{1,2,3}$ to form an $SU(2)'_L$, plus a $U(1)'$ operator Y' . For reasons that will become clear shortly, the representation we want to use for these generators has the form (left or right)

$$t_\alpha = \begin{pmatrix} \frac{1}{2} \tau_\alpha & | & 0 \\ \hline 0 & | & \frac{1}{2} \hat{\tau}_\alpha \end{pmatrix} \begin{cases} \tau_\alpha = \text{Pauli matrices} = (\tau_0, \tau_1, \tau_2, \tau_3) \\ \hat{\tau}_\alpha = \tau_2 \tau_\alpha \tau_2 = (\tau_0, -\tau_1, \tau_2, -\tau_3) \end{cases} \quad (5.10)$$

Clearly $t_{0,1,2,3}$ form a $U(2)$ algebra. Equation (5.7) is satisfied if we take $Q_w^{L,R} = t_3 + \frac{1}{3} t_0$. This suggests that the primed charge operator has the form $(F'_{3L} + F'_{3R}) + \frac{1}{3}(F'_{0L} + F'_{0R})$, determining

$$Y' = F'_{3R} + \frac{1}{3} (F'_{0L} + F'_{0R}) \quad (5.11)$$

Now, if we embed the leptons of Sec. III in a 4×4 matrix

$$\psi_D = \begin{bmatrix} \nu_L & \mu_R^\dagger & & 0 \\ e_L^- & -\nu_R & & \\ & & & ? \\ 0 & & & \end{bmatrix} \quad \psi_S = \begin{bmatrix} 0 & \mu_L^\dagger & & 0 \\ e_R^- & 0 & & \\ & & & ? \\ 0 & & & \end{bmatrix} \quad (5.12)$$

and let them transform as before (but with t_α replacing $\frac{1}{2} \tau_\alpha$ in S'_L and S'_R), we see that $F'_{0L} + F'_{0R}$ commutes with both ψ_D and ψ_S . Again only F'_{3L} assigns the value of the hypercharge Y' to the leptons as in Sec. III. The question marks (?) in (5.12) are suggestive of the presence of new heavy leptons. In fact, in order to cancel anomalies we will need new leptons which will fill the spaces marked by (?); these will also fix the fourth entry of $Q_w^{L,R}$ in Eq. (5.7) exactly as given by (5.11). Discussion of heavy leptons will be continued in Sec. VI.

So far we have seen how the determination of the charge operator has greatly restricted our choice of the $SU(2)'_L \otimes Y'$ subgroup. However, we still need to introduce the Cabibbo angle. This, and the requirement of no $\Delta S = 1$ neutral currents will finally determine the representation of the $SU(2)'_L \otimes Y'$ algebra which we need.

According to Cabibbo's theory, the weak currents which are part of an $SU(2)$ multiplet "see" a rotated picture of the hadrons. Therefore, the weak $SU(2)'_L$ group must be rotated with respect to the strong $SU(3)_L$ group in such a way that the weak gauge bosons "see" a left handed strong isospin current slightly rotated in the λ_7 direction. Weak gauge boson couplings occur in our model through the scalars M_L and M_R , which after spontaneous breakdown generate terms of the form (see later sections)

$$\text{Tr}(V_L^\mu \kappa \tilde{W}_\mu \kappa^\dagger) \quad (5.13)$$

where \tilde{W}_μ is a rotated matrix $\tilde{W}_\mu = R W_\mu R^{-1}$, $W_\mu = \sum_{\alpha=1}^3 W_\mu^\alpha t_\alpha$,

R = Cabibbo rotation. With the form of κ given in Eq.(5.2) we see that R must be chosen as

$$R = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 & 0 \\ 0 & -\sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad (5.14)$$

where θ is the Cabibbo angle.

Thus our group is now generated by the operators $\tilde{F}_{1,2,3}^{\prime L}, \tilde{Y}'$ (rotated with respect to $F_{1,2,3}^{\prime L}$ and Y'). In the simplest case of $SU(3)_L^{\prime} \otimes SU(3)_R^{\prime}$ these are represented by $\tilde{\lambda}_\alpha = R \lambda_{1,2,3,8} R^{-1}$ for the left-handed generators, and unrotated $\lambda_{3,8}$ for the right-handed generators. For the case of $SU(4)_L^{\prime} \otimes SU(4)_R^{\prime}$, we represent the left- or right-handed generators by $\tilde{t}_\alpha = R t_\alpha R^{-1}$ where t_α are as given in Eq. (5.10). The representation of Q is invariant under R in both cases. However, it is only in the latter case that the neutral operators do not (Cabibbo) rotate $\tilde{t}_3 = t_3$, $\tilde{t}_0 = t_0$. As will be seen, this is why the neutral strangeness changing currents are eliminated. For this reason, we relegate further discussion of the $SU(3)_L^{\prime} \otimes SU(3)_R^{\prime}$ to Appendix B, and continue here with the preferred $SU(4)_L^{\prime} \otimes SU(4)_R^{\prime}$ scheme.

Finally, we remark that, to maintain the couplings of the unrotated leptonic world unchanged, we need of course rotate the entire leptonic representation (so that they would not be aware of the rotated generators). Thus, we will formally introduce a fully rotated notation

$$\tilde{\psi}_D \equiv R \psi_D R^{-1}, \quad \tilde{\psi}_S \equiv R \psi_S R^{-1}, \quad (5.15)$$

$$\tilde{S}_L^{\prime} = R S_L^{\prime} R^{-1}$$

which transform as above, namely, under the "primed" group

$$\tilde{\psi}_D \rightarrow \tilde{S}_L^{\prime} \tilde{\psi}_D \tilde{S}_R^{\prime -1}, \quad M_{LR} \rightarrow M_{LR} \tilde{S}_L^{\prime -1} \quad (5.16)$$

etc. Actually, this is equivalent to saying that weak gauge bosons couple to hadrons $(M_{L,R})$ with rotated \tilde{t}_α , and with unrotated t_α to leptons. In particular, $\tilde{S}_R^{\prime} = S_R$, so some of this formalism is for notational convenience.

We have given reasons for our choice of the $SU(2)_L^{\prime} \otimes Y'$ group, and its representations. Thus much has followed from no $\Delta S = 1$ neutral currents and known hadron charges. More problems remain to be solved such as medium-strong $SU(3) \otimes SU(3)$ breaking in the presence of leptons (with no physical Goldstone bosons), cancellation of anomalies, etc. These will be discussed after we construct explicitly our model in the coming sections.

0.0.4.0.3.9.0.0.0.9.2

VI. UNIFIED MODEL

A. Construction of Lagrangian

As we saw in the last section and Appendix A, the use of $SU(3) \times SU(3)_R$ as the primed group for embedding the leptons led to trouble in general with strangeness-changing processes. Speaking generally then, we have at this juncture two possible directions: One choice, followed by most authors,⁶ is to try enlarging the hadronic group (more quarks, etc.); as explained above, we consider this unesthetic at least, and in fact such attempts do not solve the "strangeness" problems in our case anyway. Thus our choice⁸ will be, as anticipated above, to enlarge the primed group to $U(4)'_L \times U(4)'_R$.

For the sake of elegance, we will present the model in a unified (strong, weak, and electromagnetic) super-matrix notation. For example, the general local operator transformation is represented by the 14×14 super-matrix

$$\mathcal{U} = \exp i(\alpha_L \cdot F_L + \alpha_R \cdot F_R + \beta \cdot \tilde{F}'_L + \gamma Y)$$

$$S = \begin{bmatrix} S_L & 0 & 0 & 0 \\ 0 & S_R & 0 & 0 \\ 0 & 0 & \tilde{S}'_R & 0 \\ 0 & 0 & 0 & \tilde{S}'_L \end{bmatrix} \quad (6.1)$$

where, as detailed in Sec. V

$$\tilde{S}'_L = \exp i(\tilde{t} \cdot \beta + \frac{1}{3} t_0 r), \quad \tilde{S}'_R = \exp i(t_3 + \frac{1}{3} t_0) r \quad (6.2)$$

and so on. Recall that the twiddle operation is the Cabbibo rotation.

In the same notation we represent all vector mesons (strong, weak, and electromagnetic), as a similar 14×14 matrix with diagonal entries

$$V^\mu: [fv_L^\mu, fv_R^\mu, g'(t_3 + \frac{1}{3} t_0)B^\mu, g\tilde{W}^\mu + \frac{1}{3} g' t_0 B^\mu] \quad (6.3)$$

Then the unified gauge transformation on V^μ is just

$$\mathcal{U} V_\mu \mathcal{U}^{-1} = S(V_\mu + i\partial_\mu)S^{-1} \quad (6.4)$$

Similarly, for Weinberg's leptons, we introduce the $SU(2)$ doublets ψ_D and $SU(2)$ singlets ψ_S as in Eq. (5.12). To fit them into a super-matrix notation ℓ , we define rotated quantities,

$$\tilde{\psi}_D = R \psi_D R^{-1}, \quad \tilde{\psi}_S = R \psi_S R^{-1}, \quad (6.5)$$

$$\ell = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \tilde{\psi}_S & \frac{1}{\sqrt{2}} \tilde{\psi}_D^C \\ 0 & 0 & \frac{1}{\sqrt{2}} \tilde{\psi}_D & 0 \end{bmatrix} \quad (6.6)$$

(ψ_D^C : charge conjugated in Dirac space and transposed in matrix space). We then specify the transformation $\mathcal{U} \ell \mathcal{U}^{-1} = S \ell S^{-1}$.

This means that the unrotated ψ_D and ψ_S transform with the unrotated representations S'_L and S'_R :

$$\psi_D \rightarrow S'_L \psi_D S'^{-1}_R, \quad \psi_S \rightarrow S'_R \psi_S S'^{-1}_R. \quad (6.7)$$

Thus, they belong respectively to $(4, \bar{4})$ and $(1, 15)$ representations of $U(4)_L \otimes U(4)_R$. For the scalar mesons, the super-matrix notation is most symmetric,

$$M \equiv \begin{bmatrix} 0 & \Sigma & 0 & M_L \\ \Sigma^\dagger & 0 & M_R & 0 \\ 0 & M_R^\dagger & 0 & \tilde{\phi}^\dagger \\ M_L^\dagger & 0 & \tilde{\phi} & 0 \end{bmatrix}, \quad U_M U^{-1} = S M S^{-1}. \quad (6.8)$$

Here $\Sigma \equiv \sigma + i\pi$ is the usual $(3, \bar{3})$ multiplet of scalars and pseudoscalars; $M_{L,R}$ are now three-by-four complex matrices (one extra column to support the enlarged primed group), $\tilde{\phi}$ is the rotated Weinberg scalar $\tilde{\phi} = \phi_0 t_0 + i\phi \cdot \tilde{t}$. The notation emphasizes that (a) $M_{L,R}$ are the only fields in the model which transform under both the hadronic and leptonic groups and (b) Weinberg's ϕ is to the leptonic system precisely what Σ is to the hadronic. It is this symmetry which will allow us to construct a $(3, \bar{3}) + (\bar{3}, 3)$ symmetry breaking term in the model. We take leptons and quarks, for the moment at least, as discussed in the previous sections.

Covariant derivatives are formed in the usual fashion, leading to our unified gauge-invariant Lagrangian

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} \text{Tr}(F_L^{\mu\nu} F_{\mu\nu}^L + F_R^{\mu\nu} F_{\mu\nu}^R) - \frac{1}{4} F_{\mu\nu}^B F_B^{\mu\nu} - \frac{1}{4} \text{Tr}(F_{\mu\nu}^W F_W^{\mu\nu}) \\ & - i \bar{q} \not{\nabla}^\mu \gamma_\mu q - i \text{Tr}(\bar{\ell} \not{\nabla}^\mu \gamma_\mu \ell) - \frac{1}{4} \text{Tr}((\not{\nabla}^\mu M)^\dagger \nabla_\mu M) \\ & + \beta(\bar{q}_L \Sigma q_R + \text{h.c.}) + \text{Tr}(\bar{\psi}_D \phi \psi_S G + \text{h.c.}) \\ & + V(M_L) + V(M_R) + V(\Sigma) + V(\phi) \\ & + \text{Tr}(G_2 \tilde{\phi}^\dagger M_L^\dagger M_L^\dagger \tilde{\phi} + G_2 M_R^\dagger M_R^\dagger \tilde{\phi}^\dagger \tilde{\phi}) + \text{Tr}(G_1 \tilde{\phi}^\dagger M_L^\dagger \Sigma M_R + \text{h.c.}). \end{aligned} \quad (6.9)$$

Here, the $V(\dots)$'s are the usual quartic and quadratic terms¹⁵ with certain G "insertions",⁷ being 4×4 numerical matrices which, because they break only F'_R -type symmetries, do not spoil the unified gauge invariance. In particular, we write their diagonal entries

$$G : \frac{2}{\lambda}(m_e, m_\mu, ?, ?), \quad G_1 : \frac{\sqrt{2}}{2\lambda} \left(\frac{f_\pi m_\pi^2}{\kappa_1^2}, \frac{f_\pi m_\pi^2}{\kappa_1^2}, \frac{2f_K m_K^2 - f_\pi m_\pi^2}{\kappa_2^2}, d \right) \\ G_2 : (a, a, b, c). \quad (6.10)$$

The interpretation of these parameters will be clarified in the following paragraph.

We come now to the spontaneous breakdown. First, we use 21 degrees of gauge freedom (all but Q) to eliminate the 3×3 submatrices of $M_L - M_L^\dagger$ and $M_R - M_R^\dagger$, and all the components of ϕ except ϕ_0 . With an eye to the charge operator (5.3), we next assign

the charge-conserving vacuum expectation values

$$\langle \phi \rangle = \lambda t_0, \quad \langle M_L \rangle = \langle M_R \rangle = \kappa. \quad (6.11)$$

We shall return in a moment to the specific allowed form of κ , but first notice that (6.11) generates, through the last term in \mathcal{L} , a linear term in the Σ field. Thus Σ itself acquires a vacuum expectation value $\langle \Sigma \rangle \equiv v$, which is the usual $(3, \bar{3}) + (\bar{3}, 3)$ hadronic symmetry breaking in the spirit of Gell-Mann, Oakes, and Renner.¹⁶

The allowed forms for κ and v require a detailed discussion of this complicated scalar system. Such is tedious and not terribly illuminating, but can be found in Appendix A. Here we only state that to lowest order, neglecting weak effects, we can take the following isospin and strangeness conserving vacuum expectation values:¹⁷

λ arbitrary and

$$\kappa = \begin{bmatrix} \kappa_1 & 0 & 0 & 0 \\ 0 & \kappa_1 & 0 & 0 \\ 0 & 0 & \kappa_2 & 0 \end{bmatrix} \quad v = \begin{bmatrix} v_1 & 0 & 0 \\ 0 & v_1 & 0 \\ 0 & 0 & v_2 \end{bmatrix} \quad (6.12)$$

with no massless Goldstone mesons. Except for d , the interpretation of the parameters in G_1 and v is standard,¹⁵ while G_2 , d , and $V(\dots)$ can be adjusted to give arbitrarily large masses to ϕ_0 and the remaining scalars in M_L and M_R . The vacuum expectation values v_i are directly related to the pseudoscalar decay constants f_π, f_K .¹⁵

To illustrate the meaning of the κ_i , we also list (ignoring electromagnetic mixing of $\rho \phi \omega$ etc. for the moment) some (bare) vector meson masses after spontaneous breakdown¹⁸

$$\begin{aligned} m_\rho^2 &= m_\omega^2 = f^2 \kappa_1^2 \\ m_\phi^2 &= f^2 \kappa_2^2, \quad m_{K^*}^2 = \frac{1}{2} f^2 [\kappa_1^2 + \kappa_2^2 + (v_1 - v_2)^2] \\ m_{A_1}^2 &= m_{\omega_A}^2 = f^2 (\kappa_1^2 + 2v_1^2), \quad m_{\phi_A}^2 = f^2 (\kappa_2^2 + 2v_2^2) \\ m_{K_A}^2 &= \frac{1}{2} f^2 (\kappa_1^2 + \kappa_2^2 + (v_1 + v_2)^2). \end{aligned} \quad (6.13)$$

With proper choices of $\kappa_1, \kappa_2, v_1, v_2$, these formulae for the vector meson masses are well satisfied by experiment. Further, W^\pm and Z also get a small extra contribution to their masses, due to κ . Such relations should be taken together with a number of remarks (1) Electromagnetic mixing, to be discussed below, gives order e^2 corrections to $\rho \omega \phi$ masses. (2) Ignoring (1), and the presence of the remaining $M_{L,R}$ terms--which influence the known hadrons only through loops,--the hadron theory is just a familiar mass-mixing Yang-Mills σ -model. Of course, with $\kappa_1 \neq \kappa_2$, we lose the second Weinberg sum rule--so, in general we prefer $\kappa_1 = \kappa_2$, leaving ω - ϕ splitting to higher order. (3) Frankly, we do not know whether our Lagrangian will be more useful as an effective Lagrangian or as a guide to nonperturbative structure and the currents of the strong interactions. In general, we will discuss whichever view (or both) when they appear interesting.

B. Photon System and Vector-Meson Diagonalization

The structure of our theory with respect to these topics is somewhat unusual. As discussed above, the (massless, universal) photon is found as the coefficient of Q in the covariant momentum:

$$(F_\alpha = F_{\alpha L} + F_{\alpha R})$$

$$\begin{aligned}
& f(F_3 V_3 + F_8 V_8) + g F'_{3L} W_3 + g' Y' B \\
& = eQ \cdot A + \frac{f}{\cos \eta} \left[\left(\frac{\sqrt{3}}{2} F_3 + \frac{1}{2} F_8 \right) - \frac{\sqrt{3}}{2} \sin^2 \eta Q \right] V_{38} \\
& + f \left(-\frac{1}{2} F_3 + \frac{\sqrt{3}}{2} F_8 \right) V'_{38} + \frac{g}{\cos \phi} (\cos^2 \phi F'_{3L} - \sin^2 \phi Y) Z \quad (6.14)
\end{aligned}$$

where

$$e = g \sin \phi \cos \eta, \quad \tan \phi = \frac{g'}{g}, \quad \tan \eta = \frac{2g \sin \phi}{\sqrt{3} f}$$

$$A^\mu = \cos \eta (\sin \phi W_3^\mu + \cos \phi B^\mu) + \sin \eta \left(\frac{\sqrt{3}}{2} V_3^\mu + \frac{1}{2} V_8^\mu \right)$$

$$V_{38} \equiv \cos \eta \left(\frac{\sqrt{3}}{2} V_3 + \frac{1}{2} V_8 \right) - \sin \eta (\sin \phi W_3 + \cos \phi B)$$

$$V'_{38} \equiv -\frac{1}{2} V_3 + \frac{\sqrt{3}}{2} V_8. \quad (6.15)$$

with $f^2/4\pi \sim 2$ and g, g' small, we obtain approximately Weinberg's $e \sim gg'/(g^2 + g'^2)^{1/2}$.

Because of the diagonalization, our picture of electromagnetic effects is unusual. As a first indication of this, we notice that we are forced to electromagnetically mix the strong vector mesons. The $\rho \phi \omega$ mass matrix becomes

$$\text{Tr}[(\sqrt{3} Q f_1 V_{38} + f U_3 V'_{38} + f' \lambda_0 V_0)^2 \kappa^2]. \quad (6.16)$$

Here we have allowed a different universal coupling constant $f' \neq f$ for the ninth vector and axial vector mesons V_0 and A_0 ; κ stands for the 3×3 submatrix of κ [Eq. (6.12)] with $\kappa_1 \neq \kappa_2$. The other symbols are given as

$$\begin{aligned}
f_1 &= f^2 (f^2 - \frac{4}{3} e^2)^{-1/2}, & Q &= \begin{pmatrix} 2/3 & 0 & 0 \\ 0 & -1/3 & 0 \\ 0 & 0 & -1/3 \end{pmatrix}, \\
U_3 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, & \lambda_0 &= \sqrt{\frac{2}{3}} 1. \quad (6.17)
\end{aligned}$$

It turns out that f' must be close to f to obtain the usual ϕ - ω (canonical nonet) mixing angle. We also find that, aside from small electromagnetic mass corrections, the ρ - ω mixing angle can be fit to data, and is very sensitive to variations in $f'-f$ of order e^2 .

Further of course, the eigenvectors of the mass matrix which are the physical $\rho, \omega,$ and ϕ , have direct order e^2/f electromagnetic couplings to the leptons. This can be easily seen from Eq. (6.14): V_{38} is also associated with the total charge Q , like the photon. Thus electromagnetic effects will not be describable purely in terms of a $J_{EM}^\mu \cdot A_\mu$ coupling. In explicit calculation, say in electron quark (electromagnetic) scattering, we find that the hadronic vector couplings always add to the photon in just such a way as to simulate vector meson-dominated electromagnetic form factors in lowest order: e.g.,

$$\frac{e^2}{q^2} - f \left(\frac{e^2}{f} \right) \frac{1}{q^2 - m_\rho^2} = \frac{e^2}{q^2} \left(\frac{-m_\rho^2}{q^2 - m_\rho^2} \right). \quad (6.18)$$

Further, these couplings give a hadronic correction to the muonic $\frac{g-2}{2}$ of order 5×10^{-8} , agreeing with previous estimates.¹⁹

We will discuss the effect of the diagonalization on currents after specifying our prescription for the other weak vector meson couplings. These we choose not to diagonalize, leaving charged W^\pm and neutral Z terms of the form

$$\mathcal{L}' = g\text{Tr}(V_L(M_L + \kappa)\tilde{W}(M_L^\dagger + \kappa^\dagger)) + (Z \text{ terms}) \quad (6.19)$$

as they are. Thus charged lowest order currents proceed via vector exchange at low energies. Actually, of course, one can diagonalize, but this is quite lengthy, and the theory is easily interpreted without doing so.

C. Currents and Universality

Hadronic currents in our model are, of course, determined by the Lagrangian \mathcal{L} . The physical weak currents J_\pm^μ can immediately be read off as the hadronic coefficient of W_\pm^μ in \mathcal{L} (M's considered hadrons). In the limit $g, g' \rightarrow 0$ (and mass of the fourth column of M's large), these currents are just those discussed in Sec. IV. The electromagnetic current is also the hadronic coefficient of A^μ , but as stressed above, this current is not useful in the usual manner, due to the "other" electromagnetic effects from neutral strong vector mesons.

Although, as in Sec. IV, these currents can be found through hadronic considerations, it is perhaps more illuminating to consider their structure from the point of view of the "primed" transformations, and the W equations of motion. As an example, we discuss the Noether derivation of J_\pm^μ from this viewpoint. For simplicity, we consider only transformations within the unitary gauge ($(M - M^\dagger)_{33} = 0, \phi - \phi^\dagger = 0$) - in this case, those generated by $\tilde{F}_{1,2}^\dagger$. Thus ϕ , M , and the other

hadrons do not transform (otherwise we cannot maintain the gauge), while the leptons transform as usual. With respect to W , we follow Sec. IV to consider two classes of transformations (and hence two groups of currents). As in Sec. IV, the two W transformations are those with or without an extra $\tilde{S}'^{-1} \partial_\mu \tilde{S}'$. The transformation with the extra derivative term leads to the physical currents J_\pm^μ defined just above; these can be written as $J_\pm^\mu = \frac{\delta}{\delta W_\pm^\mu} (\mathcal{L}_M + \mathcal{L}_\phi)$ where $\mathcal{L}_{M, \phi}$ are the covariant kinetic energy terms for M and ϕ in the unitary gauge.

The transformations without the derivative lead to $J_{\pm N}^\mu = \frac{\delta}{\delta W_\pm^\mu} (\mathcal{L} - \mathcal{L}_M - \mathcal{L}_\phi)$, which are the ordinary Noether currents of the leptons and W fields, and not the hadron currents. As always in these algebra-of-fields-type situations (see Sec. IV), the two sets of currents are related by a total divergence, this time being $\partial_\mu F_{W_\pm}^{\mu\nu}$. Thus J_\pm^μ is related to the weak Noether current $J_{\pm N}^\mu$ through the W equations of motion; simultaneously, forms essentially J_\pm^μ are related, as in Sec. IV, to the hadronic Noether current through the strong vector meson equations of motion.

It is clear from the above discussion, that the hadronic charge algebra is that of the leptonic charges; hence universality is guaranteed.²⁰

A further remark about the electromagnetic current: It would be useful to have an "effective" electromagnetic current, that takes into account the hadronic corrections mentioned above. We would conjecture that such an object is the current coupling to Weinberg's photon (i.e., do not diagonalize; Weinberg's photon is the real photon taken at $\eta = 0$).

Neutral Strangeness-Changing Currents and a Correspondence Principle

In lowest order, we get no $\Delta S = 1$, $\Delta Q = 0$ currents, because our Cabibbo rotation (5.14) does not rotate neutral weak vector mesons [$\tilde{t}_3 = t_3$, $\tilde{t}_0 = t_0$], we have accomplished this only by increasing the size of $M_{L,R}$, without extra quarks.

On the other hand, it is clear that the four columns of $M_{L,R}$ are acting like the (p, n, λ , p') quarks of other models.⁶ In fact, we see a type of "correspondence principle" at work here in the sense that, from the structure of some n-quark "direct coupling theory" (e.g., $\bar{q} W q$), we can read an n-column "M-theory" (our models here)-- or vice versa. This principle will be useful below when we consider inclusion of other lepton models.

Preliminary calculations indicate that higher order induced strange currents are suppressed by factors of hadron masses and mass splittings divided by M_W^2 . Before vacuum expectation values, such processes are zero to order g^4 --due to a cancellation between internal p and p'-type column exchanges [as in the Glashow-Illiopoulos-Maiani SU(4) model]. Therefore, after spontaneous breakdown, these amplitudes are suppressed by hadron masses and mass differences divided by M_W^2 . These conclusions are being checked in detail and will be presented elsewhere.²¹

D. Fermions and Anomalies

As thus far presented, our model has anomalies. Further, in the presence of both strong and weak vector mesons, it does not appear possible to cancel quark versus lepton anomalies. Hence we will discuss a simple doubling scheme which, at least for the hadrons, is very much in the spirit of dual models. In particular, our approach will lead

us directly to the existence of a heavy pion. The scheme is as follows.

We introduce (heavy) q' , $\psi'_{S,D}$ that couple to gauge bosons just as q , $\psi_{S,D}$ but with the opposite sign of γ_5 . The new leptons go where we had question marks in the 4×4 lepton matrices:

$$\psi_D = \begin{bmatrix} \nu_{eL} & \mu_R^\dagger & 0 & 0 \\ e_L^- & -(\nu_\mu^C)_R & 0 & 0 \\ 0 & 0 & -\nu_{L\mu}^{\prime-} & e_R^{\prime-} \\ 0 & 0 & \mu_L^{\prime+} & (\nu_e^C)_R \end{bmatrix},$$

$$\psi_S = \begin{bmatrix} 0 & \mu_L^\dagger & 0 & 0 \\ e_R^- & 0 & 0 & 0 \\ 0 & 0 & \beta' \nu_{\mu R}^{\prime-} & e_R^{\prime-} \\ 0 & 0 & \mu_R^{\prime+} & \alpha' \nu_{eR}^{\prime-} \end{bmatrix} \quad (6.20)$$

In the leptonic system, anomalies are cancelled without complication. However now, q and q' loops with an odd number of γ_5 couplings tend to cancel (because both type of quarks are picking up masses and interactions from the same type of terms $\bar{q}_L \Sigma q_R$, $\bar{q}'_R \Sigma q'_L$)--suppressing $\pi_0 \rightarrow 2\gamma$. This we cannot allow. The only solution to this dilemma appears to require the introduction of a heavy pion'-sigma' field $\Sigma' = \sigma' + i\pi'$. For simplicity, we choose to couple q only to Σ , q' only to Σ' . (Most general couplings do not affect the conclusion.) Now it is easy to arrange that the masses of q' , Σ' are high while keeping $\frac{\langle \Sigma' \rangle}{\langle \Sigma \rangle} = \frac{v'}{v} \ll 1$ so that the new Σ' has

960016900000

negligible effect on all low lying hadrons, including V and A. Now, of course, $\pi_0 \rightarrow 2\gamma$ proceeds only through q. To get an extra factor of 3 in $\pi_0 \rightarrow 2\gamma$ amplitude there are a number of choices. We can go to sets of integrally charged quarks, with Y providing a "charm", or most perversely, e.g., introduce two more such "pairs" of cancelling quarks with large mass.²² Such schemes appear quite flexible with regard to quark classification, the only common denominator being the apparent necessity for a heavy pion. Implications of such ideas in the operatorial formulation of PCAC will be explored elsewhere.

E. Other Lepton Models

Among the other lepton models in the literature, none fits our hadrons as well as Weinberg's. However, with more scalar mesons (etc.), some other models can be incorporated, and we will make some brief remarks on this subject.

The second model of Prentki and Zumino looks good at first sight.⁸ Indeed, their leptons fit naturally into our $\psi_{D,S}$. However, due to the $(2)^{-\frac{1}{2}}$ in their neutrino classification, we would violate hadronic universality (by this factor) if we coupled directly to our hadrons above.²³ Consulting the correspondence principle again, we find that universality is restored with the addition of one more set of 3×4 matrices $M'_{L,R}$. At this stage, however, we consider this unattractive.

A more economical generalization follows lines between Weinberg and that of Georgi and Glashow. As discussed in Sec. V, the presence of the M's requires four weak gauge bosons. Keeping the same weak gauge bosons, we now classify the leptons under $SU(2)$ as Georgi and Glashow. The leptons are singlets under Y. The Georgi-

Glashow scalar fields $\phi_G = \vec{T} \cdot \vec{\phi}_G$ are needed to retain their lepton mass pattern; Weinberg's scalars ϕ_W are also needed to construct our $(\bar{3}, \bar{3})$ symmetry-breaking term (see above). Further $\langle \phi_W \rangle = \lambda$ can be taken to provide the bulk of weak gauge boson masses. To avoid Goldstone bosons in lowest order, terms like $\text{Tr}[\phi_W^\dagger \phi_G \phi_W]$ must be included in the potential. In this model then, with only three extra scalars (ϕ_G), we suppress neutrino processes in the manner of Georgi and Glashow. Without the extra U(1), the original $O(3)$ model of Georgi and Glashow does not seem possible to incorporate.

We have not found a way of incorporating (without Goldstone bosons in lowest order) the model of Lee, Prentki, and Zumino.

F. Final Remarks and Directions

We would like to discuss briefly perturbation expansion around the "hadron" theory. We choose to hold fixed the masses of W, Z, and ϕ . This leaves one parameter, say e (electric charge) to expand all weak and electromagnetic effects. As $e \rightarrow 0$, we reach the pure hadronic system, which is of interest in itself. The hadron Lagrangian is in most respects the model of Bardakci and Halpern. A notable, exception, of course, is the $(\bar{3}, \bar{3})$ symmetry-breaking term. Since $\lambda^{-1} = g M_W^{-1} \sim O(e)$, then G_1 [Eq. (6.9)] is also $O(e)$. Thus in the term $\text{Tr}(G_1(\tilde{\phi} + \lambda)M_L^\dagger \Sigma_R)$, the term $\text{Tr}(\lambda G_1 M_L^\dagger \Sigma_R)$ survives as (another e-independent) part of the hadron world. Thus all hadron symmetry breaking occurs in terms of dimension $d \leq 3$. Having required, as we have, parity, isospin and hypercharge conserving strong interactions--this conclusion about symmetry-breaking dimensionality follows directly from the structure of the leptons.

For the moment, we will not introduce the term $\det \Sigma + \det \Sigma^\dagger$, but will return later to remark on the circumstances of its inclusion. Now, we assign vacuum expectation values $\langle \phi \rangle = \lambda t_0$, $\langle M_L \rangle = \langle M_R \rangle = \kappa$, $\langle \Sigma \rangle = v$

$$\kappa = \begin{pmatrix} \kappa_1 & 0 & 0 & 0 \\ 0 & \kappa_2 & 0 & 0 \\ 0 & 0 & \kappa_3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad v = \begin{pmatrix} v_1 & 0 & 0 \\ 0 & v_2 & 0 \\ 0 & 0 & v_3 \end{pmatrix} \quad (A.3)$$

and κ_i , λ , v_i are real numbers.

Let us first deal with the particles of the fourth column of

$M_{L,R}$. Writing

$$M_{L,R} = \begin{pmatrix} \xi_{L,R} & \chi_{L,R} \\ 000 & 0 \end{pmatrix} \quad (A.4)$$

where $\xi_{L,R}$ are 3×3 matrix fields and $\chi_{L,R}$ is the fourth column, we notice that we are inducing no linear terms in χ , χ^\dagger . This is consistent with (A.3). In fact, the set of quadratic terms involving $\chi_{L,R}$ is just the usual terms from $V(M)$, plus

$$\frac{1}{2\sqrt{2}} d \chi_L^\dagger v \chi_R + \frac{\lambda^2}{4} c \chi_L^\dagger \chi_L + (L \leftrightarrow R) \quad (A.5)$$

where c, d are the numbers in the 4,4 position of $G_{1,2}$. The parameters c and d can be adjusted to give arbitrary masses to $\chi_L \pm \chi_R$. Thus, though $\chi_{L,R}$ are extremely important in the structural

connection between strong and nonstrong interactions, they play no important role in the analysis of the scalar system. In what follows, we regard $M_{L,R} \sim \xi_{L,R}$ as just 3×3 matrices.

Proceeding, we list the (matrix) relations obtained on requiring the absence of linear terms in ϕ , Σ , M :

$$\begin{aligned} \frac{1}{2} \lambda v G_1' + \frac{1}{4} \lambda^2 G_2' + \alpha + 2\beta \text{Tr}(\kappa^2) + 2\gamma \kappa^2 &= 0 \\ \frac{1}{2} \lambda \kappa^2 G_1' + [\alpha' + 2\beta' \text{Tr}(v^2) + 2\gamma' v^2] v &= 0 \end{aligned} \quad (A.6)$$

$$\text{Tr}(v \kappa^2 G_1') + [\text{Tr}(\kappa^2 G_2') + 2\delta + 16\epsilon \lambda^2] \lambda = 0$$

where $\kappa G_1'$ are the 3×3 parts of κ, G_1' . Because of the "insertions" $G_{1,2}'$, the equations are well underdetermined even allowing arbitrary diagonal κ , v , λ .

In preparation for writing down the quadratic terms, we use our 21 degrees of gauge freedom (all but Q) to eliminate the 21 scalar degrees of freedom $(\xi^\dagger - \xi)_{L,R}$ and $(\phi - \phi^\dagger)$. Then using (A.6) to simplify, we have

$$\begin{aligned} &\text{Tr}((2\gamma \kappa^2 - \frac{1}{2} \lambda v G_1') \xi_L^2 + 2\gamma \kappa \xi_L \kappa \xi_L + \frac{\lambda}{2} G_1' \xi_L v \xi_R) + 4\beta (\text{Tr} \kappa \xi_L)^2 \\ &+ \xi_L \leftrightarrow \xi_R + \text{Tr}(-\frac{1}{2} \lambda \kappa^2 v^{-1} G_1' \Sigma^\dagger \Sigma + 2\gamma' v^2 \Sigma \Sigma^\dagger + \gamma' \Sigma v \Sigma v + \gamma' \Sigma^\dagger v \Sigma^\dagger v) \\ &+ \beta' [\text{Tr} v (\Sigma + \Sigma^\dagger)]^2 + \frac{1}{4} [\epsilon \lambda^2 + \frac{2}{\lambda} \text{Tr}(v \kappa^2 G_1')] \phi_0^2 \\ &+ \text{Tr}(\lambda \kappa G_1' (\xi_L \Sigma + \Sigma \xi_R + \Sigma^\dagger \xi_L + \xi_R \Sigma^\dagger)) \\ &+ \phi_0 \text{Tr}((\xi_L + \xi_R)(v \kappa G_1' + \lambda \kappa G_2') + (\Sigma + \Sigma^\dagger) \kappa^2 G_1') \end{aligned} \quad (A.7)$$

charge combination $F'_{3R} + \frac{1}{\sqrt{3}} F'_{8R}$ is invariant under a right-handed Cabibbo rotation, we could write $\tilde{Y}_1 = \tilde{F}'_{3R} + \frac{1}{\sqrt{3}} (\tilde{F}'_{8L} + \tilde{F}'_{8R})$.

Local transformations

The general local operator

$$\mathcal{U} = \exp i(\alpha_L \cdot F_L + \alpha_R \cdot F_R + \vec{\beta} \cdot \tilde{F}'_L + \gamma_1 \tilde{Y}_1 + \gamma_2 \tilde{Y}_2) \quad (B.1)$$

is represented in a unified super matrix notation:

$$S = \begin{bmatrix} S_L & & & \\ & S_R & & \\ & & S'_R & \\ & & & \tilde{S}'_L \end{bmatrix} \quad (B.2)$$

where

$$\tilde{S}'_L(x) = \exp \frac{i}{2} [\vec{\lambda} \cdot \vec{\beta} + \frac{1}{\sqrt{3}} \tilde{\lambda}_8 (\gamma_1 + \gamma_2)] \quad (B.3)$$

$$S'_R(x) = \exp \frac{i}{2} [(\lambda_3 + \frac{1}{\sqrt{3}} \lambda_8) \gamma_1 + \frac{1}{\sqrt{3}} \lambda_8 \gamma_2] \quad \text{etc.}$$

Fields and classification

The hadronic part includes quarks, vector and axial vector mesons, $\Sigma = \sigma + i\pi$ multiplet etc. The only change from the model in the text being that M_L and M_R are now 3×3 matrices, transforming as

$$\mathcal{U} M_L \mathcal{U}^{-1} = S_L M_L \tilde{S}'_L{}^{-1}, \quad \mathcal{U} M_R \mathcal{U}^{-1} = S_R M_R S'_R{}^{-1}. \quad (B.4)$$

The weak gauge bosons are \vec{W}_μ , $B_{1\mu}$, $B_{2\mu}$ associated with the generators \vec{F}'_L , \tilde{Y}_1 , \tilde{Y}_2 with couplings g , g'_1 , g'_2 respectively.

The $SU(2)'_L$ doublet leptons ψ_D are assigned to part of the $(3, \bar{3})'$ representation while the singlet leptons ψ_S belong to $(1, 8)'$:

$$\psi_D = \begin{bmatrix} \nu_L & \mu_R^\dagger & 0 \\ e_L^- & -\nu_R & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \psi_S = \begin{bmatrix} 0 & \mu_L^\dagger & 0 \\ e_R^- & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad (B.5)$$

Defining the rotated representation for ψ_D : $\tilde{\psi}_D = e^{i\lambda_7 \theta} \psi_D$, we specify the transformations

$$\mathcal{U} \tilde{\psi}_D \mathcal{U}^{-1} = \tilde{S}'_L \tilde{\psi}_D S'_R{}^{-1} \quad \text{and} \quad \mathcal{U} \psi_S \mathcal{U}^{-1} = S'_R \psi_S S'_R{}^{-1}. \quad (B.6)$$

(This means that the unrotated ψ_D transforms with the unrotated S'_L $\psi_D \rightarrow S'_L \psi_D S'_R{}^{-1}$.) Finally we introduce a $(3, \bar{3})'$ complex 3×3 scalar ϕ ($\tilde{\phi} = e^{i\theta \lambda_7} \phi$) transforming just like ψ_D , and satisfying the invariant linear constraint

$$\text{Tr} ((\tilde{\phi} - \tilde{\phi}^\dagger)P) = 0. \quad (B.7)$$

Here $P = \begin{pmatrix} 0 & & \\ & 0 & \\ & & 1 \end{pmatrix}$ commutes with the local

$SU(2)'_L \otimes U(1)'_1 \otimes U(1)'_2$ group. This last constraint is necessary to avoid a Goldstone boson in lowest order in the spontaneous breakdown scheme we wish to consider below.

Lagrangian and spontaneous breakdown

Covariant derivatives are written with the help of the covariant momentum operators

$$P_\mu = P_\mu + f(V_{L\mu} \cdot F_L + V_{R\mu} \cdot F_R) + g \vec{F}'_L \cdot \vec{W}_\mu + g'_1 Y_1 B_{1\mu} + g'_2 Y_2 B_{2\mu} \quad (B.8)$$

The Lagrangian is constructed analogously to the $U(4)'_L \otimes U(4)'_R$ model in Sec. VI, with the G-insertions now being 3×3 matrices and in particular the G_1 insertion parametrized as

$$G_1 = \lambda^{-1} \kappa^{-2} \begin{pmatrix} f_\pi m_\pi^2 & & & \\ & f_\pi m_\pi^2 & & \\ & & f_\pi m_\pi^2 & \\ & & & 2f_K m_K^2 - f_\pi m_\pi^2 \end{pmatrix} \quad (B.9)$$

where λ and κ are respectively the vacuum expectation values

$$\langle \tilde{\phi} \rangle = \lambda = \begin{pmatrix} \lambda \\ \lambda' \\ \lambda'' \end{pmatrix}, \quad \langle M_L \rangle = \langle M_R \rangle = \kappa = \begin{pmatrix} \kappa_1 & & \\ & \kappa_1 & \\ & & \kappa_2 \end{pmatrix}$$

(We will specialize to $\lambda'' = \lambda$ below for simplicity. We do not find any interesting results with $\lambda'' \neq \lambda$.) The potential term $V(\phi)$ also must contain the necessary gauge invariant insertions that break extra symmetries by hand, thus avoiding massless Goldstone bosons in lowest order.

Further, notice, that, we can write a lepton mass generating term

$$G \text{Tr}(\bar{\psi}_D \phi \psi_S) + \text{h.c.} \quad (B.10)$$

which gives the mass ratio $\frac{m_e}{m_\mu} = \frac{\lambda'}{\lambda} \cos \theta$. (This relation can actually be broken by allowing G to be a matrix inside the trace, without breaking the gauge invariance.)

Photon diagonalization and heavy neutral weak gauge bosons

If the photon is found as in Sec. VI B, we remain with two weak neutral gauge bosons Z and B_2 associated with the following generators and couplings

$$(g^2 + g_1'^2)^{\frac{1}{2}} Z_\mu (\cos^2 \phi \tilde{F}'_{3L} - \sin^2 \phi \tilde{Y}_1) + g_2' \frac{1}{\sqrt{3}} (\tilde{F}'_{L8} + F'_{R8}) B_{2\mu} \quad (B.11)$$

where

$$Z = \cos \phi W_3 - \sin \phi B_1 \quad (B.12)$$

$$\tan \phi = g_1'/g$$

In this notation we write the mass matrix for the massive weak gauge bosons:

$$g^2 [\lambda^2 + \lambda'^2 \cos^2 \theta + \lambda''^2 \sin^2 \theta] (W_1^2 + W_2^2) + (g^2 + g_1'^2) [\lambda^2 + \lambda'^2 \cos^2 \theta + \lambda''^2 \sin^2 \theta] Z^2 + g_2'^2 (\lambda'^2 + \lambda''^2) \sin^2 \theta B_2^2 - 2 g_2' (g^2 + g_1'^2)^{\frac{1}{2}} \lambda''^2 \sin^2 \theta Z B_2 \quad (B.13)$$

Defining the neutral eigenstates Z_1, Z_2 as

$$Z = \cos \alpha Z_1 + \sin \alpha Z_2 \quad (\text{B.14})$$

$$B_2 = -\sin \alpha Z_1 + \cos \alpha Z_2 .$$

We find in the approximation $\lambda' \ll \lambda \approx \lambda''$ (remember $\frac{m_e}{m_\mu} \approx \frac{\lambda'}{\lambda}$)

$$\tan 2\alpha \approx \frac{2x - \sin^2 \theta}{x^2 - \sin^2 \theta}, \quad x = \frac{g^2 + g_1'^2}{g_2^2} . \quad (\text{B.15})$$

Strangeness changing neutral currents

The couplings of the heavy neutral weak gauge bosons to hadronic and leptonic currents are obtained as

$$\begin{aligned} \mathcal{L} \sim & \sin \theta \cos \theta [J_{L6}^h + J_{\Delta S=\Delta Q=0}^h]_\mu [(g^2 + g_1'^2)^{\frac{1}{2}} Z_\mu - g_2' B_{2\mu}] \\ & + (g^2 + g_1'^2)^{\frac{1}{2}} Z_\mu (\cos^2 \phi j_{3L}^l - \sin^2 \phi j_{3R}^l)_\mu \end{aligned} \quad (\text{B.16})$$

where, J_{L6}^h is the left-handed hadronic current associated with λ_6 , and j_{3L}^l is the left-handed leptonic current associated with λ_3 etc.

Rewriting the above in terms of $Z_{1,2}$, and using the mass matrix, we calculate the effective semi-leptonic, nonleptonic, and leptonic Lagrangians. We find that:

(1) The purely leptonic processes are almost as in Weinberg's Lagrangian with a small change of the order of $\frac{\lambda'^2}{\lambda^2} \approx \left(\frac{m_e}{m_\mu}\right)^2$.

(2) The neutral $\Delta S = 1$ currents in semi-leptonic processes are suppressed in the decay rates by a factor of $\left(\frac{\lambda'}{\lambda}\right)^4 \approx \left(\frac{m_e}{m_\mu}\right)^4$.

(3) The nonleptonic effective weak Lagrangian contains $\Delta S = 2$ pieces, which, compared to the largest $\Delta S = 0$, pieces are smaller only by a factor of $\approx (\sin \theta)^2$. There are also terms which may give an approximate $\Delta I = 1/2$ rule ($\lambda'' \neq \lambda$ may be better).

Conclusion

In spite of a lot of effort we could not improve on item (3) above within many variations of the $SU(3)_L' \otimes SU(3)_R'$ scheme. To avoid this large contribution of $\Delta S = 2$ processes in lowest order, we were finally led to consider enlarging the primed group to $U(4)' \otimes U(4)'$ as discussed in Sec. VI.

FOOTNOTES AND REFERENCES

- * Supported in part by the U. S. Atomic Energy Commission.
- † Participating guest, Lawrence Berkeley Laboratory.
- § Present address: Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania 19104.
1. P. W. Higgs, Phys. Letters 12, 132 (1964); Phys. Rev. Letters 13, 508 (1964); Phys. Rev. 145, 1156 (1966); T. W. B. Kibble, Phys. Rev. 155, 1554 (1967).
 2. S. Weinberg, Phys. Rev. Letters 19, 1264 (1967).
 3. A. Salam, Elementary Particle Physics, edited by N. Svartholm (Almquist and Wiksells, Stockholm, 1968) p. 367.
 4. G. 't Hooft, Phys. Letters 39B, 195 (1971); Nucl. Phys. B35, 167 (1971); B. W. Lee, Phys. Rev. D5, 823 (1972); B. W. Lee and J. Zinn-Justin, Phys. Rev. D5, 3121 (1972); *ibid.* 5, 3137 (1972); *ibid.* 5, 3155 (1972).
 5. M. Veltman, unpublished; D. Gross and R. Jackiw, Phys. Rev. D6, 477 (1972); C. Bouchiat, J. Iliopolis, and Ph. Meyer (unpublished); H. Georgi and S. Glashow, Phys. Rev. D6, 429 (1972).
 6. S. Weinberg, Phys. Rev. D5, 1962 (1972); Phys. Rev. D5, 1412 (1972); H. Georgi and S. Glashow, Phys. Rev. Letters 28, 1494 (1972); B. W. Lee, Phys. Rev. D6, 1188 (1972); J. Prentki and B. Zumino (to be published).
 7. K. Bardakci and M. B. Halpern, Phys. Rev. D6, 696 (1972).
 8. I. Bars, M. B. Halpern, and M. Yoshimura, Phys. Rev. Letters 29, 969 (1972).
 9. A. Neveu and J. Scherk (unpublished).

10. R. Utiyama, Phys. Rev. 101, 1597 (1956); T. W. B. Kibble, J. of Math. Phys. 2, 212 (1961).
11. C. N. Yang and R. L. Mills, Phys. Rev. 96, 191 (1954).
12. See, however, S. Coleman and E. Weinberg (unpublished).
13. F. Gürsey and G. Feinberg, Phys. Rev. 128, 378 (1962), Appendix II.
14. Following the limit procedure of K. Bardakci, Y. Frishman, and M. B. Halpern, Phys. Rev. 170, 1353 (1968), we can map the gauge theory of hadrons onto a Sugawara-like theory. The limit is $f \rightarrow 0$, $m^2 \rightarrow 0$ holding $\kappa^2 = \frac{m^2}{f^2}$ constant. The theory seems to resemble a recent model of R. Perrin (unpublished).
15. See, e.g., S. Gasiorowicz and D. A. Geffen, Rev. Mod. Phys. 41, 531 (1969), and references therein.
16. M. Gell-Mann, R. J. Oakes, and B. Renner, Phys. Rev. 175, 2195 (1968); S. L. Glashow and S. Weinberg, Phys. Rev. Letters 20, 224 (1968).
17. Actually, there is freedom to induce $\langle \sigma_3 \rangle$ as well, if desired, for a lowest order model of $\eta \rightarrow 3\pi$. See also Appendix A.
18. As mentioned in Ref. 7 and footnote 3 of Ref. 8, it is elementary to generalize to a ninth coupling $f' \neq f$. See also in this connection H. Georgi (unpublished).
19. M. Gourdin and E. de Rafael, Nucl. Phys. B10, 667 (1969). Our calculation parallels the vector contribution of the Z-exchange diagram in the weak contribution to $g-2$ (with Z-parameters replaced by ρ -parameters). See I. Bars and M. Yoshimura, Phys. Rev. D6, 374 (1972); R. Jackiw and S. Weinberg, Phys. Rev. D5, 2396 (1972).

20. M. Gell-Mann, Phys. 1, 63 (1964).
21. H. Aytac and I. Bars (in preparation).
22. This mechanism is now called "red, white, and blue quarks" by Gell-Mann and collaborators (unpublished). The mechanism was proposed independently, and at about the same time, by us in Ref. 8.
23. We thank B. W. Lee for making this point to us.
24. We thank S. Coleman for making this point to us.
25. S. Coleman and S. Glashow, Ann. of Phys. 17, 41 (1962).
26. Particle Data Group, Phys. Letters, April 1972.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720