

UC Berkeley

UC Berkeley Previously Published Works

Title

Intra-species genetic variability drives carbon metabolism and symbiotic host interactions in the ectomycorrhizal fungus *Pisolithus microcarpus*

Permalink

<https://escholarship.org/uc/item/9nd8s0dz>

Journal

Environmental Microbiology, 23(4)

ISSN

1462-2912

Authors

Plett, Krista L
Kohler, Annegret
Lebel, Teresa
[et al.](#)

Publication Date

2021-04-01

DOI

10.1111/1462-2920.15320

Peer reviewed

Intra-species genetic variability drives carbon use strategies and symbiotic host interactions in the ectomycorrhizal fungus *Pisolithus microcarpus*.

Krista L. Plett^{1*}, Annegret Kohler², Teresa Lebel³, Vasanth R. Singan⁴, Diane Bauer⁴, Guifen He⁴, Vivian Ng⁴, Igor V. Grigoriev⁴, Francis Martin², Jonathan M. Plett¹, Ian C. Anderson¹

¹ Hawkesbury Institute for the Environment, Western Sydney University, Richmond, NSW 2753, Australia

² INRA, Interactions Arbres/Microorganismes, Laboratory of Excellence ARBRE, INRA-Nancy, 54280 Champenoux, France

³ Royal Botanic Gardens Victoria, Melbourne, VIC, 3004, Australia

⁴ US Department of Energy Joint Genome Institute, Walnut Creek, CA 94598, USA

*Corresponding author:

Krista L. Plett

Hawkesbury Institute for the Environment

Western Sydney University

Locked Bag 1797

Penrith, NSW, 2751, Australia

k.plett@westernsydney.edu.au

(+61) 2 4570 1883

Running title: Intra-species genetic variability in *P. microcarpus*

24 **Originality-Significance Statement**

25

26

27 **Summary (200 words)**

28 **Introduction**

29

30 Ectomycorrhizal (ECM) fungi form beneficial symbiotic associations with the roots of
31 approximately 3% of land plant species (Smith and Read, 2008), however, as their hosts
32 include the majority of tree species dominating boreal and temperate forests, the influence of
33 ECM fungi on ecosystem function and nutrient cycling is substantial (Read and Perez-
34 Moreno, 2003; Zak *et al.*, 2019). Mycelium of ECM fungi form a mycorrhizal root tip by
35 growing around the fine roots of the host plant and then penetrating in-between root
36 epidermal cells to create the complex symbiotic structure known as the Hartig net, which acts
37 as the site of nutrient exchange between the two organisms. These fungi acquire the majority
38 of their carbon resources from their host, with as much as one third of the carbon fixed by the
39 tree transferred belowground to mycorrhizal symbionts in exchange for host growth-limiting
40 nutrients (Nehls *et al.*, 2010). Additionally, ECM fungi are increasingly seen as forest
41 decomposers, breaking down soil organic matter to acquire nutrients (Lindahl and Tunlid,
42 2015; Shah *et al.*, 2016), although this decomposing activity typically does not result in
43 sufficient amounts of carbon for extensive growth or the formation of reproductive structures,
44 leaving the fungus reliant on its host (Lindahl *et al.*, 2007; Le Tacon *et al.*, 2015). The impact
45 of ECM fungi on host tree nutrition, coupled with their role in soil organic matter
46 decomposition, highlights the need to properly integrate ECM fungi into global ecosystem
47 models (Orwin *et al.*, 2011; Sulman *et al.*, 2017; Fry *et al.*, 2019). One factor complicating
48 this goal, however, stems from the complex evolutionary history of ECM fungi. Rather than
49 descending from one common ancestor, ECM fungi are highly diverse, having evolved
50 independently 60-80 times from saprotrophic ancestors (Hibbett *et al.*, 2000; Tedersoo and
51 Smith, 2013; Martin *et al.*, 2016). Thus, while all ECM fungi form the same symbiotic

structures and exchange nutrients with their hosts, they exhibit a high level of variability in areas such as host range and nutritional benefit, growth characteristics and the ability to decay soil organic matter (Ekblad *et al.*, 2013; Clemmensen *et al.*, 2015; Pellitier and Zak, 2018).

While the importance of considering ECM function in light of fungal lineage is becoming increasingly apparent (Pellitier and Zak, 2018; Stuart and Plett, 2020), the role of intra-species genetic variation and the extent to which it may impact ECM fungal growth and function has received less focus. Previous studies have demonstrated that intra-species variation can have significant impacts on the ability of ECM fungi to scavenge N from organic sources (Anderson *et al.*, 1999; Sawyer *et al.*, 2003; Guidot *et al.*, 2005) or colonize and interact with their hosts (Plett *et al.*, 2015; Hazard *et al.*, 2017; Hortal *et al.*, 2017). Intra-species diversity in natural ecosystems may, in fact, be equally as important as inter-species diversity in ecosystem functionality and outcomes (Johnson *et al.*, 2012, Hazard *et al.*, 2017). Given these increasingly apparent effects of intra-species variation on ECM fungi and their interactions with their hosts, more research is required to better understand what is “core” to the ECM-host relationship and what is individual specific.

We chose to use the model interaction between the ECM fungus *Pisolithus microcarpus* and its host *Eucalyptus grandis* to better understand the impact of intra-specific variability on fungal traits within altered abiotic conditions. Along with the ability of *P. microcarpus* to grow in culture conditions, the availability of fully sequenced fungal (Kohler *et al.*, 2015) and host (Myburg *et al.*, 2014) genomes make this an excellent model system. Using four isolates of *P. microcarpus* collected from a similar geographical area, we investigated the influence of intra-species variability on gene regulation, growth, carbon metabolism, and symbiotic

interaction with *E. grandis* at two different levels of exogenous carbon availability. Our results uncovered larger than expected variation between these closely related isolates with substantial differences in how they behave transcriptomically, metabolically, and in symbiosis with a host. Together, these results highlight the importance of using a wider range of individuals within a species to understand the broader ecological roles of ECM fungi and their host interactions.

Results

Isolates of *P. microcarpus* show distinct patterns of gene expression

Four isolates of *Pisolithus*, originally harvested from various locations in Eastern New South Wales, Australia and maintained in pure culture (Plett *et al.*, 2015), were determined by ITS-based phylogenetic analysis to all be *P. microcarpus* (MW3, Si9, R10 and R4; Figure 1A). To assess the level of transcriptomic variability between the isolates, MW3, Si9, R10 and R4 were grown for four weeks axenically and RNA from three biological replicates for each was sequenced (Figure 1B). Principle components analysis (PCA) of the resulting normalized and regularized-log transformed transcriptomes demonstrated that gene expression within replicates of each *P. microcarpus* isolate was quite consistent, but that there were distinct differences in gene expression profiles between isolates (Figure 1C, Supplemental Figure 1A). The first three principle components together account for 96.46% of the variation. The genes most implicated in this separation (820 genes total; loading coordinates with $r > 0.75$ in a loading plot for PC1, 2 and 3; Supplemental Figure 1B) are listed in Supplemental Table 1. A GO enrichment analysis of these genes highlights that processes involved in protein modification (i.e. methylation, alkylation) and in DNA integration were what most differentiated the transcriptome between the isolates (Table 1).

P. microcarpus isolates have similar biomass accumulation but different carbon use strategies depending on exogenous glucose concentration

Given the differences in gene expression between the four *P. microcarpus* isolates, we investigated whether this resulted in similar alterations to isolate growth and biomass accumulation, carbon metabolism and ability to colonize a host. These traits were assessed at two concentrations of exogenous glucose (1.0 g/L and 0.1 g/L) to assess potential differential responses to the abiotic constraint of carbon limitation. Following two weeks of growth at the reduced glucose condition, the mycelium of all isolates formed colonies that were both smaller and less dense than those grown with higher levels of glucose (Figure 2A, B), resulting in significantly less biomass (approximately 10% of the biomass obtained in the higher glucose condition; Figure 2C). Isolate was not a significant factor in biomass accumulation, although there was a significant isolate x glucose level effect based on two-way ANOVA ($p < 0.05$). Following an additional two-weeks of growth on the media, the sugar composition of the mycelium was analysed by GC-MS (Figure 2D). While the macroscopic growth rate of each isolate was similar at a given glucose concentration, the sugar composition in the mycelium of the *P. microcarpus* isolates showed distinct differences. MW3 and Si9 were the most distinct in their carbon metabolism at a glucose concentration of 1.0 g/L exhibiting high levels of free glucose and myo-inositol (Figure 2D,E). At 0.1 g/L, this profile changed to primarily high trehalose accumulation. The sugar profiles of R10 and R4 were impacted to a lesser degree by the glucose concentration, with sugar alcohols making up a large proportion of the free sugars, along with trehalose and relatively little glucose. R4 was unique in containing substantial amounts of xylitol/arabitol when grown under 0.1g/L glucose conditions.

Given the impact of both fungal genotype and glucose concentration on the composition of free sugars in the fungal mycelium, we considered the relative expression of key enzymes in

fungal sugar metabolic pathways. *P. microcarpus* encodes homologues of key carbon metabolic enzymes as found in other ectomycorrhizal basidiomycetes (Deveau et al., 2008). While the relative expression (TPM) of these enzyme genes showed different patterns between isolates and media glucose concentration (Figure 3; Supplemental Figure 2) it is interesting to note that the differences in gene expression are not as distinct as those found in the sugar metabolic profiles. This may indicate another level of regulation controlling the activity of these enzymes.

Finally, we determined the impact of *P. microcarpus* isolate and exogenous glucose concentration on the ability to colonize *E. grandis* roots. All four isolates of *P. microcarpus* colonized *E. grandis* roots to varying extents, with R10 and R4 having the highest level of colonization under regular glucose conditions (Figure 4A). When fungal isolates were grown on 0.1g/L glucose, the percentage of *E. grandis* root tips colonized decreased for all isolates, except for R4, which was able to colonize a greater percentage of root tips at reduced glucose (Figure 4A). Microscopic observation of cross sections of mycorrhizal root tips demonstrated that all isolates were able to form mycorrhizas with normal formation of the Hartig net. The depth of penetration of the Hartig net into the root tissue varied with both isolate and media glucose concentration with MW3 penetrating into the root apoplast the deepest under 1.0g/L glucose conditions (Figure 4B). The amount of glucose in the media, the fungal isolate and their interaction were all significant factors in determining the outcome of both mycorrhization and Hartig net depth (Two-way ANOVA analysis; $p < 0.05$).

Fungal gene expression during colonization identifies both fungal isolate-independent and dependent expression patterns

Differential gene expression analysis was considered for the four *P. microcarpus* isolates after they had successfully colonized a host plant to determine if the isolate-related transcriptomic variations seen in free living mycelium would continue or if genes induced by the presence of a host were fairly uniform. RNA was isolated and sequenced from each *P. microcarpus* isolate after the formation of mature colonized root tips with *E. grandis*. Tissue was taken from the colonized root tips (ECM condition) and from attached extra-matrical mycelium (EMM condition) when grown on 1.0 g/L glucose and gene expression was compared to the free-living mycelium grown axenically without a plant present (FLM condition). All genes differentially regulated in the ECM condition as compared to FLM ($p < 0.05$, $|\log_2FC| > 1$) across all four isolates were considered for a weighted gene co-expression network analysis (WGCNA; Figure 5A; Supplemental Figure 3). Relative gene expression (TPM) for all isolates and tissue types was used in the generation of the network (34 samples in total). The network analysis demonstrates that isolate identity was more relevant in describing gene co-regulation and network modularity than tissue type, with R10 and R4 responding the most similarly to each other. A comparison of all significantly differentially regulated genes in ECM versus FLM tissues across the four isolates (Supplemental Table 2) showed that most were isolate-specific (78.9% and 72.7% of up- and down-regulated genes, respectively), although there was also a core set of isolate-independent genes (103 up-regulated, 170 down-regulated) that are similarly expressed in at least three of the four isolates (Figure 5B,C; Supplemental Figure 3,4). While almost half of these genes are unknown in function, a large portion fall under the categories of small secreted proteins (SSPs), nutrient transporters, and carbohydrate active enzymes (CAZymes; Table 2).

A large percentage of host genes regulated during *P. microcarpus* colonization are fungal isolate-specific

Given the differences in gene expression seen among the four *P. microcarpus* isolates, we considered the transcriptomic response of the host plant in colonized root tissues. Between 5,787 (Si9 colonized) and 9,708 (R4 colonized) genes were significantly differentially regulated in *E. grandis* colonized root tips as compared to axenic control roots ($p < 0.05$, $|\log_2FC| > 1$; Supplemental Table 3). Of the many differentially regulated genes in the plant roots, a large proportion (18.7-34.2%) were fungal isolate-specific – that is, significantly differentially regulated when in contact with one isolate of *P. microcarpus*, but not with any of the others (Figure 6A, B). In order to determine if these unique genes represented different parts of the same plant pathways/processes or were truly unique plant responses, we performed GO enrichment analyses on each set of fungal isolate specific genes (Supplemental Table 4). Comparison of the enriched host GO terms influenced by each fungal isolate showed that there was very little overlap in function (Figure C, D). Many of the host genes showed fungal isolate-specific enrichment in GO terms pertaining to hormone and other signalling pathways or defence responses.

Discussion

Using four isolates of *P. microcarpus*, all collected from eastern New South Wales, Australia, we examine the impacts of intra-species variation and show that isolate identity affects gene expression, carbon use and sugar metabolism and host interactions. This variation was observed most clearly in the substantially different sugar metabolic profiles in the mycelium of the fungal isolates and in their varied transcriptomic responses to interactions with host plant roots. These isolate-specific differences have the potential for more wide-spread effects as they also influenced gene regulation in the host tree in an isolate-responsive manner.

Carbon obtained by ECM fungi can be assimilated into biomass promoting fungal growth or stored as free sugars or glycogen within the fungal hyphae. Eventually this stored carbon will be returned to the atmosphere via fungal respiration or decomposition. When in symbiotic association with a host tree, carbon acquired by the ECM fungus is primarily in the form of glucose (López *et al.*, 2008; Nehls *et al.*, 2010), while free living mycelium may be reliant on carbon from other sources. Previous work in *P. microcarpus* has shown that glucose taken up into the mycelium is quickly assimilated into other forms, primarily trehalose, mannitol and other sugar alcohols such as arabitol or erythritol (Martin *et al.*, 1998). It is hypothesized that this fast conversion of glucose into sugar alcohols creates a greater sink strength for the mycelium to maximize glucose uptake from the host (López *et al.*, 2007; Wiemken 2007). In our study, isolates R4 and R10 followed this pattern with free sugars being made up primarily of trehalose, mannitol, and other sugar alcohols. Isolates Si9 and MW3, however, had only trace amounts of mannitol, and at higher levels of exogenous carbon input, contained large percentages of free glucose and myo-inositol. In another study using isolates of the ECM

222 fungus *Hebeloma* spp., Tibbett *et al.* (2002) also showed a range of sugar alcohol ratios in
223 isolates of the same species across a temperature gradient, however, generally the same types
224 of sugars were present within a species. Mannitol, in addition to being an important C storage
225 compound, has demonstrated roles in quenching reactive oxygen species and improving host
226 colonization in pathogenic fungi (Véléz *et al.*, 2008; Meena *et al.*, 2015). While it is
227 currently not known if a similar pattern exists in mutualistic fungi, it would be interesting to
228 see if the higher levels of mannitol seen in other isolates of *Pisolithus* correspond to higher
229 rates of *E. grandis* root colonization as was found in R4 and R10.

230

231 A comparison of genes differentially regulated in mycorrhizal root tips as compared to free
232 living mycelium demonstrated that about three quarters of all genes with altered expression
233 were isolate-specific – that is differentially regulated in one isolate of *P. microcarpus*, but not
234 in any of the others. Despite this surprisingly high level of genotype driven gene expression,
235 we were able to identify a core set of genes that were consistently regulated across the fungal
236 isolates in response to host colonization. This core set of genes included a number of small
237 secreted proteins, nutrient transporters and CAZymes, all of which are classes of genes that
238 have been previously highlighted as putative keystone genes involved in the establishment of
239 ECM colonization (Kohler *et al.*, 2015). Small secreted proteins induced by the presence of a
240 host (called Mycorrhizal induced Small Secreted Proteins (MiSSPs)) have been shown to be
241 crucial in ECM fungal-host interactions with several characterized MiSSPs linked to the
242 ability of the fungus to colonize its host. Within *P. albus*, PaMiSSP10b was recently shown
243 to impact root colonization through interacting with an enzyme in the plant polyamine
244 pathway (Plett *et al.*, 2020b). While the equivalent PmMiSSP10b was not in our core data set
245 (while up-regulated in all isolates, it did not meet our cut off criteria of $\log_2FC > 1$ for all

isolates), several other SSPs were consistently induced in the presence of a host suggesting important involvement in the colonization process. There were also a number of nutrient transporters included in the core set of genes, including the up-regulation of sugar transporters and an ammonium (Amt) transporter, potentially involved in the symbiotic transfer of nutrients between partners. Finally, CAZymes are involved in the degradation of cell wall constituents and most lineages of ECM fungi have a reduced complement of these enzymes as compared to their saprotrophic ancestors, highlighting their importance in the evolution of the symbiotic lifestyle (Martin *et al.*, 2016). In our data set, a number of CAZymes were down-regulated in symbiotic associations. In contrast, during the colonization of *Populus* roots, the ECM fungus *Laccaria bicolor* has increased expression of CAZymes, some of which are linked to the ability of the fungus to colonize its host (Veneault-Fourrey *et al.*, 2014, Zhang *et al.*, 2018). As these enzymes are required to loosen cell wall connections and allow for penetration of fungal hyphae between the plant cells and the formation of the Hartig net, it is unusual that these enzymes are down-regulated in *P. microcarpus*. As we tested fully colonized roots, it is possible that the up-regulation of these genes would have occurred at an earlier stage of symbiosis when mycorrhizal root tips were not yet fully formed.

In addition to the isolate-specific gene regulation seen in the four fungal isolates, we observed a similar isolate-specific variation in the transcriptomic response in the host *E. grandis*. This is significant as it points to the potential larger scale consequences of mycorrhizal intra-species variability on plant function, nutrition and carbon sequestration. In another study considering nutrient trading (Hortal *et al.*, 2017), the *P. microcarpus* isolates R10 and Si9 exhibited opposing nutrient trading strategies, with R10 donating very little nitrogen to the

270 plant while consuming more carbon and Si9 donating more nitrogen for less carbon.
271 Experiments with arbuscular mycorrhizal fungi have demonstrated that different isolates
272 within a species have significant effects on both the resulting biomass and nutritional status of
273 the host plant (Koch *et al.*, 2006; Mensah *et al.*, 2015). Altered gene regulation in colonized
274 roots can also have systemic effects on the whole plant and affect how the plant interacts with
275 other soil micro-organisms, including other competing mycorrhizal fungi (Johnson and
276 Gehring, 2007; Hortal *et al.*, 2017).

277

278 The availability of exogenous glucose to the fungal mycelium also produced variable
279 responses between isolates. While reduced glucose generally resulted in fewer mycorrhizal
280 root tips formed across the isolates tested, R4 was an exception. In another study considering
281 the effect of glucose availability on the colonization of *E. grandis* by the closely related *P.*
282 *albus*, reduced glucose availability resulted in more mycorrhizal root tips formed, similar to
283 R4 (Plett *et al.*, 2020a). Similarly, the depth of the Hartig net in individual root tips was
284 significantly affected by the interaction between glucose availability and isolate. These
285 findings demonstrate that conclusions about the function of an ECM species based on the
286 reaction of a single isolate to altered environmental conditions may need to be verified using
287 a variety of genotypes within that species.

288

289 Much of the research regarding the genetic basis of how ECM symbiosis functions is reliant
290 on studies in model systems. While these studies have provided us with invaluable
291 information on ECM function there is a need to complement them with studies across a wider
292 variety of species to understand the extent of species-induced variability (Stuart and Plett,
293 2020; Pellitier and Zak, 2018). Further to this, our present study expands upon previous

research on intra-species variation to demonstrate that multiple members of a given species should also be studied as different isolates of the same species may have substantially different traits and affect their hosts and environment in correspondingly distinct ways. The impact of this observed intra-species variability on plant performance in natural ecosystems is understudied, however, indications are that genetic diversity within an ECM species is an important mediator of plant performance (Johnson *et al.*, 2012; Hazard *et al.*, 2017). The contribution of intra-species variation may be amplified where the physiological traits of individuals are more diverse, allowing the ecosystem to respond more readily to change (Hughes *et al.*, 2008). Thus, an understanding of the range of physiological traits present within a species can assist in predicting the resilience of an ecosystem and how it may adapt to change, as well as contributing to a greater understanding of the mechanisms behind ECM function and responses to their abiotic and biotic environment.

308 **Experimental Procedures**

309

310 *Biological materials*

311 *E. grandis* seeds were obtained from the CSIRO Australian Tree Seed Centre (seedlot
312 #20974) and surface sterilized in 30% hydrogen peroxide for ten minutes and rinsed three
313 times in sterile water prior to germinating on 1% agar plates. After one month of growth
314 (25°C, 16-hour photoperiod) the seedlings were transferred onto half-strength modified Melin
315 Norkrans (MMN) media (1.0 g/L glucose) covered with a sterile cellophane membrane to
316 prevent ingrowth of the root into the media. Seedlings were allowed to grow for another
317 month under the same conditions prior to contact with *P. microcarpus*.
318 *P. microcarpus* isolates used in this experiment were MW3, Si9, R10 and R4 as described in
319 Plett *et al.* (2015). Isolates were sub-cultured onto full-strength MMN media (10g/L glucose)
320 and grown for one month and then sub-cultured again onto half-strength MMN media
321 covered in a sterile cellophane membrane containing either 1.0 g/L (regular glucose level) or
322 0.1 g/L (reduced glucose level) glucose. These colonies were allowed to grow to four weeks
323 for sugar composition analysis or RNA-sequencing of free-living mycelium. The perimeter
324 of the colony was marked at 2 and 4-weeks for growth rate.

325

326 *Mycorrhization experiments*

327 The roots of *E. grandis* seedlings were placed in contact with colonies of two-week old pre-
328 grown *P. microcarpus* isolates. Plates containing plants only or fungus only were included as
329 controls. The plates were sealed with Micropore tape to allow for gas exchange and placed at
330 a 45° angle in a growth chamber (Conviron, Winnipeg, MB, Canada; 30°C daytime/22°C
331 night, 16 hour photoperiod). The experiment was harvested at 14 days after contact.

332

333 At harvest, the percentage of mycorrhization for each root system was calculated (number of
334 mycorrhizal root tips/total lateral roots x 100%) based on a visual assessment of the lateral
335 roots in contact with fungus. The perimeter of the fungal colonies at harvest were marked
336 and photographed. The total areas of the fungal colonies at two and four weeks were
337 calculated using ImageJ software. Mycorrhizal root tips were excised and placed in 4%
338 paraformaldehyde (in 1xPBS) for microscopic analysis of the Hartig net. Three biological
339 replicates of free-living mycelium (FLM; axenically grown), extra-matrical mycelium
340 (EMM) and ectomycorrhizal roots (ECM) were collected for each fungal isolate and glucose
341 concentration for RNA and immediately frozen in liquid nitrogen. Roots of axenically grown
342 *E. grandis* were also collected for RNA. FLM tissue was also collected for sugar analysis
343 and frozen immediately in liquid nitrogen. Finally, radial portions of the fungal FLM
344 colonies were excised (area excised determined by ImageJ analysis), dried and weighed to
345 determine the density of the fungal colonies. Fungal dry weight colony biomass data was
346 calculated using each colony area multiplied by its density.

347

348 *Microscopic analysis*

349 Excised root tips in 4% paraformaldehyde were rinsed 3 times in 1 x PBS and then embedded
350 in 6% agarose. Embedded roots were sectioned on a vibratome (30 µm thick), stained with
351 wheat germ agglutinin (WGA) -FITC and propidium iodide to visualize the plant and fungal
352 cells, rinsed in 1 x PBS and mounted on microscope slides. Cross sections were visualized on
353 a confocal microscope (Leica type) and ImageJ software was used to calculate the Hartig net
354 depth. A minimum of three root tips from each sample type were observed.

355

356 *Sugar analysis*

357 Fungal mycelium from five biological replicates for each of the four *P. microcarpus* isolates,
358 at both normal and low glucose conditions, was collected and snap frozen in liquid nitrogen.
359 The sample was transferred to a 2-mL screw top vial with a glass bead and 250 μ L methanol
360 and homogenized using a bead mill (FastPrep24, MP Biomedicals). Samples were incubated
361 at 30°C for 15 minutes with shaking and then centrifuged. The supernatant was collected and
362 the remaining pellet was extracted again with 250 μ L of water at room temperature.
363 Following centrifugation, this aqueous extract was added to the methanol extract.
364 25-200 μ L of metabolite containing extract (depending on input mass) from above was added
365 into a 0.1mL micro-insert for a standard 2.0 mL GC crimp top sample vial. The metabolite
366 solution was dried down completely using a vacuum concentrator (SpeedVac, Thermo Fisher
367 Scientific) set at ambient temperature. The micro-inserts were sealed into 2.0 mL GC sample
368 vials and 20 μ L of pyridine (Sigma-Aldrich) and 20 μ L of N,O-
369 Bis(trimethylsilyl)trifluoroacetamide (BSTFA; Sigma-Aldrich) was injected into each
370 sample. Samples were incubated with shaking for 2 hours at 37°C and then 1 μ L of the
371 solution was injected onto a GC-MS instrument using a splitless injection (Agilent
372 Technologies 7890A GC system with a 5975C-XL MSD and a Gerstel Maestro Prep
373 autosampler). Separation was achieved using a Agilent HP-5MS column (30 m length; 0.250
374 mm diameter; 0.25 μ m film) with a flowrate of 1.2 mL/min and an initial oven temperature of
375 70°C, held for one minute and then increased at 7 degrees per minute to 300°C and held at
376 that temperature for 5 minutes. Standards for identified sugars were also run using the same
377 derivatization procedure to ensure correct identification of sugars based on retention times
378 and mass spectra. Ratios between integrated peak areas were used to calculate the percent
379 composition of each sample.

Generation of *Pisolithus* phylogenetic tree

RNA sequencing

RNA was extracted from three biological replicates of frozen FLM, EMM, ECM and plant control tissues using either the Bioline miRNA extraction kit or the Qiagen RNeasy Plant RNA kit. All samples for RNA sequencing were harvested 4 weeks after inoculation of the fungal isolate onto the test plate and, for plates with plants present, after the formation of mature mycorrhizal root tips (about two weeks after physical contact between the two organisms). RNA was sequenced at the Western Sydney University Next Generation Sequencing facility (FLM and ECM samples, 1.0 g/L glucose) or at the Joint Genome Institute (*E. grandis* controls, EMM and 0.1 g/L glucose FLM samples). (JGI please insert details of library preparation and sequencing here).

Transcriptomic analysis

Raw reads were trimmed for low quality (quality score 0.05), Illumina adapters and sequences shorter than 15 nucleotides and aligned to the reference transcripts available at JGI (*Pisolithus microcarpus* 441; <https://mycocosm.jgi.doe.gov/Pismi1/Pismi1.home.html>) using CLC Genomics Workbench v12 (Qiagen). The following CLC genomic workbench parameters were used for read mapping: minimum length fraction 0.9, minimum similarity fraction 0.8, Mismatch cost = 2, insertion cost = 3, deletion cost = 3, and the maximum number of hits for a read was set to 10. The unique and total mapped reads number for each

transcript were determined and then normalized to transcripts per million (TPM). The mapping for the different isolates is summarized in Supplemental table 5. DESeq2 (version 1.24) was used to determine which genes were differentially expressed between pairs of conditions (Love *et al.*, 2014). The parameters used to call a gene differentially expressed between conditions were log2 fold change > 1 and FDR $p < 0.05$. For each isolate the unique mapped reads from FLM, EMM and ECM were normalised together and the log2 fold change (log2 FC) in gene expression level between conditions was calculated for ECM versus FLM and EMM versus FLM identified.

411

Protein sequences for sugar biosynthetic enzymes from the yeast (*Saccharomyces cerevisiae*) proteome were obtained from www.uniprot.org and a protein-protein BLAST was performed against the *P. microcarpus* 441 proteome at mycocosm.jgi.doe.gov to identify the corresponding *P. microcarpus* homologues. The normalized expression of each *P. microcarpus* sugar biosynthetic enzyme homologue was determined in the four isolates from the TPM data calculated as above.

418

419 *Network generation*

We performed semi-supervised weighted gene co-expression network analysis (WGCNA) (Langfelder and Horvath, 2008), including only differential regulated transcripts (Log2 FC > 1, $p < .05$, ECM vs FLM, EMM vs ECM, normal glucose condition) and using TPM normalised values as input. The resulting 8,761 genes were binned into 20 colour-coded modules with a minimum module size of 150. Pearson correlation values were calculated for all genes across all samples, correlation matrix was transformed into connection strengths (connectivity) matrix using a power adjacency function, clusters of highly correlated genes were identified

by coupling linkage hierarchical clustering with topological overlap matrix and sample information were related to network properties (Supplemental Figure 3). Cytoscape (Cline *et al.*, 2007) was used to visualize the gene network.

430

Statistical and enrichment analyses

GO enrichment of *P. microcarpus* genes was performed in R Studio (v.1.2.1335; R Core Team, 2016) using the Bioconductor topGO package (Alexa and Rahnenfuhrer, 2020) for terms under Biological Process using the *P. microcarpus* GO annotations available at <https://mycocosm.jgi.doe.gov/Pismi1/Pismi1.home.html>. GO terms were considered as enriched with a corrected p-value of <0.05 based on Fisher's Exact tests.

GO enrichment analysis for isolate-specific differentially regulated genes in *E. grandis* was conducted using the online platform PlantRegMap (<http://plantregmap.cbi.pku.edu.cn/>; Tian *et al.*, 2020) using dcGO and Fisher's Exact tests for Biological Process using the *E. grandis* genome as the species background. GO terms were considered as enriched with a corrected p-value of <0.001.

Statistical analyses were performed in R. For PCA analysis of RNA sequencing data, mapped count data from the RNA sequencing for the free living mycelium of the four *P. microcarpus* isolates (three biological replicates per isolate) at 1.0 g/L glucose was trimmed to remove non-expressed or lowly expressed genes (<5 counts on average across the three replicates across all conditions). Gene counts were normalized and rlog transformed using the "DeSEQ2" package in R (Love *et al.*, 2014) and the resulting data frame was used for generation of the PCA plots (prcomp). The "plotly" package was used for 3D PCA visualization. Loadings data for each gene for PC1, PC2 and PC3 were acquired and those genes falling outside of a sphere of radius 0.75 on the corresponding loadings plot were

considered to contribute most to the separation observed on the PCA plot. Two-way ANOVAs (unbalanced, Type III) were used to determine the effect of isolate or glucose availability and their interaction on experimental outcomes using the “car” package (Fox and Weisberg, 2011; R Core Team, 2016). Significant differences between two experimental conditions were calculated using a single tailed Student’s T-test. Clustering of the sugar composition data used hierarchical clustering using a one minus Pearson correlation metric to calculate distances between observations. Heatmaps were generated using the Morpheus online tool at <https://software.broadinstitute.org> and Venn diagrams were generated using the online tool found at <http://bioinformatics.psb.ugent.be/webtools/Venn/>.

Acknowledgements

The authors wish to acknowledge the Australian Research Council for a Discovery grant to ICA, JMP and FM (DPXX), the Western Sydney University Next Generation Sequencing facility for RNA sequencing and the Western Sydney University Confocal facility for access to confocal microscopy. Thanks to J. Huntley for assistance with the sugar analyses.

469 **References**

470

471 Alexa, A., and Rahnenfuhrer, J. (2020) topGO: Enrichment analysis for gene ontology. R
472 package version 2.40.0.

473

474 Anderson, I.C., Chambers, S.M., and Cairney, J.W.G. (1999) Intra- and interspecific variation
475 in patterns of organic and inorganic nitrogen utilization by three Australian *Pisolithus*
476 species. Mycol Res 103:2.

477

478 Clemmensen, K.E., Finlay, R.D., Dahlberg, A., Stenlid, J., Wardle, D.A., and Lindahl, B.D.
479 (2015) Carbon sequestration is related to mycorrhizal fungal community shifts during long-
480 term succession in boreal forests. New Phytol 205: 1525-1536.

481

482 Cline, M.S., Smoot, M., Cerami, E., Kuchinsky, A., Landys, N., Workman, C., *et al.* (2007)
483 Integration of biological networks and gene expression data using Cytoscape. Nat Protoc
484 2:2366-2382.

485

486 Deveau, A., Kohler, A., Frey-Klett, P., and Martin, F. (2008) The major pathways of
487 carbohydrate metabolism in the ectomycorrhizal basidiomycete *Laccaria bicolor* S238N.
488 New Phytol 180:379-390.

489

490 Ekblad, A., Wallander, H., Godbold, D.L., Cruz, C., Johnson, D., Baldrian,
491 P., *et al.* (2013) The production and turnover of extramatrical mycelium of ectomycorrhizal
492 fungi in forest soils: role in carbon cycling. Plant and Soil 366: 1–27.

493

494 Fox, J., and Weisberg, S. (2011) An R Companion to Applied Regression, Second Edition.

495 Thousand Oaks CA: Sage.

496

497

498 Fry, E.L., De Long, J.R., Garrido, L.A., Alvarez, N., Carrillo, Y., Castañeda-Gómez, L., *et al.*

499 (2019) Using plant, microbe, and soil fauna traits to improve the predictive power of

500 biogeochemical models. *Methods Ecol Evol* 10:146-157.

501

502 Guidot, A., Verner, M.C., Debaud, J.C., and Marmeisse, R. (2005) Intraspecific variation in

503 use of different organic nitrogen sources by the ectomycorrhizal fungus *Hebeloma*

504 *cylindrosporum*. *Mycorrhiza* 15:3.

505

506 Hazard, C., Kruitbos, L., Davidson, H., Mbow, F.T., Taylor, A.F.S., and Johnson, D. (2017)

507 Strain identity of the ectomycorrhizal fungus *Laccaria bicolor* is more important than

508 richness in regulating plant and fungal performance under nutrient rich conditions. *Front*

509 *Microbiol* 8:1874.

510

511 Hibbett, D.S., Gilbert, L.B., and Donoghue, M.J. (2000) Evolutionary instability of

512 ectomycorrhizal symbioses in basidiomycetes. *Nature* 407:6803.

513

514 Hortal, S., Plett, K.L., Plett, J.M., Cresswell, T., Johansen, M., Pendall, E., et al. (2017) Role
 515 of plant-fungal nutrient trading and host control in determining the competitive success of
 516 ectomycorrhizal fungi. *ISME J* 11:12.
 517
 518 Hughes, A. R., Inouye, B.D., Johnson, M.T.J., Underwood, N., Vellend, M. (2008)
 519 Ecological consequences of genetic diversity. *Ecol Lett* 11:609-623.
 520
 521 Johnson, D., Martin, F., Cairney J.W.G., Anderson, I.C. (2012) The importance of
 522 individuals: intraspecific diversity of mycorrhizal plants and fungi in ecosystems. *New Phytol*
 523 194:614-628.
 524
 525 Johnson, N.C., and Gehring, C.A. (2007) Mycorrhizas: Symbiotic mediators of rhizosphere
 526 and ecosystem processes. In *The Rhizosphere: an ecological perspective*. Eds Cardon, Z.G.,
 527 and Whitbeck, J.L. Academic Press pp. 73-100.
 528
 529 Koch, A.M., Croll, D., and Sanders, I.R. (2006) Genetic variability in a population of
 530 arbuscular mycorrhizal fungi causes variation in plant growth. *Ecol Lett* 9:103-110.
 531
 532 Kohler, A., Kuo, A., Nagy, L.G., Morin, E., Barry, K.W., Buscot, F., et al. (2015)
 533 Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in
 534 mycorrhizal mutualists. *Nature Genet* 47:4.

535

536 Langfelder, P., and Horvath, S. (2008) WGCNA: an R package for weighted correlation
 537 network analysis. BMC Bioinformatics 9:559.

538

539 Le Tacon, F., Zeller, B., Plain, C., Hossann, C., Bréchet, C., Martin, F., *et al.* (2015) Study of
 540 nitrogen and carbon transfer from soil organic matter to *Tuber melanosporum* mycorrhizas
 541 and ascocarps using ¹⁵N and ¹³C soil labelling and whole-genome oligoarrays. Plant Soil
 542 395:1-2.

543

544 Lindahl, B.D., and Tunlid, A. (2015) Ectomycorrhizal fungi – potential organic matter
 545 decomposers, yet not saprotrophs. New Phytol 205:4.

546

547 Lindahl, B.D., Ihrmark, K., Boberg, J., Trumbore, S.E., Högberg, P., Stenlid, J., *et al.* (2007).
 548 Spatial separation of litter decomposition and mycorrhizal nitrogen uptake in a boreal forest.
 549 New Phytol 173:3.

550

551 López, M.F., Dietz, S., Grunze, N., Bloeschies, J., Weiß, M., and Nehls, U. (2008) The sugar
 552 porter gene family of *Laccaria bicolor*: function in ectomycorrhizal symbiosis and soil-
 553 growing hyphae. New Phytol 180:2.

554

555 López, M.F., Manner, P., Willmann, A., Hampp, R., and Nehls, U. (2007) Increased trehalose
 556 biosynthesis in the Hartig net hyphae of ectomycorrhizas. New Phytol 174: 389-398.

557

558 Love, M.I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and
 559 dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

560

561 Martin, F., Boiffin, V., and Pfeffer, P.E. (1998) Carbohydrate and amino acid metabolism in
562 the *Eucalyptus globulus*-*Pisolithus tinctorius* ectomycorrhiza during glucose utilization. Plant
563 Physiol 118: 627-635.

564

565 Martin, F., Kohler, A., Murat, C., Veneault-Fourrey, C., and Hibbett, D.S. (2016) Unearthing
566 the roots of ectomycorrhizal symbioses. Nature Rev Microbiol 14:760-773.

567

568 Meena, M., Prasad, V., Zehra, A., Gupta, V.K., and Upadhyay, R.S. (2015) Mannitol
569 metabolism during pathogenic fungal-host interactions under stressed conditions. Front
570 Microbiol 6: 1019.

571

572 Mensah, J.A., Koch, A.M., Antunes, P.M., Kiers, E.T., Hart, M., and Bucking, H. (2015)
573 High functional diversity within species of arbuscular mycorrhizal fungi is associated with
574 differences in phosphate and nitrogen uptake and fungal phosphate metabolism. Mycorrhiza
575 25:533-546.

576

577 Myberg, A.A., Grattapaglia, D., Tuskan, G.A., Hellsten, U., Hayes, R.D., Grimwood, J., *et al.*
578 (2014) The genome of *Eucalyptus grandis*. Nature 510:356-362.

579

580 Nehls, U., Gohringer, F., Wittulsky, S., and Dietz, S. (2010) Fungal carbohydrate support in
581 the ectomycorrhizal symbiosis: a review. Plant Biol 12:292-301.

582

583 Orwin, K.H., Kirschbaum, M.U., St John, M.G., and Dickie, I.A. (2011) Organic nutrient
 584 uptake by mycorrhizal fungi enhances ecosystem carbon storage: a model-based assessment.
 585 Ecol Lett 14:5.
 586
 587 Pellitier, P.T., and Zak, D.R. (2018). Ectomycorrhizal fungi and the enzymatic liberation of
 588 nitrogen from soil organic matter: why evolutionary history matters. New Phytol 217:1.
 589
 590 Plett, J.M., Kohler, A., Khachane, A., Keniry, K., Plett, K.L., Martin, F., *et al.* (2015) The
 591 effect of elevated carbon dioxide on the interaction between *Eucalyptus grandis* and diverse
 592 isolates of *Pisolithus* sp. is associated with a complex shift in the root transcriptome. New
 593 Phytol 206:4.
 594
 595 Plett, J.M., Plett, K.L., Wong-Bajracharya, J., de Freitas Pereira, M., Dutra Costa, M.,
 596 Kohler, A., *et al.* (2020b) Mycorrhizal effector PaMiSSP10b alters polyamine biosynthesis in
 597 *Eucalyptus* root cells and promotes root colonization. New Phytol
 598 doi.org/10.1111/nph.16759.
 599
 600 Plett, K.L., Singan, V.R., Wang, M., Ng, V., Grigoriev, I.V., Martin, F., *et al.* (2020a)
 601 Inorganic nitrogen availability alters *Eucalyptus grandis* receptivity to the ectomycorrhizal
 602 fungus *Pisolithus albus* but not symbiotic nitrogen transfer. New Phytol 226:221-231.
 603
 604 R Core Team. (2016) R: A language and environment for statistical computing. R Foundation
 605 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
 606

607 Read, D.J., and Perez-Moreno, J. (2003) Mycorrhizas and nutrient cycling in ecosystems – a
608 journey towards relevance? *New Phytol* 157:3.

609

610 Sawyer, N.A., Chambers, S.M., and Cairney, J.W.G. (2003) Variation in nitrogen source
611 utilisation by nine *Amanita muscaria* genotypes from Australian *Pinus radiata* plantations.
612 *Mycorrhiza* 13:4.

613

614 Shah, F., Nicolás, C., Bentzer, J., Ellström, M., Smits, M., Rineau, F., *et al.* (2016)
615 Ectomycorrhizal fungi decompose soil organic matter using oxidative mechanisms adapted
616 from saprotrophic ancestors. *New Phytol* 209:4.

617

618 Smith, S., and Read, D. (2008) *Mycorrhizal Symbiosis* 3rd Edition. Academic Press.

619

620 Stuart, E.K., and Plett, K.L. (2020) Digging deeper: in search of the mechanisms of carbon
621 and nitrogen exchange in ectomycorrhizal symbioses. *Front Plant Sci* 10:1658.

622

623 Sulman, B.N., Brzostek, E.R., Medici, C., Shevliakova, E., Menge, D.N.L., and Phillips, R.P.
624 (2017) Feedbacks between plant N demand and rhizosphere priming depend on type of
625 mycorrhizal association. *Ecol Lett* 20:1043-1053.

626

627 Tedersoo, L., and Smith, M.E. (2013) Lineages of ectomycorrhizal fungi revisited: Foraging
628 strategies and novel lineages revealed by sequences from belowground. *Fungal Biol Rev*
629 27:3-4.

630

631 Tian, F., Yang, D.C., Meng, Y.Q., Jin, J. and Gao, G. (2020) PlantRegMap: charting
632 functional regulatory maps in plants. *Nucleic Acids Res* 48, D1104-D1113.

633

634 Tibbett, M., Sanders, F.E., and Cairney, J.W.G. (2002) Low-temperature-induced changes in
635 trehalose, mannitol and arabitol associated with enhanced tolerance to freezing in
636 ectomycorrhizal basidiomycetes (*Hebeloma* spp.). *Mycorrhiza* 12:249-255.

637

638 Vélèz, H., Glassbrook, N.J., and Daub, M.E. (2008) Mannitol biosynthesis is required for
639 plant pathogenicity by *Alternaria alternata*. *FEMS Microbiol Lett* 285 122–129.

640

641 Veneault-Fourrey, C., Commun, C., Kohler, A., Morin, E., Balestrini, R., Plett, J., *et al.*
642 (2014) Genomic and transcriptomic analysis of *Laccaria bicolor* CAZome reveals insights
643 into polysaccharides remodelling during symbiosis establishment. *Fungal Genet Biol* 72:168-
644 181.

645

646 Wiemken, V. (2007) Trehalose synthesis in ectomycorrhizas – a driving force of carbon gain
647 for fungi? *New Phytol* 174: 228-230.

648

649 Zak, D.R., Pellitier, P.T., Argiroff, W.A., Castillo, B., James, T.Y., Nave, L.E., *et al.* (2019)
650 Exploring the role of ectomycorrhizal fungi in soil carbon dynamics. *New Phytol* 223:1.

651

652 Zhang, F., Anasontzis, G.E., Labourel, A., Champion, C., Hoan, M., Kemppainen, M., *et al.*
653 (2018) The ectomycorrhizal basidiomycete *Laccaria bicolor* releases a secreted β -1,4
654 endoglucanase that plays a key role in symbiosis development. *New Phytol* 220:1309-1321.

655 Table 1

656

GO ID	Term	P value (Fisher)
GO:00150 74	DNA integration	0.015
GO:00064 79	protein methylation	0.025
GO:00064 81	C-terminal protein methylation	0.025
GO:00082 13	protein alkylation	0.025
GO:00184 10	C-terminal protein amino acid modification	0.025
GO:00065 58	L-phenylalanine metabolic process	0.037
GO:00065 59	L-phenylalanine catabolic process	0.037
GO:00068 29	zinc ion transport	0.037
GO:00094 39	cyanate metabolic process	0.037

GO:19022	erythrose	0.037
21	4-phosphate/phosphoenolpyruvate family amino acid metabolic process	
GO:19022	erythrose	0.037
22	4-phosphate/phosphoenolpyruvate family amino acid catabolic process	

657

658 **Table 1 Gene ontology (GO) terms significantly enriched in genes most contributing to**
659 **the separation of the transcriptomes of Si9, R10, R4 and MW3 in principal components**
660 **analysis.** Gene ontology (GO) enrichment was performed with dcGO on 820 *P. microcarpus*
661 genes having loading coordinates with $r > 0.75$ in a loading plot for PC1, 2 and 3 from PCA
662 analysis of the transcriptomes of Si9, R10, R4 and MW3 (Free-living mycelium, 1.0 g/L
663 glucose). Significantly enriched terms have $p < 0.05$ based on a Fisher's exact test.

664

665 **Table 2 List of *P. microcarpus* genes significantly up- or down- regulated in the core**
666 **response of *P. microcarpus* to colonization of *E. grandis* root tips.** Genes listed are
667 similarly regulated in at least 3 of the 4 isolates considered and are from the gene categories
668 of small-secreted proteins (SSPs), transporters, or cell-wall active enzymes (CAZymes). A
669 complete list of all core genes and the expression level of each gene can be found in
670 Supplemental Figures 4 and 5.
671

Up-regulated Core Genes		Down-regulated Core Genes	
Small Secreted Proteins (SSPs)			
Protein ID	Description	Protein ID	Description
685288	Carbonic anhydrase	22402	SSP Cerato-platanin
683008	SSP	672065	Fungal hydrophobin
685028	SSP	676962	Fungal hydrophobin
687566	SSP	676970	Fungal hydrophobin
689601	SSP	677031	Fungal hydrophobin
689604	SSP	278092	SSP
683015	SSP Cerato-platanin	439320	SSP
682669	SSP Conotoxin	674112	SSP
674767	Unknown	675746	SSP
676755	Unknown SSP	688218	SSP
		688221	SSP
		679157	SSP
		97211	SSP
		676273	SSP
		672747	SSP, contains WSC domain
		690542	Unknown SSP
Transporters			
Protein ID	Description	Protein ID	Description
244660	ABC transporter	673253	Ca ²⁺ :Cation Antiporter (CaCA) Family
690664	H ⁺ or Na ⁺ translocating ATPase	422614	Major Facilitator Superfamily (MFS)
	Major Facilitator Superfamily		Monovalent Cation:Proton
22082	(MFS)	10179	Antiporter-2
680085	Major Facilitator Superfamily	678674	MOP Flippase Superfamily

91885 59181 670385 113068 675820	(MFS) Major Facilitator Superfamily (MFS) Mitochondrial Carrier (MC) Family Oligopeptide Transporter (OPT) Family Sugar transporter, (MFS) Ammonia Transporter Amt Family	261938	Oligopeptide Transporter (OPT) Family
CAZymes			
Protein ID	Description	Protein ID	Description
102510 351590	Laccase Laccase	680524 102842 104390 91890 684101 672122 678857 611459 651808 28092 636179	AA1_1 Laccase AA1_1 Laccase AA9 AA9 GH16 GH16 GH16 GH18 Chitinase GH5_15 GH71 GH79

672

673

674

675 **Figure legends**

676

677 **Figure 1. Genomic and transcriptomic variation between *P. microcarpus* isolates MW3,**
678 **Si9, R10 and R4.** (A) Phylogenetic tree based on ITS sequence showing the relationship of
679 the four studied *P. microcarpus* isolates (in blue) with other *Pisolithus* isolates including
680 Australian isolates (in red) and *P. microcarpus* isolate 441 (in green). (B) Photos of the
681 studied *P. microcarpus* isolates following two weeks of growth on ½ strength MMN media
682 with 1.0 g/L glucose. (C) 3D PCA plot of normalized and rlog transformed transcriptomic
683 data (3 biological replicates per condition) for MW3, Si9, R10 and R4 grown for four weeks
684 on ½ strength MMN media.

685

686 **Figure 2. Glucose availability affects fungal biomass accumulation and sugar**
687 **metabolism.** Two-week old colonies of *P. microcarpus* isolate R10 growing on media
688 containing 1.0 g/L glucose (A) or 0.1 g/L glucose (B). (C) Dry weight biomass of *P.*
689 *microcarpus* isolates after two weeks of growth with normal glucose levels (1.0 g/L; light
690 grey bars) or low glucose levels (0.1 g/L; dark grey bars). +/- SE. Results from two-way
691 ANOVA assessing the impact of glucose availability, fungal isolate and their interaction on
692 experimental outcomes are indicated. (D) Percent composition of free sugars in *P.*
693 *microcarpus* mycelium after four weeks of growth on normal (N; 1.0 g/L) or low (L; 0.1 g/L)
694 glucose media. (E) Hierarchical clustering of the *P. microcarpus* isolates at different sugar
695 concentrations based on the percent composition of free sugars in mycelium from (D).

696

697 **Figure 3. Expression of key enzymes in *P. microcarpus* sugar metabolic pathways.** (A)
698 Simplified sugar metabolic pathway for *P. microcarpus* with key enzymes indicated with a

number corresponding to enzymes in (B). Bolded sugars are those analysed for and detected in fungal mycelium samples. (B) *P. microcarpus* gene ID, closest yeast homologue and identity of the numbered enzymes from (A). The average normalized expression (TPM; n = 3) of each enzyme in each condition is indicated by heat map (heat map with individual replicates given in Supplemental Figure 2).

Figure 4. Extent of root colonisation is dependent on both glucose availability and fungal isolate. A) Percentage of lateral roots colonised after two weeks of fungal contact on media with either 1.0 g/L (light grey bars) or 0.1 g/L (dark grey bars) glucose; n=7-11. B) Depth of Hartig net in colonised root tips after two weeks of fungal contact on media with either 1.0 g/L (light grey bars) or 0.1 g/L (dark grey bars) glucose. +/- standard error, *= significant difference between media types. Results from two-way ANOVA assessing the impact of glucose availability, fungal isolate and their interaction on experimental outcomes are indicated.

Figure 5 Differentially expressed genes in *P. microcarpus* when colonizing *E. grandis* are isolate dependent but preserve a common core. (A) Visualization of weighted gene co-expression network analysis (WGCNA) for *P. microcarpus* gene expression. WGCNA was generated using all significantly differentially regulated genes (ECM vs FLM condition, p<0.05) from MW3, R10, R4 and Si9 transcriptomes of FLM, EMM and ECM tissue types at normal levels of glucose (1.0 g/L). Major gene clusters are identified with the represented *P. microcarpus* isolate (Bold text) and tissue type (Italic text). (B, C) Venn diagrams describing overlap of significantly differentially up- (B) and down- (C) regulated genes for each *P. microcarpus* isolate in the ECM condition versus FLM. Genes describing the core response

of *P. microcarpus* to its host (regulated in at least 3 of the 4 isolates) are outlined. FLM = Free living mycelium; EMM = Extra-matrical mycelium; ECM = Ectomycorrhizal.

Figure 6 Colonized *E. grandis* roots differentially express genes assigned to unique GO processes depending on *P. microcarpus* isolate identity (A, B) Venn diagrams showing the overlap of genes significantly up- (A) or down- (B) regulated in *E. grandis* when colonized by *P. microcarpus* isolates Si9, R4, R10 or MW3. Isolate specific genes are indicated with black circles. (C, D) Venn diagrams showing the overlap of GO terms (Biological Function) significantly enriched ($p < 0.05$) in the up- (C) and down- (D) regulated genes uniquely regulated by each *P. microcarpus* isolate. Main functions found enriched within the GO terms for each isolate are indicated in italics.

Supplemental Figure 1 PCA analysis and loadings graphs for transcriptomes of *P. microcarpus* isolates.

Supplemental Figure 2 Heatmap of expression (TPM) of key genes in sugar biosynthesis pathways in isolates of *P. microcarpus*.

Supplemental Figure 3 WGCNA module trait relationships and p- values for *P. microcarpus* isolates.

Supplemental Figure 4 Up-regulated Core Genes in *P. microcarpus* colonizing *E. grandis*.

746 **Supplemental Figure 5** Down-regulated Core Genes in *P. microcarpus* colonizing *E.*
747 *grandis*.

748

749 **Supplemental Table 1** List of genes, with their annotations, contributing to the separation of
750 *P. microcarpus* isolate transcriptomes in PCA analysis.

751

752 **Supplemental Table 2** List of all genes significantly differentially expressed in *P.*
753 *microcarpus* isolates Si9, R10, R4 or MW3 colonizing *E. grandis* roots as compared to
754 axenic controls.

755

756 **Supplemental Table 3** List of all genes significantly differentially expressed in *E.grandis*
757 roots when colonized by *P. microcarpus* isolates Si9, R10, R4 or MW3 as compared to
758 axenic controls.

759

760 **Supplemental Table 4** List of significantly enriched GO terms resulting from genes uniquely
761 up- or down- regulated in *E. grandis* roots when colonized by *P. microcarpus* isolates

762

763 **Supplemental Table 5** *P. microcarpus* mapping statistics