

UC Irvine

UC Irvine Previously Published Works

Title

Correction: Hom et al. Cognitive Function during the Prodromal Stage of Alzheimer's Disease in Down Syndrome: Comparing Models. *Brain Sci.* 2021, 11, 1220

Permalink

<https://escholarship.org/uc/item/9nh3r0t1>

Journal

Brain Sciences, 12(7)

ISSN

2076-3425

Authors

Hom, Christy L
Kirby, Katharine A
Ricks-Oddie, Joni
et al.

Publication Date

2022

DOI

10.3390/brainsci12070925

Peer reviewed

Correction

Correction: Hom et al. Cognitive Function during the Prodromal Stage of Alzheimer's Disease in Down Syndrome: Comparing Models. *Brain Sci.* 2021, 11, 1220

Christy L. Hom ^{1,*} , Katharine A. Kirby ², Joni Ricks-Oddie ^{2,3}, David B. Keator ¹, Sharon J. Krinsky-McHale ⁴ , Margaret B. Pulsifer ⁵, Herminia Diana Rosas ⁶, Florence Lai ⁶, Nicole Schupf ^{7,8}, Ira T. Lott ⁹ and Wayne Silverman ⁹

- ¹ Department of Psychiatry and Human Behavior, University of California, Orange, CA 92868, USA; dbkeator@uci.edu
- ² Center for Statistical Consulting, University of California, Irvine, CA 92697, USA; kathark@uci.edu (K.A.K.); jricksod@uci.edu (J.R.-O.)
- ³ Institute for Clinical and Translational Sciences, University of California, Irvine, CA 92697, USA
- ⁴ New York State Institute for Basic Research in Developmental Disabilities, Staten Island, NY 10314, USA; sharon.krinsky-mchale@opwdd.ny.gov
- ⁵ Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114, USA; mpulsifer@mgh.harvard.edu
- ⁶ Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114, USA; rosas@helix.mgh.harvard.edu (H.D.R.); flai@partners.org (F.L.)
- ⁷ Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, NY 10027, USA; ns24@cumc.columbia.edu
- ⁸ Department of Epidemiology, School of Public Health, Columbia University, New York, NY 10027, USA
- ⁹ Department of Pediatrics, University of California, Orange, CA 92868, USA; itlott@uci.edu (I.T.L.); wsilverm@uci.edu (W.S.)
- * Correspondence: homc@uci.edu; Tel.: +1-714-456-2927



Citation: Hom, C.L.; Kirby, K.A.; Ricks-Oddie, J.; Keator, D.B.; Krinsky-McHale, S.J.; Pulsifer, M.B.; Rosas, H.D.; Lai, F.; Schupf, N.; Lott, I.T.; et al. Correction: Hom et al. Cognitive Function during the Prodromal Stage of Alzheimer's Disease in Down Syndrome: Comparing Models. *Brain Sci.* 2021, 11, 1220. *Brain Sci.* 2022, 12, 925. <https://doi.org/10.3390/brainsci12070925>

Received: 23 June 2022

Accepted: 24 June 2022

Published: 15 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

We would like to submit the following correction to our recently published paper [1].

Error in Tables

In the original publication, there were mistakes in Tables 2 and 5 as published. The standard deviation listed for the CS group on the DSMSE language test was mistakenly listed as 0.07 in Table 2. The *p*-value listed for the RADD-2 Expressive Language in Table 5 was mistakenly listed as <.001 in Table 5. The Estimate value for the Purdue Pegboard was mistakenly listed as 0615 in Table 5. The corrected Tables 2 and 5 appear below.

Table 2. Means and standard deviations of raw test scores by AD status and their hypothesized cognitive domain.

| Variable (Range of Scores) | Domain | CS (n = 103) | MCI-DS (n = 41) | Mann-Whitney U (p-Value) |
|--|------------|---------------|-----------------|--------------------------|
| Block Design (0–54) | Visuomotor | 23.79 (10.49) | 17.37 (12.78) | 3.11 (0.002) |
| Boston Naming (0–27) | Language | 15.88 (5.51) | 13.23 (6.91) | 2.04 (0.041) |
| Category Fluency (0–17) | EF | 8.23 (3.18) | 6.82 (3.68) | 2.21 (0.027) |
| Cats and Dogs Switch (–17.00–61.80) [†] | EF | 9.80 (11.22) | 5.05 (11.21) | 1.82 (0.069) |
| Cued Recall (3–35) | Memory | 28.61 (6.83) | 21.38 (9.22) | 4.43 (0.0001) |
| DSMSE Language (3–52) | Language | 37.10 (9.07) | 30.80 (9.63) | 3.28 (0.001) |
| DSMSE Memory (0–23) | Memory | 14.08 (4.68) | 9.80 (4.57) | 4.62 (0.0001) |
| DSMSE Visual Spatial (2–8) | Visuomotor | 6.18 (1.09) | 5.59 (1.01) | 2.98 (0.003) |
| mMMSE-DS Anomia (4–20) | Language | 18.18 (2.20) | 16.49 (4.19) | 1.55 (0.122) |
| mMMSE-DS Concentration (0–6) | EF | 3.69 (2.10) | 2.39 (2.14) | 3.15 (0.002) |

Table 2. Cont.

| Variable (Range of Scores) | Domain | CS (n = 103) | MCI-DS (n = 41) | Mann–Whitney U (p-Value) |
|----------------------------------|------------|--------------|-----------------|--------------------------|
| mMMSE-DS Fine Motor (1–10) | Visuomotor | 8.04 (1.29) | 7.05 (2.28) | 2.58 (0.010) |
| mMMSE-DS Orientation (5–30) | EF | 25.46 (5.33) | 20.87 (6.41) | 4.75 (0.0001) |
| Purdue Pegboard Both Hands (0–8) | Visuomotor | 2.62 (1.82) | 1.51 (1.71) | 3.14 (0.002) |
| RADD-2 Digit Span Forward (0–8) | EF | 4.02 (1.66) | 2.97 (1.80) | 2.84 (0.005) |
| RADD-2 Expressive Lang. (0–16) | Language | 11.24 (3.92) | 8.80 (4.01) | 3.34 (0.001) |
| RADD-2 Receptive Lang. (0–12) | Language | 7.56 (2.51) | 6.80 (2.99) | 1.08 (0.279) |
| RADD-2 Sensorimotor (0–7) | Visuomotor | 6.70 (0.50) | 5.85 (1.50) | 3.87 (0.0001) |
| RADD-2 Similarities (0–4) | Language | 2.51 (1.49) | 1.56 (1.47) | 3.40 (0.001) |
| Rivermead Recognition (0–10) | Memory | 5.42 (3.77) | 2.21 (3.40) | 3.90 (0.0001) |
| Selective Reminding Test (1–24) | Memory | 15.50 (5.62) | 9.18 (4.32) | 5.70 (0.0001) |
| Tinetti Gait (4–12) | Visuomotor | 10.75 (1.62) | 10.37 (1.88) | 1.47 (0.143) |
| VMI (1–25) | Visuomotor | 15.45 (3.22) | 13.98 (3.40) | 1.75 (0.081) |

† Cats and Dogs is measured in seconds; all other scores are measured in points (number correct). EF = executive function.

Table 5. The relationship between sex, premorbid ID, MCI-DS, and cognitive functioning.

| Cognitive Domain | Predictor Variable | Estimate | SE | CR | Standardized Regression Weight | p-Value |
|------------------|----------------------------|----------|-------|-------|--------------------------------|---------|
| Structural | | | | | | |
| Language/EF | Sex | −0.010 | 0.075 | −0.13 | −0.01 | 0.893 |
| Language/EF | PID | −0.413 | 0.067 | −6.15 | −0.41 | <0.001 |
| Language/EF | MCI-DS | −0.238 | 0.074 | −3.23 | −0.24 | 0.001 |
| Memory | Sex | 0.008 | 0.076 | 0.11 | 0.01 | 0.916 |
| Memory | PID | −0.241 | 0.075 | −3.21 | −0.24 | 0.001 |
| Memory | MCI-DS | −0.471 | 0.67 | −7.07 | −0.47 | <0.001 |
| Visuomotor | Sex | −0.093 | 0.081 | −1.14 | −0.09 | 0.255 |
| Visuomotor | PID | −0.302 | 0.077 | −3.91 | −0.30 | <0.001 |
| Visuomotor | MCI-DS | −0.303 | 0.78 | −3.90 | −0.30 | <0.001 |
| Measurement | | | | | | |
| Language/EF | Boston Naming | 0.879 | 0.023 | 38.18 | 0.88 | <0.001 |
| Language/EF | DSMSE Language | 0.878 | 0.023 | 37.81 | 0.88 | <0.001 |
| Language/EF | mMMSE Anomia | 0.742 | 0.041 | 18.14 | 0.74 | <0.001 |
| Language/EF | mMMSE Concentration | 0.724 | 0.043 | 16.53 | 0.72 | <0.001 |
| Language/EF | RADD-2 Digit Span Forward | 0.782 | 0.036 | 21.93 | 0.78 | <0.001 |
| Language/EF | RADD-2 Expressive Language | 0.874 | 0.023 | 37.52 | 0.87 | <0.001 |
| Language/EF | RADD-2 Receptive Language | 0.683 | 0.047 | 16.53 | 0.68 | <0.001 |
| Language/EF | RADD-2 Similarities | 0.760 | 0.038 | 20.05 | 0.76 | <0.001 |
| Memory | Cued Recall | 0.673 | 0.053 | 12.65 | 0.67 | <0.001 |
| Memory | DSMSE Memory | 0.858 | 0.032 | 27.09 | 0.86 | <0.001 |
| Memory | Rivermead Recognition | 0.702 | 0.051 | 13.87 | 0.70 | <0.001 |
| Memory | Selective Reminding Test | 0.845 | 0.033 | 25.65 | 0.85 | <0.001 |
| Visuomotor | Block Design | 0.849 | 0.033 | 26.09 | 0.85 | <0.001 |
| Visuomotor | DSMSE Visual Spatial | 0.797 | 0.038 | 17.05 | 0.80 | <0.001 |
| Visuomotor | Purdue Pegboard | 0.615 | 0.59 | 10.39 | 0.62 | <0.001 |
| Visuomotor | RADD Sensorimotor | 0.624 | 0.058 | 10.70 | 0.62 | <0.001 |
| Visuomotor | VMI | 0.708 | 0.048 | 14.77 | 0.71 | <0.001 |

CR = critical ratio, EF = executive function, MCI = mild cognitive impairment, PID = premorbid level of ID, SE = standard error.

The authors apologize for any inconvenience caused and state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Reference

1. Hom, C.L.; Kirby, K.A.; Ricks-Oddie, J.; Keator, D.B.; Krinsky-McHale, S.J.; Pulsifer, M.B.; Rosas, H.D.; Lai, F.; Schupf, N.; Lott, I.T.; et al. Cognitive Function during the Prodromal Stage of Alzheimer's Disease in Down Syndrome: Comparing Models. *Brain Sci.* **2021**, *11*, 1220. [[CrossRef](#)] [[PubMed](#)]