

# UCLA

## UCLA Previously Published Works

### Title

Re-engagement in HIV care among mothers living with HIV in South Africa over 36 months post-birth

### Permalink

<https://escholarship.org/uc/item/9nj2r4xf>

### Journal

AIDS, 29(17)

### ISSN

0269-9370

### Authors

Rotheram-Borus, Mary Jane  
Tomlinson, Mark  
Scheffler, Aaron  
[et al.](#)

### Publication Date

2015-11-01

### DOI

10.1097/qad.0000000000000837

Peer reviewed



Published in final edited form as:

AIDS. 2015 November ; 29(17): 2361–2362. doi:10.1097/QAD.0000000000000837.

## Re-engagement in HIV care among Mothers Living with HIV in South Africa over 36 Months Post-birth

Mary Jane Rotheram-Borus<sup>1</sup>, Mark Tomlinson<sup>2</sup>, Aaron Scheffler<sup>1</sup>, and Ingrid M. Le Roux<sup>3</sup>

<sup>1</sup>Department of Psychiatry and Biobehavioral Medicine, Semel Institute, University of California at Los Angeles, 10920 Wilshire Blvd., Suite 350, Los Angeles, California 90024 USA

<sup>2</sup>Department of Psychology, Stellenbosch University, Private Bag X1 Matieland, 7602 South Africa

<sup>3</sup>Philani Maternal, Child Health and Nutrition Project, PO Box 40188, Elonwabeni, Cape Town, 7791 South Africa

### Summary

PMTCT services are integrated into antenatal care in South Africa, but post-birth care is at HIV clinics. Almost all Mothers Living with HIV (MLH) in 24 township neighborhoods (N=324) reported engagement in HIV care from pregnancy to 36 months post-birth. Less than half re-engaged in HIV care at 6 months (45%), 52.5% at 18 months, and 62.5% at 36 months. Most were prescribed ARVs if reengaged in care, yet only about half (53%) are on ARVs at 36 months post birth. Implementation of Option B+ will require substantially better engagement in care.

---

In Cape Town, South Africa, 26-30% of pregnant women are infected with HIV. Fortunately, South Africa has broad availability and uptake of HIV testing and ARV therapies during the perinatal period for both mothers and children [1]. The country is planning to implement the B+ protocol which will guarantee all Mothers Living with HIV (MLH) lifelong ARV [2]. The B+ protocol will heighten the importance of maintaining HIV care over time. Services to Prevent-Mother-to-Child-Transmission (PMTCT) are generally integrated into antenatal care. However, following the birth of their child, MLH must re-engage in HIV care at HIV specialty clinics [3]. During pregnancy, about 99% of pregnant women in Cape Town are tested for HIV [4]. While the rates of uptake of tasks to PMTCT are known, it is difficult to estimate how many new MLH transition to HIV care post birth. In trials evaluating PMTCT the loss to follow-up ranges from 19% to 89% [5]. Loss to follow-up among adults in Johannesburg indicate that 30% are disengaged from care by 12 months [6]. This article summarizes MLH's reports of their re-engagement in care over the 36 months following their children's birth, suggesting the scope of the challenge service providers face in implementing B+ protocols.

---

Corresponding author: Mary Jane Rotheram-Borus, 10920 Wilshire Blvd., Suite 350, Los Angeles, California 90024 USA, Tel: +1 (310) 794-8278, Fax: +1 (310) 794-8297, CCHPublications@mednet.ucla.edu.

[Trial Registration](#). [ClinicalTrials.gov](http://ClinicalTrials.gov) registration # NCT00972699

No Conflicts of Interest are declared.

The data on MLH's re-engagement in care were gathered in the context of a cluster randomized controlled trial (RCT) in 24 township neighborhoods [7]. No intervention effects were observed on any measure of re-engagement in care and, therefore, the RCT is not described in this brief report. Other papers summarize the maternal and child outcomes from this RCT [4,8]. The Institutional Review Boards of University of California Los Angeles, Stellenbosch University, and Emory University approved the study [[ClinicalTrials.gov](http://ClinicalTrials.gov) (NCT00996528)].

Neighborhood clusters (N=24) of 450-600 households were matched outside Cape Town, and 98% of pregnant women were recruited from May 2009 to September 2010 at an average 26 weeks of pregnancy (range, 3–40 weeks).

We assessed 1238 women at baseline. Follow-up assessments were conducted at two weeks post-birth (92%; mean=1.9 weeks; SD=2.1 weeks; median=1.1; range=0.1-14.9); six months (88%; mean=6.2 months, SD=0.7; median=6.0; range=4.2-11.7); 18 months (84%; mean=19.1 months; SD=3.0; median=18.0; range=13.6-34.4); and 36 months (86%; mean = 37.3 months; median =37; SD = 2.2 months; range = 34 to 42 months). All assessments were completed by 72% of mothers.

MLH were on average about 26 years old, with about a 10<sup>th</sup> grade education, and 18.8% were employed [4]. More than half (56%) were living with a partner, although 89% had recent sexual partners. Most mothers lived in informal housing (shacks, 68.7%), 53.3% had water, 55.2% had a toilet, and 89.8% had electricity on their premises. Almost half the mothers (49.5%) and 29.1% of their children had gone hungry in the last week. Most mothers had previously had children (65.9%) and 76.9% had attended an antenatal clinic at the time of recruitment. Among non-primiparous mothers, 15.9% had previously had a low birth weight infant (<2500 grams).

Almost all women had been tested for HIV (99.2%). At recruitment, 26.2% (n=324) of women were HIV seropositive, with 92% learning of their serostatus during this pregnancy at the antenatal clinic. Information regarding HIV status was self-reported and was able to be validated post-birth on the child's government issued Road to Health card which documents maternal HIV status, as well as clinic visits and hospitalizations. Information on linkage and retention in HIV care and ARV utilization was only obtained by maternal self-reports. At completed follow-up assessments, some MLH did not report HIV-related clinic utilization, even though they reported other aspects of their health status and care utilization.

By 36 months post birth, 13 MLH had died, 23 of their children (6.2%) children were seropositive, and an additional 47 mothers identified themselves as HIV seropositive. Almost all MLH (93%; 279/300) had received antenatal care and PMTCT, as previously reported [4]. By six months post-birth 45% of MLH (126/278) had sought care at an HIV clinic and 70.6% of these MLH (89/126) had been prescribed ARV (32% of the overall MLH [89/278]). At 18 months, 52.5% (147/280) had sought HIV care; 89% of these were receiving ARV (46% of the overall MLH [130/280]). At 36 months, 62.5% (n=165/264) were receiving HIV care, with 85% of these receiving ARV (140/165). Only 31% of MLH had been linked to HIV care at the 6, 18, and 36 month assessments (86/280). From 18-36

months, 43% (119/275) had been receiving care. Among surviving MLH, 53% are receiving ARV at 36 months following their children's birth.

These are disturbing data, given the broad accessibility of ARV in the Western Cape of South Africa and the relatively high quality of medical care in the region. B+ regimens may improve re-engagement, but the poor follow-up rates in PMTCT trials [5], the failure to integrate HIV into primary health care clinics in Africa, and HIV stigma (which may be responsible for this study's missing self-reports) must be anticipated.

## Acknowledgments

**Funding:** This study was funded by NIAAA, Grant # 1R01AA017104 and supported by NIH grants MH58107, 5P30AI028697, and UL1TR000124.

## References

1. Luque-Fernandez MA, Van Cutsem G, Goemaere E, Hilderbrand K, Schomaker M, Mantangana N, et al. Effectiveness of patient adherence groups as a model of care for stable patients on antiretroviral therapy in Khayelitsha, Cape Town, South Africa. *PLoS One*. 2013; 8(2):e56088. [PubMed: 23418518]
2. Coutsoydis A, Goga A, Desmond C, Barron P, Black V, Coovadia H. Is option B+ the best choice? *Lancet*. 2013; 14(1):8–10. [PubMed: 23122783]
3. Boule A, Van Cutsem G, Hilderbrand K, Cragg C, Abrahams M, Mathee S, et al. Seven-year experience of a primary care antiretroviral treatment programme in Khayelitsha, South Africa. *AIDS*. 2010; 20(24):563–572. [PubMed: 20057311]
4. le Roux IM, Tomlinson M, Harwood JM, O'Connor MJ, Worthman CM, Mbewu N, et al. Outcomes of home visits for pregnant township mothers and their infants in South Africa: a cluster randomized control trial. *AIDS*. 2013; 27:1461–1471. [PubMed: 23435303]
5. Kalembo FW, Zgambo M. Loss to followup: A major challenge to successful implementation of prevention of mother-to-child transmission of HIV-1 programs in Sub-Saharan Africa. *ISRN AIDS*. 2012; 589817
6. Clouse K, Pettifor AE, Maskew M, et al. Patient retention from HIV diagnosis through one year on antiretroviral therapy at a primary healthcare clinic in Johannesburg, South Africa. *J Acquir Immune Defic Syndr*. 2013; 62(2):e39–e46. [PubMed: 23011400]
7. Rotheram-Borus MJ, Le Roux IM, Tomlinson M, Mbewu N, Comulada WS, le Roux K, et al. Philani Plus (+): A mentor mother community health worker home visiting program to improve maternal and infants' outcomes. *Prev Sci*. 2011; 12(4):372–388. [PubMed: 21850488]
8. Rotheram-Borus MJ, le Roux I, Harwood J, O'Connor M, Worthman C, Tomlinson M. A cluster randomised controlled effectiveness trial evaluating perinatal home visiting among South African mothers/infants. *PLoS One*. 2014; 9:e105934. [PubMed: 25340337]