

Lawrence Berkeley National Laboratory

Recent Work

Title

Capillary Surfaces in Exotic Containers

Permalink

<https://escholarship.org/uc/item/9p00q5hq>

Authors

Concus, P.
Finn, R.

Publication Date

1991-07-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Physics Division


Mathematics Department

Presented at the International Symposium on Hydromechanics and Heat/Mass Transfer in Microgravity, Moscow, USSR, July 6-14, 1991, and to be published in the Proceedings

Capillary Surfaces in Exotic Containers

P. Concus and R. Finn

July 1991



1 LOAN COPY 1
1 Circulates 1
1 for 4 weeks 1

Bldg. 50 Library.

LBL-31200

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. Neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California and shall not be used for advertising or product endorsement purposes.

Lawrence Berkeley Laboratory is an equal opportunity employer.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Capillary Surfaces in Exotic Containers

Paul Concus

**Mathematics Department
Physics Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720**

and

Robert Finn

**Department of Mathematics
Stanford University
Stanford, CA 94305**

July 1991

This work was supported in part by the National Aeronautics and Space Administration under Grant No. NAG3-1143, by the National Science Foundation under Grant No. DMS89-02831, and by the Mathematical Sciences Subprogram of the Office of Energy Research, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

CAPILLARY SURFACES IN EXOTIC CONTAINERS

Paul Concus

*Lawrence Berkeley Laboratory and Department of Mathematics
University of California, Berkeley, CA 94720, U.S.A.*

Robert Finn

*Department of Mathematics
Stanford University, Stanford, CA 94305, U.S.A.*

ABSTRACT

A survey is presented of results to date for capillary surfaces in "exotic" containers. These containers have the property that each one admits a continuum of distinct equilibrium free surfaces, all bounding with the container walls the same volume of fluid, making the same contact angle at the triple interface curve, and having identical mechanical energies. The containers can be so designed that they are themselves axially symmetric but that the fluid configurations of minimizing energy cannot be axially symmetric.





Key words: *capillary surfaces, free surfaces, microgravity, symmetry breaking, space experiments*

The free surface of a liquid that partly fills a container under the action of surface and gravitational forces may assume, in general, one of several possible equilibrium configurations. An example for which only one configuration is possible is a vertical homogeneous cylindrical container of general cross-section, with gravity either absent or directed downward into the liquid; if the boundary of the free surface lies entirely on the cylindrical walls, then the surface is determined uniquely by its contact angle and the liquid volume.^{1,2} Examples of other containers can be given for which there exist two or more distinct equilibrium configurations. Our interest here is in certain container shapes

having the striking property that there is an entire continuum of such configurations.


Axially symmetric containers having this property can be constructed for any prescribed contact angle and Bond number, in such a way that all configurations in the family have identical volumes and yield identical mechanical energies, and so that no two of the free surfaces in the continuum are congruent to each other.^{3,4,5} The procedure requires integration of a system of differential equations, in which the coefficients are determined as solutions of an auxiliary (nonlinear) differential equation system. Nevertheless, the global existence can be proved rigorously. It is shown in Refs. 4 and 6 that the families are all unstable, in the sense that the horizontal flat surface, which all families contain, can be deformed locally into a surface of smaller mechanical energy. It is possible to construct such "exotic" containers in such a way that no symmetric configuration can be energy minimizing. The configurations in Figure 1 show seven surface meridians of the continuum (dashed lines) in the particular case of contact angle 60° , at varying Bond numbers B ; the solid curves are container meridians. The curves in the figures have been normalized so that the containers have maximum radius unity.

Further particular cases are calculated and displayed in Ref. 5. It would be difficult to observe the phenomenon in the earth's gravitational field, as the dimensions of the apparatus would have to be so small as to preclude ac-




XBL 903-989

Figure 1. Meridian of container (solid curve) for contact angle 60° and several Bond numbers showing meridians of some of the symmetric equilibrium solution surfaces (dashed curves), all having the same contact angle and energy, and enclosing the same volume of liquid.



CBB 900-8661



CBB 900-8659

Figures 2a-b. Drop-tower experiment: 2a. (upper) Flat interface initial configuration; 2b. (lower) Non-symmetric terminal configuration.

curate measurements. However, if gravity is absent the equations describing the containers are scale invariant, and experiments can be designed of any size, subject only to the usual laboratory constraints. Even in a microgravity environment that is not strictly zero-gravity, adequately large length scales for accurate observation and measurement are possible. An extended space experiment has been planned and is scheduled to be carried out on the NASA United States Microgravity Laboratory flight (USML-1) in 1992.

Preliminary drop tower experiments have been conducted by M. Weislogel on the five-second drop tower at NASA Lewis Research Center in Cleveland, Ohio.⁷ An exotic container was fabricated corresponding to a contact angle of 80° (the contact angle for the materials used - acrylic plastic and water) and $B = 0$. The container, initially vertical, was filled with a volume of liquid corresponding to a horizontal planar equilibrium free surface meeting the container wall with contact angle 80° , as shown in Figure 2a. This is an equilibrium solution for any gravity level, but for zero g the interface is unstable in this container according to the mathematical theory, as are all members of the symmetric equilibrium family corresponding to the ones depicted in Figure 1. A resulting configuration after a five-second period of free fall subsequent to release of the container in the drop tower is depicted in Figure 2b. The liquid appeared essentially to have settled down to an equilibrium configuration by that time. The free surface exhibits a shape that is obviously non-symmetric, in accordance with the mathematical theory.

Independently, numerical studies were carried out in collaboration with M. Callahan. These studies are based on the Surface Evolver program developed by K. Brakke,⁸ which deforms a given surface in such a way as to decrease (if that is possible) a discrete approximation to the total energy. Three local minima are suggested by the numerical investigation, of which the one with least energy looks remarkably like the surface in Figure 2b observed by Weislogel. It meets the container wall at the join between the bulge and the upper cylindrical wall for a good portion of the circumference, drops rapidly to the join with the lower cylindrical wall, and meets the lower join for the remainder of the circumference. The other locally energy-minimizing solutions are similar in nature, but have two or three excursions from top to bottom of the bulge (and back), instead of just one.⁹

Detailed derivations and further discussion can be found in the references.

This work was supported in part by the National Aeronautics and Space Administration under Grant NAG3-1143, by the National Science Foundation under Grant DMS89-02831,

and by the Mathematical Sciences Subprogram of the Office of Energy Research, U. S. Department of Energy, under Contract Number DE-AC03-76SF00098.

REFERENCES

1. R. Finn, *Equilibrium Capillary Surfaces* (Springer-Verlag, New York 1986), Chapter 5. Russian edition: (Mir Publishers, Moscow 1988).
2. T. I. Vogel, Uniqueness for certain surfaces of prescribed mean curvature, *Pacific J. Math.* **134** (1988) 197-207.
3. R. Gulliver and S. Hildebrandt, Boundary configurations spanning continua of minimal surfaces, *Manuscr. Math.* **54** (1986) 323-347.
4. R. Finn, Nonuniqueness and uniqueness of capillary surfaces, *Manuscr. Math.* **61** (1988) 347-372.
5. P. Concus and R. Finn, Exotic containers for capillary surfaces, *J. Fluid Mech.* **224** (1991) 383-394; Corrigenda, *J. Fluid Mech.* (1991), to appear.
6. P. Concus and R. Finn, Instability of certain capillary surfaces, *Manuscr. Math.* **63** (1989) 209-213.
7. P. Concus, R. Finn, and M. Weislogel, Drop-tower experiments for capillary surfaces in an exotic container, *AIAA J.* (1991), to appear.
8. K. A. Brakke, *Surface Evolver Program*, Geometry Supercomputer Project, University of Minnesota, Minneapolis, MN, 1990.
9. M. Callahan, P. Concus, and R. Finn, Energy minimizing capillary surfaces for exotic containers, in *Computing Optimal Geometries* (with accompanying videotape), J. E. Taylor, ed., AMS Selected Lectures in Mathematics (Amer. Math. Soc., Providence, RI, 1991).

LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
INFORMATION RESOURCES DEPARTMENT
BERKELEY, CALIFORNIA 94720