

UC Irvine

UC Irvine Previously Published Works

Title

Microglial APOE4: more is less and less is more.

Permalink

<https://escholarship.org/uc/item/9rb9g1kj>

Journal

Molecular Neurodegeneration, 18(1)

Authors

Eskandari-Sedighi, Ghazaleh

Blurton-Jones, Mathew

Publication Date

2023-12-19

DOI

10.1186/s13024-023-00693-6

Peer reviewed

EDITORIAL

Open Access



Microglial APOE4: more is less and less is more

Ghazaleh Eskandari-Sedighi^{1,2} and Mathew Blurton-Jones^{1,2,3*}

Abstract

Apolipoprotein E (APOE) is the single greatest genetic risk factor for late onset Alzheimer's disease (AD). Yet, the cell-specific effects of APOE on microglia function have remained unclear. Fortunately, two comprehensive new studies published in the latest issue of *Nature Immunology* have employed complementary gain-of-function and loss-of-function approaches to provide critical new insight into the impact of microglial APOE on AD pathogenesis.

Keywords Apolipoprotein E, Microglia, APOE4, APOE3, Alzheimer's disease, Lgals3, TGFβ

For over three decades, researchers have known that polymorphisms in Apolipoprotein E (APOE) represent the single greatest genetic risk factor for late-onset Alzheimer's Disease (AD) [3]. Since then, the APOE4 risk allele has been shown to influence an array of diseased-associated processes including amyloid aggregation and clearance, tau-induced neurodegeneration, glucose metabolism, synaptic degeneration, and cerebral amyloid angiopathy (reviewed in detail in [9, 4], and [1]). Yet, the precise impact of APOE4 on microglial function has remained unclear. As a member of the apolipoprotein family of lipid binding proteins, human APOE is encoded by three differing alleles: ε2, ε3 and ε4. In comparison to the most common allele, ε3, a single copy of ε4 is associated with a threefold increased risk of developing AD, whereas two copies of ε4 leads to a remarkable 15-fold increase in disease risk [3]. In contrast, the ε2 allele confers a protective ~sevenfold decreased risk of AD. The

apoE protein has important roles in lipid homeostasis including the shuttling of cholesterol and lipids between cells. Each isoform also exhibits distinct binding affinities for different classes of lipoproteins. While astrocytes are the predominant producer of APOE within the healthy brain [10], microglia can significantly upregulate APOE expression in response to inflammatory insults, including AD pathology [5, 6, 8]. Understanding cell-specific roles of apoE isoforms within the brain will therefore provide invaluable new mechanistic insight that could likely inform the development of more effective therapeutics.

Fortunately, two highly complementary and comprehensive studies from the labs of Guojun Bu [7] and Oleg Butovsky [11], have now greatly advanced our understanding of the effects of APOE polymorphisms on microglial function. Led by co-first authors Liu and Wang, Yin and Rosenzweig, and published in *Nature Immunology*, each study applied opposing, yet highly complementary approaches to deconvolve the role of microglial APOE in AD pathogenesis. Using a Cx3cr1^{creERT2/+} mouse line to drive inducible expression of human APOE3 or APOE4 on a murine ApoE-knockout background, Liu et al. created a brain environment in which apoE3 or apoE4 could be specifically and exclusively expressed by microglia and CNS-associated macrophages (CAMs), also referred to as border-associated macrophages (BAMs) [7]. In contrast, Yin et al. crossed Cx3cr1^{creERT2/+} mice with mice expressing

*Correspondence:

Mathew Blurton-Jones
mblurton@uci.edu

¹ Institute for Memory Impairments and Neurological Disorders, University of California Irvine, Irvine, CA 92697, USA

² Sue and Bill Gross Stem Cell Research Center, University of California Irvine, Irvine, CA 92697, USA

³ Department of Neurobiology & Behavior, University of California Irvine, Irvine, CA 92697, USA



This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

floxed versions of human APOE3 or APOE4, providing a paradigm to specifically delete APOE3 or APOE4 only within microglia and BAMs, while preserving human APOE expression in other cell types [11]. Both studies then crossed these inducible models with APP/PS1 mice to examine the impact of microglial-specific APOE4 or APOE3 expression [7] or deletion [11] on the microglial response to AD pathology. Despite their differing models and experimental designs, both groups reach the same ultimate conclusion: expression of human apoE4 restricts microglia in a more quiescent state, impairing their ability to mount a protective response to AD-associated pathology and increasing neurodegeneration (Fig. 1).

Having validated the specificity of their model system, Liu et al. began by inducing microglial/CAM-specific expression of either apoE3 or apoE4 in 6-month-old APP/PS1 mice [7]. Three months later, they found that microglial apoE3 expression led to an increased number of plaque-associated microglia while concurrently reducing insoluble amyloid levels, dystrophic neurites, and microglial lipid droplets. Using single cell sequencing they further demonstrated that apoE3 induced a more reactive transcriptomic signature, promoting expression of neurodegenerative microglial (MGnD) transcripts. In

contrast, microglial-specific expression of apoE4 produced opposing effects, worsening each of these pathological endpoints, impairing lipid metabolism, and promoting a stress-related microglial transcriptomic signature. Further analysis revealed apoE4-mediated induction of eIF2 signaling, oxidative phosphorylation and mitochondrial dysfunction in APP/PS1 mice, as well as downregulation of genes associated with the complement pathways, and lysosomal degradation.

Next, Liu and colleagues thoroughly assessed the cell-autonomous and cell-non-autonomous effects of microglial apoE on astrogliosis. Accordingly, they observed that in the absence of endogenous mouse Apoe, microglial expression of apoE3 reduced A β -associated cortical astrogliosis whereas apoE4 promoted astrocytic activation. While GFAP and A β levels exhibited a strong positive correlation in mice expressing microglial apoE3, no correlation was observed in apoE4 expressing mice, suggesting that microglial apoE4 can influence astrogliosis in an A β -independent manner. To further understand the potential impacts of Apoe expression in other cells they combined their inducible system with 5xFAD mice that express endogenous murine Apoe. In this model, the number of GFAP-positive astrocytes




Fig. 1 A schematic representation of the complementary gain-of-function and loss-of-function approaches used to examine the impact of human APOE on microglial function in AD. APOE4^{+/+} restricts microglia in a non-responsive state (left part of the circle), resulting in exacerbated amyloid pathology (red) and dystrophic neurites (green). In contrast, deletion of APOE4 in microglia rescues these deficits (right part of the circle)

was again increased by microglial expression of apoE4, but no changes in GFAP were observed in 5xFAD mice expressing microglial apoE3 on a murine apoE expressing background. Taken together, the authors concluded that these differences are likely due to the cell-autonomous and non-cell-autonomous effects of apoE as well as the development of diffused versus compact amyloid plaques. Importantly, microglial apoE4 also influenced non-amyloid-dependent functions, reducing microglial response to a localized laser ablation, impairing cognition, and disrupting long term potentiation in comparison to apoE3 expressing microglia.

Taking an opposite approach, Yin and colleagues began their study by investigating the effects of global humanized apoE3 and apoE4 expression, finding that microglia isolated from apoE4 knockin mice exhibited increased expression of homeostatic genes in comparison to apoE3 mice [11]. When challenged by intracerebral injection of apoptotic neurons, apoE3 microglia mounted a robust phagocytic response, upregulating MGnD, interferon-responsive, and antigen presentation-associated transcripts, while apoE4 microglia failed to induce these disease-responsive genes. Furthermore, microglial-targeted deletion of apoE4 rescued the response to apoptotic neurons, elevating expression of MGnD genes while suppressing homeostatic transcripts.

To examine the effects of microglial-specific apoE deletion in the context of AD, Yin et al. also employed the APP/PS1 model as well as the P301S model of tauopathy [11]. Consistent with their apoptotic neuron results, microglia isolated from 9-month-old P301S:APOE3-KI mice exhibited an induction of several MGnD genes in response to tau pathology and a corresponding downregulation of homeostatic transcripts including *Tgfb1* and *Smad3*. In contrast, P301S:APOE4-KI microglia exhibited a blunted MGnD response and increased TGF β -signaling. Importantly, deletion of microglial APOE4 reversed many of these transcriptional deficiencies, reduced tau hyperphosphorylation and prevented cortical neuronal loss. Turning to the APP/PS1 model, the group found that APP/PS1:APOE3-KI mice readily adopt an MGnD signature, expressing high levels of *Lgals3* and *Clec7a*, whereas apoE4-expressing microglia again exhibited an impaired MGnD response mediated via upregulation of microglial homeostatic checkpoint genes, including *ITGB8* and *Inpp5d*. Microglial-specific deletion of APOE4 in turn restored the MGnD signature, increased plaque associated microglia and plaque compaction, and reduced both A β pathology and dystrophic neurites. They also reported an apoE4-mediated induction of *ITGB8*-transforming growth factor- β (TGF β) signaling within microglia that drove upregulation of

homeostatic checkpoint genes, reducing the MGnD phenotype.

Given the role of astrocyte derived *ITGB8* in TGF β activation and Butovsky's prior work showing the importance of TGF β in microglial homeostasis [2], Yin and colleagues next examined the effects of manipulating *ITGB8* expression. Using both knockout approaches and intracerebral injection of *ITGB8* neutralizing antibody, they found that reduction of *ITGB8* signaling led to enhanced Ab phagocytosis, elevated microglial expression of antigen presentation and interferon genes, increased GFAP and *Clec7a*, and decreased plaque load. Taken together, they concluded that the MGnD state is regulated by the reciprocal induction of APOE signaling and suppression of TGF β and introduce the microglial APOE4–*ITGB8*–TGF β pathway as a negative regulator of the neuroprotective microglial responses to AD pathology.

A mutual discovery of both studies is the convergence of microglial apoE-driven effects on *Lgals3*, the gene encoding galectin-3, a lectin with immunoregulatory functions. This encouraged both groups to examine the impact of microglial apoE on astrocytes. Using GFAP immunostaining, Liu et al. observed that microglial expression of apoE3 reduced A β -associated astrogliosis, whereas apoE4 promoted A β -independent astrocytic activation. However, Yin et al. reported decreased astrocytic activation in the presence of microglial apoE4. These differences could be due to distinct approaches each group used to quantify astrocyte activation as well as the differing apoE background of their mouse models. In Liu et al. the mice had no expression of apoE within astrocytes and the amyloid deposits were diffuse in nature. In contrast, the model used by Yin et al. included astrocytic expression of human apoE and exhibited compact amyloid plaques. Differences in plaque compaction can in turn effect the astrocytic response. Regardless of this minor difference, both studies support a critical role for microglial apoE in modulating microglia-astrocyte crosstalk that converges on microglial-derived *Lgals3* signaling.

Lastly and importantly, both groups further validated their observations in postmortem human AD brain samples; Liu et al. investigated the plaque-associated microglia and observed a significant reduction in *Lgals3*-positive responsive microglia in *APOE4* carriers compared to *APOE3*. Yin et al. used publicly available RNAseq datasets and reported lower levels of MGnD-associated genes as well as increased TGF β signaling in *APOE4* carriers, further supporting their hypothesis that *ITGB8*–TGF β signaling could influence the development of AD by inhibiting MGnD induction. Taken together, these back-to-back studies used complementary gain-of-function and loss-of-function approaches to provide

critical new insight into the role of microglial apoE in AD, ultimately finding that when it comes to microglial APOE4, more is less and less is more.

Abbreviations

APOE	Apolipoprotein E
AD	Alzheimer's disease
CAMs	CNS-associated macrophages
BAMs	Border-associated macrophages
KI	Knockin
TGFβ	ITGB8-transforming growth factor-β
MGNd	Microglial neurodegenerative phenotype

Authors' contributions

G.E.S. and M.B.J. drafted the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by NIH U19-AG069701 and Cure Alzheimer's Fund to M.B.J.

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 28 November 2023 Accepted: 4 December 2023

Published online: 19 December 2023

References

1. Bu G. APOE targeting strategy in Alzheimer's disease: lessons learned from protective variants. *Mol Neurodegeneration*. 2022;17:51.
2. Butovsky O, Jedrychowski M, Moore C, et al. Identification of a unique TGF-β-dependent molecular and functional signature in microglia. *Nat Neuroscience*. 2014;17:131–43.
3. Corder EH, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*. 1993;261(5123):921–3.
4. Fernández-Calle R, Konings SC, Frontiñán-Rubio J, et al. APOE in the bullseye of neurodegenerative diseases: impact of the APOE genotype in Alzheimer's disease pathology and brain diseases. *Mol Neurodegeneration*. 2022;17:62.
5. Keren-Shaul H, et al. A unique microglia type associated with restricting development of Alzheimer's disease. *Cell*. 2017;169(7):1276–1290. e1217.
6. Krasemann S, et al. The TREM2-APOE pathway drives the transcriptional phenotype of dysfunctional microglia in neurodegenerative diseases. *Immunity*. 2017;47(3):566–581. e569.
7. Liu CC, Wang N, Chen Y, et al. Cell-autonomous effects of APOE4 in restricting microglial response in brain homeostasis and Alzheimer's disease. *Nat Immunol*. 2023;24:1854–66.
8. Mathys H, et al. Single-cell transcriptomic analysis of Alzheimer's disease. *Nature*. 2019;570(7761):332–7.
9. Raulin AC, Doss SV, Trotter ZA, et al. ApoE in Alzheimer's disease: pathophysiology and therapeutic strategies. *Mol Neurodegeneration*. 2022;17:72.

10. Xu Q, et al. Profile and regulation of apolipoprotein E (ApoE) expression in the CNS in mice with targeting of green fluorescent protein gene to the ApoE locus. *J Neurosci*. 2006;26(19):4985–94.
11. Yin Z, Rosenzweig N, Kleemann KL, et al. APOE4 impairs the microglial response in Alzheimer's disease by inducing TGFβ-mediated checkpoints. *Nat Immunol*. 2023;24:1839–53.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

