

UCSF

UC San Francisco Previously Published Works

Title

Hospital Culture and Intensity of End-of-Life Care at 3 Academic Medical Centers

Permalink

<https://escholarship.org/uc/item/9s01s3hn>

Journal

JAMA Internal Medicine, 183(8)

ISSN

2168-6106

Authors

Dzeng, Elizabeth

Batten, Jason N

Dohan, Daniel

et al.

Publication Date

2023-08-01

DOI

10.1001/jamainternmed.2023.2450

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Hospital Culture and Intensity of End-of-Life Care at Three Academic Medical Centers: A Comparative Ethnographic Study

Elizabeth Dzeng, MD, PhD, MPH^{1,2,3}, Jason N. Batten, MD, MA^{4,5}, Daniel Dohan, PhD², Jacob Blythe, MD, MA⁶, Christine Ritchie, MD, MSPH^{7,8}, J. Randall Curtis, MD, MPH^{9,10}

¹ Division of Hospital Medicine, Department of Medicine, University of California, San Francisco, San Francisco, California, USA.

² Philip R. Lee Institute for Health Policy Studies, University of California, San Francisco, San Francisco, California, USA.

³ Cicely Saunders institute, King's College London, London, United Kingdom.

⁴ Department of Anesthesia, Perioperative, and Pain Medicine, Stanford University, Stanford, California, USA.

⁵ Stanford Center for Biomedical Ethics, Stanford University, Stanford, California, USA.

⁶ Department of Radiology, Massachusetts General Hospital, Boston, Massachusetts, USA.

⁷ Division of Palliative Care and Geriatric Medicine, Department of Medicine, Massachusetts General Hospital, Boston, Massachusetts, USA.

⁸ Mongan Institute Center for Aging and Serious Illness, Department of Medicine, Massachusetts General Hospital, Boston, Massachusetts, USA.

⁹ Division of Pulmonary and Critical Care Medicine, Department of Medicine, University of Washington, Seattle, Washington, USA.

¹⁰ Cambia Palliative Care Center of Excellence, University of Washington, Seattle, Washington, USA.

Corresponding Author:

Elizabeth Dzeng, MD, PhD, MPH
UCSF Division of Hospital Medicine
521 Parnassus Avenue, 5th floor, Box 0131
San Francisco, CA, 94143
(410) 207-9923
liz.dzeng@ucsf.edu

Word count: 2992

41 **KEY POINTS:**

42 **Question:** What is the role of hospital culture and institutional structures in the provision of
43 potentially non-beneficial high-intensity life-sustaining treatments near the end of life?

44 **Finding:** We conducted 113 semi-structured, in-depth interviews with hospital-based clinicians
45 and administrators at three academic hospitals in California and Washington. Hospital culture
46 tended to be aligned with institutional structures (e.g., policies, practices, protocols, and
47 resources), and together shaped the provision of potentially non-beneficial life-sustaining
48 treatments near the end-of-life at each site.

49 **Meaning:** Institutional cultures should be considered when developing policies and
50 interventions to mitigate non-beneficial high-intensity life-sustaining treatments.

51
52

ABSTRACT:

Importance: There is significant institutional variability in the intensity of end-of-life care that is not explained by patient preferences. Hospital culture and institutional structures (e.g., policies, practices, protocols, resources), might contribute to potentially non-beneficial high-intensity life-sustaining treatments near the end of life.

Objective: To understand the role of hospital culture in the everyday dynamics of high-intensity end-of-life care.

Design: Comparative ethnographic study. Data were deductively and inductively analyzed using thematic analysis through an iterative coding process.

Setting: Three academic hospitals in California and Washington that differed in end-of-life care intensity based on measures in the Dartmouth Atlas.

Participants: Hospital-based clinicians, administrators, and leaders

Main Outcome and Measure: Institution-specific policies, practices, protocols, and resources that shape hospital culture and their role in the everyday dynamics of potentially non-beneficial high-intensity life-sustaining treatments.

Results: We conducted 113 semi-structured, in-depth interviews with inpatient-based clinicians and administrators between December, 2018 and June, 2022. Respondents at all hospitals described default tendencies to provide high-intensity treatments that they believed was universal in American hospitals. They also reported that pro-active, concerted efforts among

multiple care teams were required to de-escalate high-intensity treatments. Efforts to de-escalate were vulnerable to being undermined at multiple points during a patient's care trajectory by any individual or entity. Respondents described institution-specific policies, practices, protocols, and resources that engendered broadly-held understandings of the importance of de-escalating non-beneficial life-sustaining treatments. Respondents at different hospitals reported different policies and practices that encouraged or discouraged de-escalation. They described how these institutional structures contributed to the culture and everyday dynamics of end-of-life care at their institution.

Conclusions and Relevance: Clinicians, administrators, and leaders at the hospitals we studied report that they work in a hospital culture where high-intensity end-of-life care constitutes a default trajectory. Institutional structures and hospital cultures shape the everyday dynamics by which clinicians may de-escalate end-of-life patients from this trajectory. Individual behaviors or interactions may fail to mitigate potentially non-beneficial high-intensity life-sustaining treatments if extant hospital culture or lack of supportive policies and practices undermine individual efforts. Hospital cultures need to be considered when developing policies and interventions to decrease potentially non-beneficial high-intensity life-sustaining treatments.

BACKGROUND

Ethical challenges are common around the provision of potentially non-beneficial high-intensity life-sustaining treatments near the end of life such as intensive care unit (ICU) admission, mechanical ventilation, cardiopulmonary resuscitation, and continuous renal replacement therapy.^{1–3} The United States is unique in the proportion of older adults with serious illness who are admitted to the ICU near the end of life^{4–8} despite minimal evidence that the benefits outweigh the burdens^{9–11}. Many of these patients receive treatments that are perceived to be non-beneficial, the majority of whom do not survive hospitalization^{12–14}. The US's market-based healthcare system encourages overtreatment¹⁵. Indeed, ICU beds per capita and the use of high-intensity life-sustaining treatments continue to rise^{8,16–21}. Interventions aimed at improving the quality of end-of-life care in the ICU have had mixed results^{22–25}.

Understanding how and why potentially non-beneficial high-intensity life-sustaining treatments occur is complicated by significant variation in end-of-life treatment intensity throughout the US^{26–31}. This observed variation is not explained by patient preferences, regional differences, or a clinicians' failure to recognize poor prognoses^{2,5,22,28,29,32–35}. This suggests that potentially non-beneficial high-intensity life-sustaining treatments may be driven by currently understudied systemic factors.

Drivers of potentially non-beneficial high-intensity life-sustaining treatments include the ethical, social, and cultural aspects of healthcare institutions^{27,31,36–39}. Culture is defined as shared beliefs, values, and practices of a group of people (e.g., organization, institution, or profession) and influences the typical way of doing things^{40,41}. Institutions vary in treatment intensity, which appears to be stable over time^{32,42}. Prior studies suggest that hospital cultures might influence the provision of non-beneficial high-intensity life-sustaining treatments^{27,31,36,38,43}. There is a gap in our understanding of how hospital cultures are related to institution-specific structures –

defined as the hospital's policies, practices, protocols, and resource allocation – as well as how it might influence the intensity of end-of-life care. Institutional structures and hospital culture influence one another cyclically and iteratively⁴⁴. In sociological scholarship, this feedback loop perpetuated over time is described as “recursion,” where one entity cyclically reproduces and strengthen another. The objective of this study was to elucidate our understanding of the complex, recursive relationships between hospital culture, institutional structures, and the provision of potentially non-beneficial high-intensity life-sustaining treatments.

METHODS

Design: This is a comparative ethnographic study conducted at three academic hospitals in California and Washington selected for differences in intensities of end-of-lifecare (e.g., high, medium, and low) based on the Dartmouth Atlas (See Table 1). We interviewed clinicians and administrators with different clinical backgrounds and organizational responsibilities. This project was guided by a conceptual framework based upon literature review and on prior work by the authors (See Figure 1)^{31,45–47}. This conceptual framework was continuously refined throughout the project as more data was collected and analyzed.

Data collection: One interviewer (ED), a hospitalist and PhD-trained sociologist, conducted semi-structured, in-depth interviews with inpatient-based clinicians, leaders, and administrators. Interviews were conducted in-person until the COVID-19 pandemic when interviews transitioned to Zoom video-conferencing. Respondents were purposively sampled by profession, which contributed to understanding everyday dynamics within each hospital. Recruitment occurred through group e-mail solicitations, individual requests, and snowball sampling^{48,49}. Participation rate could not be calculated because persons were contacted through unsolicited e-mails and list-serves, along with direct solicitations. Sampling occurred until theoretical saturation was

reached, the point where interviews generated no new insights.^{50,51} We include further details of our methods using the Consolidated Criteria for Reporting Qualitative Research (COREQ) in eTable 1.

An interview guide was used across all sites; minor adaptations were made for different roles (See eAppendix 1 for interview guide). The initial guide derived from our conceptual framework and evolved during pilot and subsequent interviews. The interviews were open-ended; participants were encouraged to explore topics they considered relevant. Interviews were audiotaped, transcribed, and anonymized.

Analysis: Data collection and analysis occurred concurrently. During data collection, emerging findings were incorporated into ongoing interviews for further exploration to determine their consistency, robustness, and salience. We continually self-reflecting around our own lens (e.g., perspectives, assumptions, positionality) and how that might influence the research process and interpretation. We paid careful attention to counterfactual data, which were used to inform ongoing data collection, refine emerging themes, and enhance rigor. Respondents' diverse perspectives provide a composite view of clinical care at each hospital. Though no one respondent was able to fully explain or recognize what is attributable to hospital culture, analysis of the corpus of interviews provided these broader insights⁵².

The research team (ED, DD, JRC, JNB, TM) thematically coded a subset of interviews to create an initial codebook.^{53–55} We deductively and inductively generated codes through line-by-line analysis and discussion and came to consensus on coding definitions.^{55,56} 20% of the interviews were subsequently double-coded by TM, LP, JNB, JB, and CB, which contributed to further refinement of the codebook. Analysis was conducted using ATLAS.ti software. Disagreements were resolved through discussion and clarification of code definitions until consensus was

achieved. Member-checking occurred through presentations, discussions, and review of manuscript drafts with clinicians of similar background to the respondents. Additional interviews were conducted following the initial drafts of this manuscript to finalize evolving hypotheses.

To characterize organizational similarities and differences among the three hospitals, we examined thematic divergences and convergences among individual respondents within and across each institution. The UCSF Institutional Review Board approved this study and participants underwent a written or verbal consent process. Our IRB protocol called for anonymization of respondents and hospitals.

RESULTS

Table 2 describes the demographic characteristics of the 113 interviewees (66 in person, 47 via video-conferencing). The mean and median interview length was 47 minutes. During analyses, we noted similarities between experiences at low and medium-intensity hospitals as distinct from the high-intensity hospital. As such, we characterized the study sites as high-intensity and “lower-intensity”, which includes the low and medium-intensity hospitals.

Defaults of high-intensity care and consistency between Dartmouth Atlas-measured care intensity and respondents’ accounts of hospital culture

Potentially non-beneficial high-intensity life-sustaining treatments occurred at all hospitals (Table 2, Quote 1-3) as well as defaults towards high-intensity care (Q4-6). Respondents noted that this default reflected cultural norms in American society (Q7). However, respondents at each of the three sites also described distinct hospital cultures around the intensity of end-of-life care that differed between sites and were broadly consistent with the intensity indicated by the Dartmouth Atlas data. High-intensity care was particularly notable at the high-intensity hospital

(Q3) as was the receptiveness towards palliative care and a mindset towards de-escalation at the lower-intensity hospitals (Q8).

Consensus and coordination required to de-escalate

In the absence of patient or surrogate preferences, respondents at all sites defaulted towards ICU admission if the patient met criteria, regardless of whether it was beneficial. At the high-intensity hospital, this led to persistent escalation whereas at lower-intensity hospitals, there were multi-disciplinary efforts to engage in shared decision-making within the ICU setting to de-escalate (Q9, 10). Institutional structures and hospital cultures at the lower-intensity hospitals facilitated de-escalation; some respondents perceived it as relatively easy to de-escalate life-sustaining treatments (Q11). Respondents at the lower-intensity hospitals but not at the high-intensity hospital felt comfortable not offering or strongly recommending against non-beneficial treatments including dialysis, pressors, and feeding tubes (Q12).

At all hospitals, consensus and coordination was required amongst clinicians and families to *de-escalate* life-sustaining treatments (Q13). At the high-intensity hospital, respondents described consensus around de-escalation to be challenging to achieve (Q14). Respondents at the lower-intensity hospitals described alignment and teamwork amongst and within care teams to achieve consensus to de-escalate and present a unified message to families. Potentially non-beneficial high-intensity treatments at lower-intensity hospitals were discussed and sometimes resisted (Q15-16). In contrast, consensus was not always required to *escalate* life-sustaining treatments (Q17).

Cultural norms around crucial decision points

At the lower-intensity hospitals, respondents noted that because multiple people were involved in decision-making, there were multiple checks to ensure that treatments provided were

beneficial. . Respondents described a cultural norm that encouraged a shared desire towards appropriate de-escalation (Q18-19). They recognized that defaults existed towards high-intensity life-sustaining treatments and described intentional mindsets and actions to resist which required time and effort (Q9). At the high-intensity hospital, respondents noted that the involvement of multiple people in decision-making led to a diffusion of decision-making responsibility that allowed for unchecked momentum towards a high-intensity care (Q20).

Efforts to de-escalate non-beneficial high-intensity life-sustaining treatments can be undermined

Despite efforts to de-escalate, respondents described undermined attempts at de-escalation of non-beneficial high-intensity life-sustaining treatments at all hospitals by external entities such as consultants, ethics committees, or hospital administrators (Q21-23). In our respondent sample, however, the theme of undermined de-escalation was particularly notable at high-intensity hospitals (Q21-22). Transitions of care such as attending physician turnover were described as occasions where re-escalation of high-intensity treatments occurred (Q24).

The ways that ethics committees made decisions and supported clinical teams appeared to be aligned with measured end-of-lifecare intensity. At the high-intensity hospital, respondents described the ethics committee's approach as favoring patient autonomy and placing additional burdens on clinical teams seeking to de-escalate (Q25). At the lower-intensity hospitals, respondents reported that the ethics committees worked with clinicians and institutional leadership to support clinicians' clinical judgement (Q26). At one of the lower-intensity hospitals, respondents reported that risk management encouraged clinicians to make decisions based on the patient's best interest and supported them institutionally and legally to do so (Q27).

Institutional policies, protocols, practices, and resources shape hospital culture

At the lower-intensity hospitals, protocolized approaches and hospital policies counteracted momentum towards potentially non-beneficial high-intensity life-sustaining treatments. Respondents felt that institutional structures (e.g., policies, protocols, practices, resources) successfully created and sustained hospital cultures that supported goal-concordant end-of-life care at these hospitals (Table 3, Q28-29). Respondents felt that palliative care services demonstrated their value as important and trusted sources of support and education not only for patients and families, but also for clinicians (Q30). At the lower-intensity hospitals, respondents described consultants (Q16), social services (Q9), hospital leadership (Q31-32), ethics committees (Q9, 26), risk management (Q27), and other entities as facilitating consensus rather than undermining de-escalation efforts. These institutional entities actively encouraged clinicians to de-escalate when ethically appropriate and deferred to clinician judgement (Q32).

The relational and emotional aspects of end-of-life care

There was a notable emotional toll, especially at the high-intensity hospital, associated with challenging cases which affected respondents' future willingness to attempt de-escalation. Respondents described feeling powerlessness, particularly when institutional entities thwarted attempts at de-escalation and asked them to provide potentially non-beneficial treatments. This led to beliefs that their efforts were not worth the fight (Q33-34). Although emotionally fraught and challenging cases that involved intense conflict were relatively infrequent, the specter of these conflicts loomed large in a clinicians' minds long afterwards. Respondents noted that the negative emotional valence surrounding prior efforts resulted in less willingness to expend effort to de-escalate care in the future.

One theme that emerged only at the lower intensity hospitals was the way end-of-life decision-making occurred. Respondents at these hospitals reported that they achieved de-escalation by seeking nuance between the extremes, finding ethical middle grounds between providing all

possible treatments and unilaterally not offering non-beneficial high-intensity life-sustaining treatments. Respondents described focusing on relationship building and aligning with the family. They recognized the potentially traumatic and burdensome nature of these decisions. Although an informed assent approach, where clinicians explicitly offer the choice to defer to clinicians' judgement about withdrawing or withholding life-sustaining treatments, was an accepted norm at the lower-intensity hospitals, it was used within a context of empathetic, values-oriented goals of care discussions. Respondents frequently noted that this approach sometimes took more time but was the right thing to do (Q35-36).

DISCUSSION

At the three hospitals we studied, hospital culture appeared to recursively shape and reflect institutional structure, as manifested by its policies, practices, protocols, and resource allocation. Each hospital's culture and institutional structure either supported or undermined attempts to de-escalate against the default of high-intensity life-sustaining treatments in American medicine. A clinicians' attempt to de-escalate appeared to be more effective if they operated within a hospital culture that discouraged high-intensity life-sustaining treatments. The aggregate effects of a hospital's culture and institutional structures appeared to coalesce into an institutionally specific equilibrium which create and reproduce hospital culture. Overall, these observed patterns were consistent with our conceptual model (Figure 1) with regards to how hospital culture and institutional structures recursively reproduce and strengthen one another.

In Figure 2, we illustrate a prototypical patient trajectory of how recursive reproduction and strengthening between institutional structures and hospital cultures might impact the ease of de-escalation. A default towards high intensity treatments occurred unless every element of a hospital's culture, as manifested by specific structural and procedural factors (i.e., institutional structures), were aligned to resist this default. We described in our results instances where

hospital leaders, administrators, subspecialty consultants, and ethics committees undermined de-escalation. As such, successful de-escalation of non-beneficial high-intensity life-sustaining treatments required every individual involved to be aligned towards de-escalation. If one individual undermined an attempted de-escalation, respondents reported that the care trajectory regressed to the default of high-intensity treatments.

It is well known that the US is an outlier in its default towards high-intensity treatments^{57,58}. Respondents at all sites needed to employ active and concerted efforts to de-escalate non-beneficial high-intensity life-sustaining treatments. The tendency across hospitals was to escalate treatments unless there was explicit agreement with all relevant decision-makers to de-escalate. A relevant concept is that of clinical momentum, or clinical practice norms and patterns of usual care that promote the accumulation of multiple interventions over time⁵⁹. Even when clinicians, patients, or families resisted this default towards high-intensity care, multiple overlapping institutional structures (e.g., clinical practices of the primary/consultant, hospital policies) made it challenging to do so.

It was notable that dynamics that encouraged escalation at the high-intensity institution were similar to dynamics that facilitated de-escalation at lower-intensity hospitals. As we described in our results (e.g., Q20), respondents at the high-intensity hospital described a diffusion of decision-making responsibility to multiple clinicians as encouraging escalation. Respondents at lower-intensity institutions described (e.g., Q16,18) similar involvements multiple decision-makers and teams as positive checks and balances which facilitated multiple opportunities to raise concerns. Another example of similar dynamics resulting in divergent clinical practice patterns occurred when respondents at the high-intensity hospital (e.g., Q22) described consultants as undermining de-escalation, whereas consultants helped mitigate unchecked clinical momentum at the lower-intensity hospitals (e.g., Q15). Studies examining clinical

practice patterns often focus on team dynamics and inter-team structures^{60,61}. Our results suggest that hospital culture should also be considered. Hospital culture manifests in many ways, such as the convergence of ethical perspectives around appropriate care amongst all members of the multi-disciplinary team, or a preponderance of clinicians willing to attempt de-escalation.

These findings provide insights into the recursive relationship between institutional policies and practices designed to mitigate potentially non-beneficial high-intensity life-sustaining treatments and institutional culture by creating feedback loops. Deliberate hospital policies and protocols that encourage thoughtful pauses around treatment escalation decisions help mitigate unchecked clinical momentum. The “comfort care huddle” described at one of the lower-intensity hospitals allows for all members of the team to convene at regular intervals to discuss treatment de-escalation opportunities. This is reminiscent of a protocolized process in France (“limitation et arrêt des traitements”) which we had previously described as beneficial in their ability to align team members towards a unified message with family⁴⁶. The ways these interventions, policies, and practices cyclically reinforce and are reinforced by hospital culture to influence care intensity is an example of the recursiveness of social processes.

Though extreme and rare, our results demonstrate that challenging clinical cases had an outsized impact on clinicians’ perceptions and willingness to engage in future difficult cases. This is another example of recursion, but in this case between the individual and the hospital’s culture. Shared beliefs and experiences recursively reproduce cultural orientations to strengthen hospital culture. This “feedback – feedforward” reproductive process is a general feature of social life as noted by Antony Giddens’ Structuration Theory⁴⁴ whereby individuals’ experiences and subsequent actions are influenced by institutional constraints in ways that reinforce hospital culture.

Limitations

This study is limited in its ability to ascribe causation from hospital cultures to care intensity. While this study helps elucidate the dynamics between hospital culture, institutional structures, and care intensity, there may be other differences between hospitals such as uptake of palliative care, economic incentives, and patient population demographics that influence care intensity. While broader macro-sociological, political, and economic forces influence individuals and institutions, we were not able to specifically examine those phenomena in this manuscript beyond a general perception by respondents that these forces contributed to a default of high-intensity care. Our observations were based on three urban academic medical centers and may not generalize to community hospitals.

Conclusion

We describe the significance of hospital culture and institutional structures in resisting the default towards high-intensity life-sustaining treatments. This study highlights the importance of the deliberate design of institutional structures – policies, practices, protocols, and resource allocation – to mitigate the harmful impact of entrenched societal forces and defaults within the American healthcare system. These efforts should include purposeful consideration of how institutional values might reflect and be reinforced by specific policies and procedures, such as ethics committee decision-making processes and the structure of co-management and consultation services. We also suggest careful attention to the relational aspects of care including risk management values, thoughtful and consistent dialogue between institutional leaders and clinical teams, and attention to the nature of administrator oversight in relation to high-stakes clinical decisions in extreme cases. While these entities' values and structures may appear to be relatively distant to day-to-day clinical decision-making, their impact can have potential wide-ranging intended and unintended consequences.

382

383 This study illustrates how hospital culture might undermine the impact of interventions that
384 narrowly target individuals or groups of individuals (e.g., team dynamics, communications,
385 decision-making). Consideration of hospital culture and its impact on individuals and clinical
386 practice patterns should be incorporated into institutional policies, practices, and interventions.
387 The design of institutional policies, protocols, practices, and resource allocation have the
388 potential to be a powerful shaper of hospital culture and, thereby, individual clinician behavior
389 and patient and family experiences.

390

391 **Acknowledgements:** The authors would like to thank Catherine Burke, Steven Chang,
392 Solomon Liao, Thea Matthews, Lorraine Pereira, Amy Rosenwohl-Mack, Neil Wenger, and
393 Sherry Xu for their assistance with the project. We would also like to thank the study
394 respondents who took the time to provide us with their insights. This manuscript is dedicated to
395 the memory of J. Randall Curtis, and is a culmination of a decade of mentorship, support, and
396 enthusiasm from this most beloved mentor and friend (ED).

397

398 **Funding:** Research reported in this manuscript was supported by an Alzheimer's Disease
399 Research Award from the California Department of Public Health (ED, DD), the National
400 Palliative Care Research Center (NPCRC) Junior Investigator Career Development Award (ED),
401 the National Institute on Aging under Award Number 1R03AG060098 (ED), the National
402 Institute of Nursing Research under Award Number 3R01NR018161-02S1 (ED, JRC), and the
403 National Center for Advancing Translational Sciences of the NIH under Award Number
404 KL2TR001870-02 (ED). Its contents are solely the responsibility of the authors and do not
405 necessarily represent the official views of the NIH.

406

407 **REFERENCES**

- 408 1. Institute of Medicine. *Dying in America Improving Quality and Honoring Individual*
409 *Preferences Near the End of Life*. (2014).
- 410 2. Turnbull, A. E., Bosslet, G. T. & Kross, E. K. Aligning use of intensive care with patient
411 values in the USA: past, present, and future. *Lancet Respir. 7*, 626–638 (2019).
- 412 3. Khandelwal, N. *et al.* Pragmatic methods to avoid intensive care unit admission when it
413 does not align with patient and family goals. *Lancet Respir. 7*, 613–625 (2019).
- 414 4. Gomes, B., Calanzani, N., Gysels, M., Hall, S. & Higginson, I. J. Heterogeneity and
415 changes in preferences for dying at home: a systematic review. *BMC Palliat. Care 12*,
416 (2013).
- 417 5. Barnato, A. E. *et al.* Are Regional Variations in End-of-Life Care Intensity Explained by
418 Patient Preferences? A Study of the US Medicare Population. *Med. Care 45*, 386–393
419 (2007).
- 420 6. Angus, D. C. *et al.* Use of intensive care at the end of life in the United States: an
421 epidemiologic study. *Crit. Care Med. 32*, 638–643 (2004).
- 422 7. Teno, J. M. *et al.* Change in End-of-Life Care for Medicare Beneficiaries: Site of Death,
423 Place of Care, and Health Care Transitions in 2000, 2005, and 2009. *J. Am. Med. Assoc.*
424 **309**, 470–477 (2013).
- 425 8. Teno, J. M. *et al.* Site of death, place of care, and health care transitions among US
426 medicare beneficiaries, 2000-2015. *JAMA - J. Am. Med. Assoc. 320*, 264–271 (2018).
- 427 9. Bekelman, J. *et al.* Comparison of Site of Death, Health Care Utilization, and Hospital
428 Expenditures for Patients Dying With Cancer in 7 Developed Countries. *JAMA 315*, 272–
429 283 (2016).
- 430 10. Wunsch, H. *et al.* Use of Intensive Care Services during Terminal Hospitalizations in
431 England and the United States. *Am J Respir Crit Care Med 180*, 875–880 (2009).
- 432 11. Anesi, G. L., Admon, A. J., Halpern, S. D. & Kerlin, M. P. Understanding irresponsible use
433 of intensive care unit resources in the USA. *Lancet Respir. 7*, 605–612 (2019).

- 434 12. Huynh, T. N. *et al.* The frequency and cost of treatment perceived to be futile in critical
435 care. *JAMA Intern. Med.* **173**, 1887–94 (2013).
- 436 13. Palda, V. A., Msw, K. W. B., Mclean, R. F. & Chapman, M. G. ‘Futile’ care: Do we provide
437 it? Why? A semistructured, Canada-wide survey of intensive care unit doctors and
438 nurses. *J. Crit. Care* **20**, 207–213 (2005).
- 439 14. Piers, R. D., Max, A., Michalsen, A., Maia, P. A. & *et al.* Perceptions of Appropriateness
440 of Care Among European and Israeli Intensive Care Unit Nurses and Physicians. *J. Am.*
441 *Med. Assoc.* **306**, 2694–2703 (2011).
- 442 15. Brownlee, S. *Overtreated: Why too much Medicine is Making us Sicker and Poorer.*
443 (Bloomsbury, 2007).
- 444 16. Teno, J. *et al.* Association of Increasing Use of Mechanical Ventilation Among Nursing
445 Home Residents With Advanced Dementia and Intensive Care Unit Beds. *JAMA Intern.*
446 *Med.* 1–8 (2016) doi:10.1001/jamainternmed.2016.5964.
- 447 17. Halpern, N., Debra, G., Tan, K. S. & Pastores, S. Trends in critical care beds and use
448 among population groups and Medicare and Medicaid beneficiaries in the United States:
449 2000-2010. *Crit. Care Med.* **44**, 1490–1499 (2016).
- 450 18. Zilberberg, M. D., Wit, M. De & Shorr, A. F. Accuracy of previous estimates for adult
451 prolonged acute mechanical ventilation volume in 2020: Update using 2000 –2008 data.
452 *Crit. Care Med.* **40**, 18–20 (2012).
- 453 19. Zilberberg, M. D. & Shorr, A. F. Prolonged acute mechanical ventilation and hospital bed
454 utilization in 2020 in the United States: implications for budgets , plant and personnel
455 planning. *BMC Health Serv. Res.* **8**, 1–10 (2008).
- 456 20. Kahn, J. *et al.* The Epidemiology of Chronic Critical Illness in the United States. *Crit. Care*
457 *Med.* **43**, 282–287 (2015).
- 458 21. Earle, C. C. *et al.* Trends in the aggressiveness of cancer care near the end of life. *J.*
459 *Clin. Oncol.* **22**, 315–21 (2004).

- 460 22. SUPPORT Principal Investigators. A Controlled Trial to Improve Care for Seriously Ill
461 Hospitalized Patients. *J. Am. Med. Assoc.* **274**, 1591–1597 (1995).
- 462 23. Carson, S. S. *et al.* Effect of Palliative Care–Led Meetings for Families of Patients With
463 Chronic Critical Illness: A Randomized Clinical Trial. *J. Am. Med. Assoc.* **316**, 51–62
464 (2016).
- 465 24. Curtis, J. R. *et al.* Effect of communication skills training for residents and nurse
466 practitioners on quality of communication with patients with serious illness: a randomized
467 trial. *JAMA* **310**, 2271–81 (2013).
- 468 25. Schneiderman, L. *et al.* Effect of Ethics Consultations on Nonbeneficial Life-Sustaining
469 Treatments in the Intensive Care Setting. *J. Am. Med. Assoc.* **290**, 1166–1172 (2003).
- 470 26. Nicholas, L. H., Langa, K. M., Iwashyna, T. & Weir, D. Regional Variation in the
471 Association Between Advance Directives and End-of-Life Medicare Expenditures. *JAMA*
472 **306**, 1447–1453 (2011).
- 473 27. Barnato, A. E., Tate, J. A., Rodriguez, K. L., Zickmund, S. L. & Arnold, R. M. Norms of
474 Decision Making in the ICU: A Case Study of Two Academic Medical Centers at the
475 Extremes of End-of-life Treatment Intensity. *Intensive Care Med.* **38**, 1886–1896 (2012).
- 476 28. Quill, C. M., Ratcliffe, S. J., Harhay, M. O. & Halpern, S. D. Variation in Decisions to
477 Forgo Life-Sustaining Therapies in US ICUs. *Chest* **146**, 573–82 (2014).
- 478 29. Hart, J. L. *et al.* Variability Among US Intensive Care Units in Managing the Care of
479 Patients Admitted With Preexisting Limits on Life-Sustaining Therapies. *JAMA Intern.*
480 *Med.* **19104**, (2015).
- 481 30. Fulton, A. T., Gozalo, P., Mitchell, S. L., Mor, V. & Teno, J. M. Intensive Care Utilization
482 among Nursing Home and Severe Functional Impairment. *J. Palliat. Med.* **17**, 313–317
483 (2014).
- 484 31. Dzeng, E. *et al.* Influence of institutional culture and policies on do-not-resuscitate
485 decision making at the end of life. *JAMA Intern. Med.* **175**, 812–820 (2015).

- 486 32. Kruser, J. M. *et al.* Assessment of Variability in End-of-Life Care Delivery in Intensive
487 Care Units in the United States. *JAMA Netw. Open* **2**, 1–16 (2019).
- 488 33. Halpern, N., Pastores, S. M., Chou, J. F., Chawla, S. & Thaler, H. Advance Directives in
489 an Oncologic Intensive Care Unit: A Contemporary Analysis of their Frequency, Type,
490 and Impact. *J. Palliat. Med.* **14**, 483–489 (2011).
- 491 34. Hartog, C. S. *et al.* Are written advance directives helpful to guide end-of-life therapy in
492 the intensive care unit? A retrospective matched-cohort study. *J. Crit. Care* **29**, 128–133
493 (2014).
- 494 35. Goodman, M., Tarnoff, M. & Slotman, G. Effect of advance directives on the management
495 of elderly critically ill patients. *Crit. Care Med.* **26**, 701–704 (1998).
- 496 36. Dzeng, E. *et al.* Homing in on the Social : System-Level Influences on Overly Aggressive
497 Treatments at the End of Life. *J. Pain Symptom Manage.* **55**, 282-289.e1 (2018).
- 498 37. Barnato, A. E. *et al.* Advance Care Planning Norms May Contribute to Hospital Variation
499 in End-of-Life ICU Use: A Simulation Study. *Med. Decis. Mak.* **34**, 473–484 (2014).
- 500 38. Curtis, J. R., Engelberg, R. A. & Teno, J. M. Understanding variability of end-of- life care
501 in the ICU for the elderly. *Intensive Care Med.* **43**, 94–96 (2017).
- 502 39. Van Den Bulcke, B. *et al.* Ethical decision-making climate in the ICU: Theoretical
503 framework and validation of a self-assessment tool. *BMJ Qual. Saf.* 1–9 (2018)
504 doi:10.1136/bmjqs-2017-007390.
- 505 40. Swidler, A. Culture in Action: Symbols and Strategies. *Am. Sociol. Rev.* **51**, 273–286
506 (1986).
- 507 41. Bourdieu, P. *Outline of a Theory of Practice.* (1977).
- 508 42. Dartmouth Atlas Project: End of Life Care. *Dartmouth Atlas Project*
509 <https://www.dartmouthatlas.org/interactive-apps/data-by-hospital/> (2020).
- 510 43. Barnato, A. E. *et al.* Advance care planning norms may contribute to hospital variation in
511 end-of-life ICU use: a simulation study. *Med. Decis. Mak.* **34**, 473–84 (2014).

- 512 44. Giddens, A. *The Constitution of Society: An Outline of the Theory of Structuration*. (Polity
513 Press, 1984).
- 514 45. Dzung, E. *et al.* Homing in on the Social: System-Level Influences on Overly Aggressive
515 Treatments at the End of Life. *J. Pain Symptom Manage.* (2017)
516 doi:10.1016/j.jpainsymman.2017.08.019.
- 517 46. Blythe, J. A. *et al.* An Interprofessional Process for the Limitation of Life-Sustaining
518 Treatments at the End of Life in France. *J. Pain Symptom Manage.* **00**, 1–11 (2021).
- 519 47. Dzung, E., Bein, T. & Curtis, J. R. The role of policy and law in shaping the ethics and
520 quality of end - of - life care in intensive care. *Intensive Care Med.* (2022)
521 doi:10.1007/s00134-022-06623-2.
- 522 48. Biernacki, P. & Waldorf, D. Snowball Sampling: Problems and Techniques of Chain
523 Referral Sampling. *Sociol. Methods Res.* **10**, 141–163 (1981).
- 524 49. Naderifar, M., Goli, H. & Ghaljaie, F. Snowball Sampling: A Purposeful Method of
525 Sampling in Qualitative. *Strides Dev Med Ed* **14**, (2017).
- 526 50. Bowen, G. a. Naturalistic inquiry and the saturation concept: a research note. *Qual. Res.*
527 **8**, 137–152 (2008).
- 528 51. Guest, G., Bunce, A. & Johnson, L. How Many Interviews Are Enough?: An Experiment
529 with Data Saturation and Variability. *Field methods* **18**, 59–82 (2006).
- 530 52. Lofland, J., Snow, D., Anderson, L. & Lofland, L. *Analyzing Social Settings: A Guide to*
531 *Qualitative Observation and Analysis*. (2006).
- 532 53. Emerson, R., Fretz, R. & Shaw, L. *Writing Ethnographic Fieldnotes*. (University of
533 Chicago Press, 2011).
- 534 54. Hammersley, M. & Atkinson, P. *Ethnography: Principles in Practice*. (Tavistock
535 Publications, 1983).
- 536 55. Braunn, V. & Clarke, V. Using thematic analysis in psychology. *Qual. Res. Psychol.* **3**,
537 77–101 (2006).

- 538 56. Creswell, J. W. *Research Design: Qualitative, Quantitative, and Mixed Methods*
539 *Approaches*. (Sage Publications, 2003).
- 540 57. Kaufman, S. *And a Time to Die: How American Hospitals Shape the End of Life*.
541 (University of Chicago Press, 2006).
- 542 58. Kaufman, S. *Ordinary Medicine: Extraordinary Treatments, Longer Lives, and Where to*
543 *Draw the Line*. (Duke University Press, 2015).
- 544 59. Kruser, J. M., Cox, C. E. & Schwarze, M. L. Clinical momentum in the intensive care unit:
545 A latent contributor to unwanted care. *Ann. Am. Thorac. Soc.* **14**, 426–431 (2017).
- 546 60. Hua, M. *et al.* What Affects Adoption of Specialty Palliative Care in Intensive Care Units :
547 A Qualitative Study. *J. Pain Symptom Manage.* **00**, 1–10 (2021).
- 548 61. Michalsen, A. *et al.* Interprofessional shared decision-making in the ICU: A systematic
549 review and recommendations from an expert panel. *Crit. Care Med.* **47**, 1258–1266
550 (2019).

Table 1: Hospital selection based upon Dartmouth Atlas indices around intensities of end-of-life care⁴²

Variable	High Intensity	Medium Intensity	Low Intensity
Inpatient Medicare Reimbursements per Patient during Last Two Years of Life	107130	97534	39170
Hospital Days per Patient during Last Six Months of Life	17.4	14.3	8.6
Total ICU days per Patient during Last Six Months of Life	11.1	3	2.5
Percent of deaths which included ICU admission	30	18.7	34.3
Percent of deaths occurring in hospital	43	32	31.5
Number of Different Physicians Seen per Decedent in Last Six Months of Life	16.8	13.6	10.9
Hospital Care Intensity Index	0.98	0.72	0.09

*Hospital Care Intensity Index = Based upon number of days patient spent in the hospital and the number of physician encounters they experienced as inpatients, and constitutes the ratio of a given hospital's utilization rate compared to the national average.

560 **Table 2:** Table of illustrative quotations
561

Hospital	Quotation	Quote number
<i>Defaults of high-intensity care and consistency between Dartmouth Atlas-measured care intensity and respondents' accounts of hospital culture</i>		
Medium Intensity	"Sometimes I feel like it's just gotten to the point where it's like bordering on true cruelty...I would say almost everything almost always gets offered...and then we're put in a situation of having to carry out all of these things that we don't really agree with and that can be a really distressing situation." (ICU MD6)	1
Low Intensity	"[Treatments are] very high intensity...we've had occasions where pretty much everybody agreed that the care that we were providing was inappropriate... the reality is that we have to work within a framework of the way our system operates." (ICU MD4)	2
High Intensity	"There is a culture from the top of the healthcare system down. We are going to go to eleven on every situation, appropriate or not. It's not explainable that each individual physician has independently arrived at this very aggressive practice pattern...I think it's the institution wants to be aggressive and so that just kind of steeps everyone in doing that kind of stuff." (ICU MD2)	3
High Intensity	"If...the son is freaking out and saying that you're killing and murdering her and that you need to code her, they're going to ask you to code her because that's the path of least resistance...you're probably going to end up coding her because [it's] the easier thing to do." (ICU Nurse) (*Details have been changed to preserve patient anonymity)	4
Medium Intensity	"[If] we don't have a family member...or some other form of documentation...the assumption [is] of full court press and do everything that we can." (ED MD1)	5
Low Intensity	"Until the family members or the patient [decides], then [care] will be typically intensive." (ICU MD2)	6
Low Intensity	"In the United States...everyone has the opportunity to receive full care to the maximum amount of our abilities until they say no, or until, essentially, their body tells us that they can no longer tolerate it...that is 'the right thing to do'....it is [an] American right...There's a lot of conversations about 'If you don't do this, I will report you. I'll bring you to the media'...'Sue you,' kind of thing. That conversation comes up with disgruntled family members here often. (ICU APP2)	7
Low Intensity	"I think it's almost like passive diffusion of the knowledge and style and vision of how palliative care. I see that there's a lot more palliative care consults happening...I think you can't help but learn their approach...some of that knowledge or management rubs off on the primary teams." (Hosp MD2)	8
<i>Consensus and coordination required to de-escalate</i>		
Low Intensity	"The default [for unbefriended patients] is the aggressive care pathway. But I think what ends up happening is we get to know them and their character...We try to reach out to case managers, to SNF providers... people in the community...but that takes time... [if] we are unable to track anyone down...we ultimately get our ethics	9

	committee involved and do a deeper dive into how we should proceed.” (PC MD5)	
Low Intensity	“We are not interested in blocking or making the upfront decision-making hard to get into the ICU. We’d much prefer to be open to bringing people into the ICU and then do a good job with care decision-making there”. (ICU MD 3)	10
Low Intensity	“I think in our institution, It’s pretty easy to move to CMO (Comfort Measures Only)...and deescalate. That tends to be well-established. (ED MD2)	11
Low Intensity	“We don’t have to offer certain things...we may in the course of discussions say, ‘This is perhaps not the most appropriate patient to offer renal replacement therapy to.’” (ICU MD2)	12
Low Intensity	“Generally it’s if there is not consensus between the teams, the default is to then allow whatever decision the family makes to just ride...we’ll just go ahead and allow that to evolve.” (Neuro ICU MD)	13
High Intensity	“I think it can be really distressing to see...[patients] suffering, and to see them getting treatments that you know that they wouldn’t want...it’s just really hard to see people getting forced into things that you know that they wouldn’t want here. But you have essentially no control to stop it. Because if the doctors aren’t going to say it, I can try, but then they may or may not listen to me, and then the family doesn’t want to. I don’t have a choice. I’m forced to do it.” (ICU Nurse2)	14
Medium Intensity	“Most of the time when we say we’re not going to offer a treatment such as dialysis, I can’t remember a family pushing really hard to do it. Especially when we all come together as a team...There are usually multiple teams involved to reinforce that we don’t think there’s going to be any benefit from this.” (ICU MD6)	15
Low Intensity	“I would say more often than not, it feels like the various teams are on the same page. There are a few instances in which nephrology is the team to raise the alarm bells of, ‘Hey, we’re being asked to offer this intervention and it just doesn’t make sense. And we don’t feel comfortable doing it.’...we have a discussion with our colleagues and we’re all on the same page about what makes the most sense.” (IM PGY-3)	16
Low Intensity	“I call it ‘the system wins’...the system dictates where it goes...I can say, ‘I don’t want to do this. I don’t think it’s in best interest; I talk to the patient and the family.’ The nocturnist who’s cross covering, sees the labs and doesn’t feel comfortable...and then next thing I know, they’re like, ‘I just decided that we should consult nephrology.’...I don’t think our system’s created in a way that unless everybody’s fundamentally feeling it’s futile, It’s almost like a majority gets what they want for the patient.” (Hospitalist MD2)	17
<i>Cultural norms around crucial decision points</i>		
Medium Intensity	“There’s global input coming in from the [various] teams...looking at the whole global picture and saying, ‘This is not right. We need to really start putting the brakes on this thing.’...It’s interesting because you’ll start to bring this up, and then you’ll start to see the gearbox move, and then in a day or two you can start to see it swing. The idea is implanted. I think part of that clinical momentum is that if you don’t	18

	have somebody else to look at that, then that idea never gets planted. (ICU MD5)	
Low Intensity	"I feel like there are a lot of checks in place to prevent untethered clinical momentum in the ICU...It just seems like things don't naturally just continue to accumulate and progress unchecked over time. We talk with our team members and with families pretty frequently about how things are going and then what makes the most sense to do next based on what has happened so far...Everyone's very thoughtful and thorough about making medical decisions and reassessing those decisions over time. (IM PGY3)	19
High Intensity	"[Doctors feel] it's not your job to decide whether or not someone's eligible and appropriate for a cath, it's the cardiologist's...We keep thinking that the sub-specialist is going to decide for us. And overwhelmingly [the] specialists want the primary [team], to decide before they call...all the specialists I talk to have a lot of grief about feelings about forced to do these procedures." (Palliative Care MD)	20
<i>Efforts to de-escalate nonbeneficial high-intensity life-sustaining treatments can be undermined</i>		
High Intensity	"Sometimes the ethics team will make a recommendation to make the patient a unilateral DNR if the patient has a poor prognosis...so we make [that recommendation]. The family then goes to the hospital leadership, and there are times when leadership has actually stopped us or stopped that unilateral DNR and said, 'Let's give it another week', which to be honest with you, it makes it challenging." (ICU Nurse)	21
High Intensity	"The family would not let go. She was dying for months and had no other treatment options left and [she] suffered horrendously...even though she had a DNR, I knew that it wouldn't be followed...Our attending said, 'She is no longer a candidate for [treatment] so if she decompensates... do not offer [treatment]...' So I stood in front of the room for hours, and every doctor that came in, I said, 'We're not offering [treatment]. Don't bring it up.'...Finally [the family started] to accept it...Later in the day, I came back and the [specialist] attending was in the room. The family said, 'Well she's not a candidate for [treatment].' [The [specialist] attending] then said, 'Well, Yes, we can do that [give treatment].' I was devastated. I'm not sure I've ever been so devastated in my whole life...[The attendings] got into a screaming match in the hallway...She eventually went back on [treatment], he coded and died. They did CPR on her." (ICU Nurse) (*Details have been changed to preserve patient anonymity)	22
Low Intensity	"The ethics committee was like, '[De-escalation is]...appropriate', but when they went to the hospital administration, they said, 'We can't. What if some family member emerges out of the background and then we made a decision to withdraw care. Let's just not escalate.'...even if all the clinicians agree, the institution puts a block on certain decisions because of their concerns around litigation." (PC MD 5)	23
High Intensity	"It made the whole system look like a fool. The whole ethics [committee] mechanism kind of fell apart before our eyes [in] that we brought this to [Ethics] Committee, [the] committee agreed with us,	24

	and then the next doctor was like, "Let's just trach and PEG and get them out of the ICU. This is not worth the fight." (ICU MD6)	
High Intensity	"Bringing [cases] to the Ethics Committee is not a small deal. It's made up of clinicians and community members, so you present the case and the family presents the case, but it almost seems like you're sitting in court. The members are asking you questions and questioning every single judgment and stuff like that...[the ethics chair] tells me that the ethics mechanism is supposed to be hard. It's a big deal to override a family, so you want to make sure that all your i's are dotted and all your t's are crossed." (ICU MD4)	25
Low Intensity	"There's still this discomfort around pushing against the aggressive care motto that we tend to default into. The ethics committee...can be this extra layer of support to clinicians to validate and verify the decisions that are being made...They liaise very closely with risk management because sometimes in these scenarios, a decision to not escalate or a decision to withdraw interventions can raise these flags for our institution...the ethics committee manages up and down and makes everyone feel like the decision is okay on multiple levels...They have a very low threshold to sort of get the institutional leadership involved...they have a good relationship and partnership with the administrators who also happen to be phenomenal clinicians themselves." (PC 5)	26
Medium Intensity	"The most surprising thing about the role that we play in risk management is the fact that unlike many risk managers we're often telling physicians "Please don't give care that's medically ineffective if it goes against your conscience or isn't in the best interest of the patient." It would be easier sometimes to just let it go on and on, but we advocate for people to please do the right thing, and that is not very common for a risk manager. The profession is reputed as being more risk adverse so I think my approach and my office's approach is kind of different than a lot of risk managers." (Risk Administrator 2)	27

562
563

564 **Table 3:** Table of illustrative quotations
565

<i>Institutional policies, protocols, practices, and resources shape hospital culture</i>		
Medium Intensity	"It's just part of our culture to consider palliative care...there are changes in the ICU like the comfort care huddle to have a more systematic approach...before meeting with the family to make sure that [everyone is] on board...Have we explored all the options? Do we all feel that there's consensus on a particular recommendation?" (ICU MD4)	28
Low Intensity	"There are now little checkpoints for teams who have previously been on the train that keeps going and doesn't stop...We have this [automated] algorithm that says make sure that if patients meet these criteria that you consult palliative care...within the first 24 hours." (PC MD 5)	29
Low Intensity	"death rounds [is where the]...palliative team provides for our residents to talk about really difficult cases or deaths...it's really important that they provide that space for the resident team...it was just so traumatizing. They would also just be present within our COVID ICU...[to] check in with our team and offer debrief sessions...that kind of support...was really important for our team as a whole." (ICU APP2)	30
Medium Intensity	"I have never experienced a time when medical administration has in any way got involved in any of these decisions. I wouldn't say that I've ever felt financial pressure or press pressure...it's just absent at this institution..." (Ethics Consult)	31
Low Intensity	"And it's normally not a 'you shouldn't do this.' [from hospital leadership], it's usually, 'our suggestion right now would be to wait or like to give more time'...it's a supportive culture...[for] decision making...it's mostly on our team." (ICU APP2)	32
<i>The relational and emotional aspects of end-of-life care</i>		
High Intensity	"I think it takes away a piece of your soul. You feel horrible about it because you're not accomplishing anything good in the long run, which is why I think a lot of physicians don't call ethics. They don't even fight that battle. Why fight that battle? Just do your shift and move on" (ICU MD5)	33
High Intensity	"It's like an exercise in futility almost...I think those cases are few. It's just that when they occur it just takes so much time, and energy, and work. Sometimes I wonder...is it worth it to even go through that whole process?" (ICU MD5)	34
Low Intensity	"I really value ending the situation on the same team as the family or the patient... I've got to align with this family. That's my number one priority...if I am at odds with the patient or the family and they feel like they're not getting the care that they want, I feel like that's a much bigger loss in my mind than someone getting CPR when they're 85 years old...it's just far sadder to me when they say, 'Those doctors didn't care' or 'they didn't try to save my mom'." (PC MD1)	35
Medium Intensity	"I think I'm aligned with the culture here...I don't know anybody who would disagree with the fundamental principles of building relationships...I really try to reserve unilateral decision-making as a last resort. As physicians we're making the final sacrifice to do what we think is best for the patient. But there's so much struggle and	36

	suffering that is going to continue after a patient dies with the family...if we can align with the families...I'm willing to wait and go the extra mile to try to achieve that." (Hospitalist MD1)	
--	---	--

566

Figure 1: Conceptual framework for the influence and interactions between hospital and national cultures and policies with illustrative examples from the data. (Adapted from Dzeng, 2022)⁴⁷





Figure 2: Model describing a prototypical patient trajectory for how institutional structures (e.g., policies, protocols, practices, and resource allocation) might impact the ease of de-escalation of non-beneficial high-intensity life-sustaining treatments.



All individuals and structures need to be aligned towards de-escalation for low-intensity cultures to succeed. Any one entity within the hospital can thwart de-escalation. As such, there are multiple ways to achieve high-intensity care but one pathway where all entities are aligned towards de-escalation for de-escalation to succeed.