Lawrence Berkeley National Laboratory

Recent Work

Title

TWO DIMENSIONAL ZERO FIELD NMR AND NQR

Permalink

<https://escholarship.org/uc/item/9s44m9gh>

Authors

Thayer, A.M.

Millar, J.M.

Pines, A.

Publication Date

1985-12-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

Submitted to Chemical Physics Letters

TWO DIMENSIONAL ZERO FIELD NMR AND NQR

A.M. Thayer, J.M. Millar, and A. Pines

December 1985

RECEIVED
LAWRENCE
BERKELEY LABORATORY

May 1986

LIBRARY AND
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*

Prepared for the U.S. Department of Energy under Contract DE-AC03-76SF00098

LBL-20932
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

TWO DIMENSIONAL ZERO FIELD NMR AND NQR

A.M. Thayer, J.M. Millar[†] and A. Pines

Department of Chemistry, University of California, Berkeley and
Materials and Molecular Research Division, Lawrence Berkeley Laboratory,
Berkeley, California 94720

[†]Current Address: Department of Chemistry, Yale University, New Haven, CT

Abstract

Correlations between quadrupolar frequencies in a spin $I=1$ system are determined through a two-dimensional version of the zero field NMR experiment. Results presented on a selectively deuterated polycrystalline solid illustrate this for the quadrupolar lines corresponding to inequivalent deuterium sites.

1. Introduction

Two dimensional NMR methods involve recording the NMR signal as a function of two time variables with subsequent transformation to produce a spectrum described by two frequency variables.¹ Applications of these experiments are generally to weakly coupled liquids and the frequencies observed depend upon the specific excitation pulse sequence used. Common examples are those which show correlations between chemical shifts and scalar couplings,² between different chemical shifts^{1,3} exchanging or cross relaxing dipolar coupled nuclei,⁴ and multiple quantum transitions.^{1b} By measuring the connectivities between spectral transitions, a determination of structure, conformation, dynamics or assignment of an otherwise intractable one dimensional spectrum is possible.

In Zero Field NMR and NQR, dipolar and quadrupolar interactions in the absence of an applied magnetic field are observed.⁵ These experiments yield well-resolved, sharp line spectra in polycrystalline solids. In zero field, quadrupolar frequencies label specific chemical sites while dipolar couplings should induce connectivities between zero field transitions of neighboring spins. In this communication, we demonstrate the principles of two-dimensional NMR to the detection of quadrupolar nuclei in zero applied magnetic field. Through a combination of these techniques, the connectivities in the NQR transitions of a spin $I = 1$ nucleus are shown.

2. Zero Field NQR Spectra

A spin $I = 1$ quadrupolar nucleus, such as deuterium, has three energy levels in zero field.⁶ For a non-axially symmetric electric field gradient, these levels are non-degenerate linear combinations of the high field states. The allowed transitions are labeled ν_0 , ν_- and ν_+ , in order of in-

creasing frequency. The dependence of these frequencies on quadrupole coupling constants (e^2qQ/h) and asymmetry parameter (η) are given by⁶:

$$\nu_+ = (3 + \eta)K$$

$$\nu_- = (3 - \eta)K$$

$$\nu_0 = 2\eta K = \nu_+ - \nu_-$$

in which $K = e^2qQ/4h$ for $I = 1$. Values of η and e^2qQ/h , which are descriptive of a quadrupolar system, may be calculated for a given chemical site assuming that the pair of corresponding ν_- and ν_+ lines is distinguishable. If there are two or more inequivalent sites, the NQR spectrum will consist of a superposition of six or more lines. Therefore, resolution of the difference frequency ν_0 lines at very low frequencies is essential in assigning the one dimensional spectrum.

The one dimensional zero field NQR spectrum of the methylene group of polycrystalline diethylterephthalate is shown in Figure 1. The spectrum was obtained using the selective indirect detection method which has been described in detail elsewhere.⁷ Essentially this experiment involves using the $^1H - ^2H$ level crossings which occur as a consequence of the field cycle to polarize and detect the NQR spectrum.⁸ Pulsed dc fields are used to selectively excite the zero field evolution of only the deuterons. The two deuterons of a particular methylene group are inequivalent resulting in the six line spectrum. The low frequency lines are resolved and allow for calculation of quadrupole coupling constants and asymmetry parameters for the inequivalent sites. If more sites were present, it becomes evident that overlapping lines would make assignment difficult.

3. Two Dimensional Zero Field Experiment

The correlations between the NQR frequencies can also be observed by probing the connectivities in the spin $I = 1$ manifold through a two dimensional version of the field cycle. The simplest form of the field cycle, in which the zero field interval is divided in half, is shown in Figure 2a. The sample is demagnetized to an intermediate field, B_{int} , which is switched off suddenly to initiate evolution. The system evolves under the quadrupolar Hamiltonian for the time t_1 . Application of a short dc pulsed field will transfer coherence between the energy levels of the spin 1 system. Evolution in zero field continues after the pulse for a time, t_2 . Reapplication of the field and remagnetization provides for sampling of the magnetization in high field. In successive field cycles, the zero field periods are incremented independently to produce a time domain signal as a function of t_1 and t_2 which when Fourier transformed produces the two dimensional zero field spectrum.

Other initial conditions are also possible in this experiment. In actual practice, for the increase in sensitivity and potentially shorter 1H T_1 relaxation times, the signal can be detected indirectly via the protons as described previously. The field cycle of Figure 2b uses demagnetization to zero field followed by pulsed dc magnetic fields to initiate and terminate zero field evolution (rather than the sudden transition of B_{int}). Of course, the behavior of the spin system has the same dependence on the natural quadrupolar frequencies, as in the version with sudden transitions in the intermediate field, although now the intensities of the spectrum will also be a function of the initial and final dc pulse angles and the dynamics of the level crossings. A short dc pulsed field can be applied to mix coherences in the same manner as in Figure 2a.

In addition to observing quadrupolar frequencies, connectivities between dipolar coupled groups of spins might also be established by altering the mixing period of the experimental field cycle. Assuming a deuteron-deuteron dipolar coupling of $\nu_D \sim 1$ kHz, application of an intermediate field, as in Figure 2a, for ~ 1 msec ($\sim 1/\nu_D$) or longer should allow a coherence transfer between dipolar coupled spins. Thus correlations based on spatial proximity can also be developed. In cases where the coherences do not persist for longer than a few milliseconds, signal might be conserved if the evolving magnetization is stored as populations by application of a dc pulse as shown in Figure 2c. Thus the decay is now described by a time constant related to T_{1z} which is expected to be longer than the decay of the coherences in solids. By slowly cycling the field to an intermediate value, the energy levels of inequivalent deuterons may be brought into contact. This is similar to the signal enhancement approach via repeated level crossings used in frequency domain double resonance NQR experiments.⁹

As an experimental verification of the applicability of these field cycles, the two-dimensional zero field spectrum of the high frequency ν_- and ν_+ lines of the same methylene sites in polycrystalline diethylterephthalate- d_4 is shown in Figure 3. The field cycle of Figure 2b was employed to indirectly and selectively detect only the deuterium NQR signal. The diagonal peaks fall along the $\nu_1 = \nu_2$ line and reproduce the one dimensional spectrum. Off-diagonal peaks correspond to the ν_-/ν_+ connectivities in the spin $I = 1$ manifold and are illustrated by the connecting lines. One connected pair of lines belongs to one deuteron of the CD_2 group and the second pair belongs to the other inequivalent deuteron.

4. Summary

NQR transitions in frequency domain experiments are often assigned on the basis of weakly allowed double transition peaks,¹⁰ double irradiation of two NQR lines⁹, or by the shift in frequencies due to application of a low field¹¹ since the low frequency lines are often obscured or unobserved.⁹ Time domain techniques⁵ have the low frequency detection capabilities, resolution and selectivity to assign transitions on the basis of ν_0 lines. All of these approaches are plagued by the problem of increased complexity of ^{the} spectrum with increasing numbers of quadrupolar sites. Two dimensional zero field NMR experiments can address many of these problems by utilizing the cross-peak correlations to determine connectivities. A large variety of experimental conditions, produced with different field cycles, can be envisioned with the use of pulsed magnetic fields to manipulate the spins in zero field.

Acknowledgements:

We would like to thank D.B. Zax for many useful discussions throughout the course of this work and D. Hugi-Cleary for help with the use of computer programs supplied by M. Gochin. This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Material Sciences Division of the U.S. Department of Energy and by the Director's Program Development Funds of the Lawrence Berkeley Laboratory under Contract No. DE-AC03-76SF00098.

References

1. (a) J. Jeener, Ampere International Summer School, Basko Polje, Yugoslavia, (1971).
(b) W.P. Aue, E. Bartholdi and R.R. Ernst, J. Chem. Phys. 64 (1976) 2229.
2. (a) L. Müller, A. Kumar and R.R. Ernst, J. Chem. Phys. 63 (1975) 5490.
(b) W.P. Aue, J. Karhan and R.R. Ernst, J. Chem. Phys. 64 (1976) 4226.
(c) K. Nagayama, P. Bachmann, R.R. Ernst and K. Wüthrich, Biochem. Biophys. Res. Commun. 86 (1979) 218.
3. (a) A.A. Maudsley and R.R. Ernst, Chem. Phys. Lett. 50 (1977) 368.
(b) G. Bodenhausen and R. Freeman, J. Magn. Reson. 28 (1977) 471.
(c) A.A. Maudsley, L. Müller and R.R. Ernst, J. Magn. Reson. 28 (1977) 463.
4. J. Jeener, B.H. Meier, P. Bachmann and R.R. Ernst, J. Chem. Phys. 71 (1979) 4546.
5. D.B. Zax, A. Bielecki, K.W. Zilm, A. Pines and D.P. Weitekamp, J. Chem. Phys. 83 (1985) 4877.
6. (a) M.H. Cohen and F. Reif, Solid State Phys. 5 (1957).
(b) T.P. Das and E.L. Hahn, Solid State Phys. Supp. 1, (1958).
7. J.M. Millar, A.M. Thayer, A. Bielecki, D.B. Zax and A. Pines, J. Chem. Phys. 83 (1985) 934.
8. (a) A.G. Redfield, Phys. Rev. 130 (1963) 589.
(b) R.L. Strombotne and E.L. Hahn, Phys. Rev. A133 (1964) 1616.
(c) R.E. Slusher and E.L. Hahn, Phys. Rev. 166 (1968) 332.
(d) J. Koo and Y.N. Hsieh, Chem. Phys. Letts. 9 (1971) 238.
(e) Y. Hsieh, J.C. Koo and E.L. Hahn, *ibid*, 13 (1972) 563.

9. D.T. Edmonds, Physics Reports C29 (1977) 233.
10. D.T. Edmonds and A.A.L. White, J. Magn. Reson. 31 (1978) 149.
11. (a) E.A. Keiter, Y. Hiyama and T.L. Brown, J. Mol. Struct. 111 (1983) 1.
(b) D.A. d'Avignon and T.B. Brown, J. Phys. Chem. 85 (1981) 4073.

Figure Captions

Figure 1. One dimensional zero field deuterium NQR spectrum of selectively deuterated diethylterephthalate- d_4 . The spectrum was detected indirectly via $^1\text{H} - ^2\text{H}$ level crossings. Six lines are resolved corresponding to the $\nu_0/\nu_-/\nu_+$ transitions of two inequivalent methylene deuterons. Through the use of the low frequency ν_0 lines, combinations of peaks may be assigned as the ν_-/ν_+ pair corresponding to a single quadrupolar site.

Figure 2. Field cycles for two dimensional zero field experiments. a) The sample is demagnetized to an intermediate field, B_{int} , which is switched off suddenly to initiate evolution in zero field. The evolution period is divided into two time periods, t_1 and t_2 , by reapplication of the field. If the dc field is applied as a brief pulse, it will act as a rotation on the spin system and mix coherences. Evolution then continues for t_2 and is terminated with a sudden reapplication of the intermediate field and remagnetization to high field. The signal is sampled as a function of the independently incremented time variables. If the applied field is longer in duration, ~ 0.5 -1 msec, correlations between dipolar coupled groups of spins should develop. b) Using demagnetization to zero field, indirect detection of the zero field NQR spectrum is possible. The short pulsed dc magnetic fields are used to initiate zero field evolution for t_1 , to mix coherences as in (a), and to terminate evolution after t_2 . Remagnetization after the zero field period again provides for the $^1\text{H} - ^2\text{H}$ level crossings and the detection of the signal in high field. c) The preparation of the spin system and t_1 evolution period are identical to that shown in (a). Applying a brief dc pulsed field will effectively store the magnetization, and

prevent the decay of coherences under the applied field. Cycling the field slowly to an intermediate level should provide for cross relaxation between groups of spins. Evolution is reinitiated for t_2 by a second dc pulse then halted and detected in the same manner as in (a) and (b).

Figure 3. Two dimensional zero field NQR spectrum of the high frequency ν_-/ν_+ transitions of polycrystalline diethylterephthalate- d_4 . The spectrum was obtained through indirectly detecting the ^2H signal via the protons as described in Figure 2b. By using pulses which were multiples of 2π for the protons, selective excitation of only the quadrupolar transitions is possible. The transfer of coherence between states in the spin $I = 1$ manifold was produced with a short dc pulsed field. Diagonal peaks along the $\nu_1 = \nu_2$ line correspond to the one dimensional spectra as shown in the projections. The peak positions of the one dimensional spectrum are indicated by the stick spectra. Cross peaks indicate correlations between the ν_+ and ν_- transition of an individual deuteron site. The connectivities are illustrated by the connecting lines.

XBL 862 - 10303

Figure 1
Thayer et. al.

XBL 863-11048

Figure 2
Thayer et. al.

XBL 864-11052

Figure 3
Thayer et. al.

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*