

UCLA

Proceedings of UCLA Health

Title

Carotid Epithelioid Hemangioendothelioma — A Management Dilemma

Permalink

<https://escholarship.org/uc/item/9sh365jf>

Journal

Proceedings of UCLA Health, 23(1)

Authors

Chi, Dorcas

Kazakova, Vera

Kozyreva, Olga

Publication Date

2020-02-11

CLINICAL VIGNETTE

Carotid Epithelioid Hemangioendothelioma – A Management Dilemma

Dorcas Chi, MD, Vera Kazakova, MD and Olga Kozyreva, MD

Case Report

A 58-year-old male presented to a local ENT with several months of progressive dysphagia to solid food, odynophagia, and transient painless vision impairment. Physical exam was noticeable for subtle right neck fullness without obvious lymphadenopathy. Neurological exam was normal. Flexible indirect laryngoscopy showed paralysis of right true vocal fold. CT angiography (CTA) of the neck followed by duplex ultrasound demonstrated a mass within the right carotid bifurcation and proximal internal carotid artery concerning for a carotid bifurcation tumor. A cervical MRA demonstrated a mass described as possible thrombosed pseudo-aneurysm at right carotid artery bifurcation. He was placed on dual-antiplatelet therapy, but continued to deteriorate with worsening dysphagia and tongue weakness. He had multiple admissions due to aspiration pneumonia and eventually underwent feeding tube placement. Repeat CTA two months later showed growth of the soft tissue mass at the bifurcation raising concern for a malignant etiology. He underwent radical surgical resection of the mass, and pathology revealed epithelioid hemangioendothelioma (EHE) of the carotid artery with extravascular invasion into the jugular vein, hypoglossal and vagal nerve. Given the extent of his disease, he completed adjuvant concurrent chemo-radiation with weekly carboplatin and paclitaxel as radiation sensitizers. His symptoms gradually improved and he has been disease free for over one year.

Discussion

Epithelioid hemangioendothelioma (EHE) is a rare malignant vascular neoplasm that derives from endothelial cells. It can be variably aggressive with biological behavior between benign hemangioma and malignant hemangiosarcoma. Peak incidences are between the ages 30 to 50 years. The incidence in the US is about 20 cases per year with primary vascular EHE accounting for about 50% of cases. EHE can also occur in soft tissues, bones, liver and lungs. Over 30 vascular EHE cases have been previously reported predominantly from veins rather than arteries.¹ The diagnosis of intravascular EHE remains extremely challenging due to its rarity, and the difficulty in distinguishing it from thrombosis, as their radiographic characteristics are rather nonspecific. There are no specific imaging features or preferred imaging modality to differentiate primary intravascular EHE from atherosclerotic, inflammatory and thrombotic lesions.

Our patient had EHE involving the carotid artery and the tumor mimicked a thrombosed carotid artery aneurysm on serial imaging studies. Despite his presenting symptoms concerning for cranial nerve involvement, the diagnosis was delayed due to imaging challenges. We have found only two published cases of EHE arising from the carotid artery.^{2,3} Clinicians should be aware of carotid artery EHE to include in the differentials of carotid body masses. This is especially important when cranial nerve palsies are present. Further efforts are needed for a better radiologic characterization of such intravascular tumors as timely diagnosis and complete surgical intervention remains the primary treatment. The effectiveness of adjuvant chemotherapy and/or radiation therapy remains ambiguous. Adjuvant radiation after surgical resection can be explored for localized EHE in order to control the residual disease given the risk of recurrence. Systemic chemotherapy is primarily for widespread disease.

REFERENCES

1. **Charette S, Nehler MR, Whitehill TA, Gibbs P, Foulk D, Krupski WC.** Epithelioid hemangioendothelioma of the common femoral vein: Case report and review of the literature. *J Vasc Surg.* 2001 May;33(5):1100-3. Review. PubMed PMID: 11331856.
2. **Osawa S, Saito A, Shimizu H, Ogawa T, Watanabe M, Tominaga T.** A case of intravascular epithelioid hemangioendothelioma occurring 14 years after coil embolization for an extracranial internal carotid artery aneurysm. *J Vasc Surg.* 2012 Jan;55(1):230-3. doi: 10.1016/j.jvs.2011.06.108. Epub 2011 Sep 13. PubMed PMID: 21917400.
3. **Abraham MJ, Gregory ND, Chennupati SK.** Epithelioid hemangioma of the internal carotid artery: a case report supporting the reactive pathogenesis hypothesis of this vascular tumor. *Int J Pediatr Otorhinolaryngol.* 2014 Jul;78(7):1186-9. doi: 10.1016/j.ijporl.2014.04.050. Epub 2014 May 6. PubMed PMID: 24837865.