

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

Fatal Tension Pneumoperitoneum Due to Non-Accidental Trauma

Permalink

<https://escholarship.org/uc/item/9sn6572n>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 16(5)

ISSN

1936-900X

Authors

Thornton, Stephen L
Scott, Mark
Hunter, Jeremy

Publication Date

2015

DOI

10.5811/westjem.2015.7.28022

License

<https://creativecommons.org/licenses/by/4.0/> 4.0

Peer reviewed

Fatal Tension Pneumoperitoneum Due to Non-Accidental Trauma

Stephen L. Thornton, MD
Jeremy Hunter, DO
Mark Scott, DO

Department of Emergency Medicine, University of Kansas Hospital, Kansas City, Kansas

Section Editor: Sean O. Henderson, MD

Submission history: Submitted July 9, 2015; Accepted July 17, 2015

Electronically published October 20, 2015

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2015.7.28022

[West J Emerg Med. 2015;16(5):788-789.]

A previously healthy two-year-old boy presented to the emergency department with vomiting. He was cyanotic with mottling of both lower extremities. He was in respiratory distress with retractions and diminished breath sounds. His abdomen was distended and rigid. He had a pulse of 170 beats per minute, blood pressure of 144/69mmHg and respiratory rate of 42 breaths per minute. He was endotracheally intubated. Chest and abdominal radiographs demonstrated a tension pneumoperitoneum (Figure 1).

Abdominal decompression was performed with a 16-gauge

needle in the left lower quadrant. Bilateral tube thoracotomies were also performed. Post-decompression radiograph demonstrated continued free air but normal lie of organs and viscera (Figure 2). The patient then went into cardiopulmonary arrest. Chest compressions, epinephrine, bicarbonate, atropine and calcium gluconate were administered, but he did not regain spontaneous circulation. Subsequent autopsy and investigation determined the patient had been a victim of non-accidental trauma resulting in gastric rupture.

In pediatric patients tension pneumoperitoneum is a rare




Figure 1. White arrows demonstrate medial displacement of viscera. Free air is present.



Figure 2. After decompression viscera demonstrate normal lie. Free air is still present.

complication described after reduction of intussusceptions, mouth-to-mouth breathing, iatrogenic bowel perforations, and positive pressure ventilation.¹⁻⁴ It has not been described as a complication of non-accidental trauma. The increase in intra-abdominal pressure causes multiple physiologic derangements including decreased cardiac return via compression of the inferior vena cava and respiratory failure due to splinting of the diaphragms.³

Initial symptoms include abdominal pain and distension followed by hypoxia and shock.¹⁻⁴ Diagnosis is clinical, but radiographs will demonstrate free air and medial displacement of the solid organs and viscera.

If not recognized and promptly treated tension pneumoperitoneum can rapidly lead to cardiopulmonary arrest. Treatment is emergent needle decompression followed by definitive laparotomy repair.⁴ Emergency medicine clinicians should be familiar with tension pneumoperitoneum as a cause of respiratory distress and cardiovascular collapse in the pediatric patient, as early recognition and treatment is critical in improving survival.

Address for Correspondence: Stephen L. Thornton, MD, University of Kansas Hospital, Department of Emergency Medicine, 3901 Rainbow Blvd, 4045 Delp, Kansas City, KS 66160. Email: sthornton@kumc.edu.

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

Copyright: © 2015 Thornton et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. O'Hanlon KP. Gastric rupture with pneumoperitoneum after mouth-to-nose breathing in an infant. *J Emerg Med.* 2010;39(3):312-5.
2. Fallon SC, Kim ES, Naik-Mathuria BJ, et al. Needle decompression to avoid tension pneumoperitoneum and hemodynamic compromise after pneumatic reduction of pediatric intussusception. *Pediatr Radiol.* 2013;43(6):662-7.
3. Hughes DB, Judge TN, Spigland NA. Tension pneumoperitoneum in a child resulting from high-frequency oscillatory ventilation: a case report and review of the literature. *J Pediatr Surg.* 2012;47(2):397-9.
4. Chiapponi C, Stocker U, Körner M, et al. Emergency percutaneous needle decompression for tension pneumoperitoneum. *BMC Gastroenterol.* 2011;11:48.