

UC Berkeley

UC Berkeley Previously Published Works

Title

Coordination of 2,2'-(Trifluoroazanediy)bis(N,N'-dimethylacetamide) with U(VI), Nd(III), and Np(V): A Thermodynamic and Structural Study

Permalink

<https://escholarship.org/uc/item/9ss469qw>

Journal

Inorganic Chemistry, 58(23)

ISSN

0020-1669

Authors

Gao, Yang
Parker, Bernard F
Dau, Phuong V
[et al.](#)

Publication Date

2019-12-02

DOI

10.1021/acs.inorgchem.9b02469

Peer reviewed

Coordination of 2,2'-(Trifluoroazanediy)bis(N,N'-dimethylacetamide) with U(VI), Nd(III), and Np(V): A Thermodynamic and Structural Study

Yang Gao,^{a,b} Bernard F. Parker,^{a,c} Phuong V. Dau,^a Phuong D. Dau,^a Trevor D. Lohrey,^{a,c} John K. Gibson,^a John Arnold,^{a,c} Simon J. Teat,^{a,d} Andrea Melchior,^e Zhicheng Zhang,^{a,*} Linfeng Rao^{a,*}

^aChemical Sciences Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, California 94720, USA. E-mail: lrao@lbl.gov

^bCollege of Nuclear Science and Technology, Harbin Engineering University, Harbin, 150001, China

^cDepartment of Chemistry, University of California – Berkeley, Berkeley, CA 94720, USA

^dAdvanced Light Source, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, California 94720, USA

^eDipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Laboratori di Chimica, via delle Scienze 99, 33100 Udine, Italy

Abstract

Thermodynamic properties of the complexation of 2,2'-(trifluoroazanediy)bis(N,N'-dimethylacetamide) (CF₃ABDMA) with U(VI), Nd(III), and Np(V) have been studied in 1.0 M NaNO₃ at 25 °C. Equilibrium constants of the complexation were determined by potentiometry and spectrophotometry. In comparison with a series of structurally related amine-bridged diacetamide ligands, including 2,2'-(benzylazanediy)bis(N,N'-dimethylacetamide) (BnABDMA), 2,2'-azanediybis(N,N'-dimethylacetamide) (ABDMA), and 2,2'-

(methylazanediyl)bis(N,N'-dimethylacetamide) (MABDMA), CF₃ABDMA forms weaker complexes with U(VI), Nd(III), and Np(V) due to the lower basicity of the center N atom in CF₃ABDMA resulting from the attachment of the strong electron-withdrawing CF₃- moiety. The complexation strength of CF₃ABDMA with the three metal ions follows the order: UO₂²⁺ > Nd³⁺ > NpO₂⁺, consistent with the order of the “effective” charges of the metal ions. Structural information of the U(VI)/CF₃ABDMA complexes in solution and in solid was obtained by theoretical computation, single crystal X-ray diffractometry, ¹⁹F NMR, and electrospray ionization mass spectrometry. The structural data indicate that, similar to the three previously studied amine-bridged diacetamide ligands (BnABDMA, ABDMA, and MABDMA), the CF₃ABDMA ligand coordinates to UO₂²⁺ in a tridentate mode, through the center nitrogen and the two amide oxygen atoms.

Key Word: *Uranium, Neodymium, Neptunium, Amine-bridged diacetamide, Complexation, Thermodynamics, Coordination mode*

1. INTRODUCTION

Large quantities of minor actinides (MAs = Am, Np, and Cm) and fission products including trivalent lanthanides (Ln(III)) coexist in high level nuclear waste (HLW) generated during the reprocessing of spent nuclear fuel. Efficient separation of these radioactive elements from HLW can not

only recover valuable isotopes but also reduce the long-term heat loading and radioactive hazards in HLW, which is very beneficial to the final treatment and disposal of the nuclear wastes in geological repositories.

In an advanced nuclear fuel cycle, a strategy of partitioning and transmutation (P&T)^[1,2] is proposed. With the P&T strategy, the long-lived minor actinides are separated from the trivalent lanthanides and converted into isotopes that are stable or short-lived. Among the long-lived radionuclides, ²³⁷Np is of great concern because it is predicted to become a major radiation hazard, contributing 67% of the total radiation dose from HLW 75000 years after the closure of the repository.^[3] The pentavalent state of neptunium, Np(V), is the most stable oxidation state and exists as NpO_2^+ ions in solution. Due to its low ionic charge and large ionic radius, NpO_2^+ has a weak tendency toward hydrolysis, precipitation, or sorption by geomeia so that it is highly mobile in the environment. Also, NpO_2^+ does not form strong complexes with many ligands and is difficult to separate by many traditional extracting agents in actinide separation processes. Besides neptunium, separation of trivalent actinides (An(III)) from lanthanides is a challenging task because of the extremely high similarity of their chemical and physical properties. In addition, separation of uranium is always a significant task in the management of nuclear wastes and spent nuclear fuel reprocessing because uranium is the most abundant radioactive element in these processes. As a result,

there has been substantial interest in the development of efficient ligands and extractants for the separation of actinides and lanthanides to help develop an advanced nuclear fuel cycle.

In the past two decades, a group of alkyl-substituted diglycolamides (DGA), including N,N,N',N'-tetraoctyl-diglycolamide (TODGA),^[4,5] N,N,N',N'-tetrakisobutyl-diglycolamide (TiBDGA),^[6] and N,N'-dimethyl-N,N'-dihexyl-diglycolamide (DMDHDGA)^[7] have become a subject of high interest in the area of actinide partitioning.^[8] Actinides in nitric acid solutions can be effectively extracted into organic solvents containing these ligands and readily stripped back by diluted acid solutions.^[4,5,6,7,8,9,10,11] In addition, the diglycolamide ligands are completely combustible because they are composed of only C, H, O, and N atoms. Therefore, the final amount of solid radioactive waste can be greatly reduced by using these CHON ligands to replace the traditional extractants containing phosphorus and/or sulfur. To help understand the thermodynamic and structural properties of the actinide complexes with DGA ligands, a number of studies have been conducted in recent years by using small alkyl groups (e.g., methyl) to render the DGA ligands water-soluble so that a variety of techniques, such as potentiometry, spectrophotometry, and calorimetry, are applicable to the studies.^[12,13,14,15]

The alkyl-substituted DGA ligands are usually tridentate to the metal ions, using the two amide oxygen atoms and the center ether oxygen.

[12,13,14,15] Considering that no substitutional groups can be attached to the ether oxygen of DGA to vary the basicity and binding strength of the ligands, we have previously synthesized and conducted the studies of a series of structurally related amine-bridged diacetamide ligands, including 2,2'-(benzylazanediyl)bis(N,N'-dimethylacetamide) (BnABDMA), 2,2'-azanediylbis(N,N'-dimethylacetamide) (ABDMA), 2,2'-(methylazanediyl)bis(N,N'-dimethylacetamide) (MABDMA).^[16,17,18] The structures of these ligands are similar to that of N,N,N',N'-tetramethyldiglycolamide (TMDGA), but differ in the linkage between the two amide groups: the two amide groups are bridged by an amine group in the former, but an ether oxygen in the latter. As a result, different substitutional groups (e.g., a benzyl group in BnABDMA, a proton in ABDMA, and a methyl group in MABDMA) can be attached to the amine nitrogen, so that the basicity of the amine nitrogen and the binding strength of the amine-bridged diacetamides can be fine-tuned. It has been demonstrated that these ligands form stronger complexes with Nd^{3+} ,^[16] UO_2^{2+} ,^[17] or NpO_2^+ ,^[18] than TMDGA.^[12,13,15] In a tridentate mode similar to the DGA ligands, the amine-bridged diacetamides coordinate to the metal ions via the two amide oxygen atoms and the central nitrogen.

Among the three amine-bridged diacetamide ligands that were previously studied, the metal binding strength generally follows the order: MABDMA > ABDMA > BnABDMA,^[16,17,18] which is attributed to the electron-

donating ability of the methyl group in MABDMA and the electron-withdrawing ability of the benzyl group in BnABDMA. In this work, a new diacetamide ligand with a CF_3 - group substituting on the central nitrogen, 2,2'-(trifluoroazanediy)bis(N,N'-dimethylacetamide) (CF_3ABDMA , Figure 1), was synthesized. The CF_3 - group is known to be strongly electron-drawing and we hypothesized that the attachment of the CF_3 - group could make the electronegativity on the amine nitrogen even lower than that in BnABDMA. To test this hypothesis and extend the structure-property relationship in metal complexation with the amine-bridged diacetamides, thermodynamic properties of the complexation of CF_3ABDMA with UO_2^{2+} , Nd^{3+} , and NpO_2^+ were determined by potentiometry and spectrophotometry. Techniques including Density Functional Theory (DFT) computation, ^{19}F NMR, electrospray ionization mass spectrometry (ESI/MS), and X-ray crystallography were used to provide the structural information of the complexes. The thermodynamic and structural data from this work were discussed in comparison with the data for the three previously studied amine-bridged diacetamide ligands.





Figure 1. From left to right: CF₃ABDMA (this work), BnABDMA, ABDMA, and MABDMA.^[16,17,18] All ligands are neutral and donated as L, with their protonated forms donated as HL⁺, in this paper.

2. EXPERIMENTAL

2.1 Chemicals

2,2'-(trifluoroazanediy)bis(N,N'-dimethylacetamide), denoted as CF₃ABDMA in this paper, was prepared according to Scheme 1. 2-chloro-N,N-dimethylacetamide (13.4 g, 110 mmol) was added to a suspension of potassium iodide (18.2 g, 110 mmol) in acetonitrile (200 mL) and the mixture was stirred vigorously for 3 hours at room temperature under an atmosphere of dry nitrogen to produce 2-iodo-N,N-dimethylacetamide *in situ*. 2,2,2-trifluoroethylamine (5.0 g, 50 mmol) and sodium carbonate (28 g, 200 mmol) were then added, and the mixture was stirred for 6 hours at room temperature followed by heating to reflux for 16 hours. The mixture was cooled to room temperature, filtered, then the solvent removed under reduced pressure to give a viscous brown oil. The oil was extracted with chloroform (50 mL × 4), washed with 50 mL water, 20 mL portions of 10% Na₂S₂O₃ to remove I₂ until the solution was colorless, and 50 mL brine. The solution was dried over magnesium sulfate, passed through a plug of neutral alumina, and the solvent slowly evaporated at room temperature to give CF₃ABDMA as a very pale yellow solid (4.25 g, 32%). ¹H NMR (CDCl₃, 500 MHz): δ = 2.94 (s, 6H), 2.99 (s, 6H), 3.44 (q, ³J_{HF} = 10 Hz), 3.75

(s, 4H); ^{13}C NMR (CDCl_3 , 500 MHz): $\delta = 35.7, 36.7, 56.0, 169.8$, CH_2CF_3 carbon atoms not observed; ^{19}F NMR (CDCl_3 , 400 MHz): $\delta = -70.6$ (t, $^3J_{\text{FH}} = 10$ Hz).



Scheme 1. Synthesis of 2,2'-(trifluoroazanediyloxy)bis(N,N'-dimethylacetamide).

Milli-Q water was boiled and cooled (to avoid CO_2 contamination) before being used for the preparation of all solutions. All experiments were conducted at 25 °C and an ionic strength of 1.0 M NaNO_3 . The stock solutions of U(VI) and Nd(III) were prepared, respectively, by dissolving $\text{UO}_2(\text{NO}_3)_2 \cdot \text{H}_2\text{O}(\text{s})$ in dilute nitric acid solution and $\text{Nd}_2\text{O}_3(\text{s})$ (Aldrich, 99.9%) in HNO_3 under low heating. The concentrations of UO_2^{2+} and Nd^{3+} in the stock solutions were determined by complexometry titrations using ethylenediaminetetraacetic acid (EDTA).^[19] The stock solution of Np(V) was prepared by following a procedure slightly modified from that described elsewhere.^[20] The concentration of Np(V) was determined by the optical absorbance at 980.2 nm ($\epsilon = 395 \text{ M}^{-1} \text{ cm}^{-1}$). The concentrations of free acid in the stock solutions of U(VI), Nd(III), and Np(V) were determined by Gran's titration.^[21] The stock solution of CF_3ABDMA was prepared by

dissolving appropriate amount of the ligand in 1.0 M NaNO₃. The concentration of the ligand was calculated from the weight and verified by potentiometry. All chemicals are reagent grade or higher.

2.2 Potentiometry

Potentiometric titrations were conducted to determine the equilibrium constants of the protonation and complexation of the ligand with U(VI). The titration system, controlled by a computer, consists of a water-jacketed cell, a pH meter (Metrohm Model 713) equipped with a combination pH electrode (Orion Model 8102), and an automatic burette (Metrohm Model 765). The titrations were conducted at a constant temperature (25.0 ± 0.1)°C. A constant flow of argon was applied over the solution to avoid contamination of CO₂. The pH electrode was filled with 1 M NaCl to reduce the electrode junction potential.

In acidic and basic regions, the electromotive force (E , in millivolts) can be expressed by eq.(1) and eq.(2), respectively.

$$E = E^0 + RT/F \ln[H^+] + \gamma_H[H^+] \quad (1)$$

$$E = E^0 + RT/F \ln(Q_w/[OH^-]) + \gamma_{OH}[OH^-] \quad (2)$$

where the electrode parameters, E^0 , γ_H , and γ_{OH} , are determined by an acid-base calibration titration with standard HCl and NaOH solutions prior

to each titration. In equations (1) and (2), T is the temperature in Kelvin, F is the Faraday constant, R is the gas constant, and $Q_w = [H^+][OH^-]$ and the value of Q_w is $10^{-13.77}$ for the ionic medium of 1.0 M $NaNO_3$ in this work.

Duplicate or triplicate titrations were conducted with solutions of varying concentrations of the ligand, acidity and U(VI). For the determination of the protonation constant of the ligand, the cup solution containing the ligand at different acidities was titrated with NaOH or HNO_3 . For the determination of the equilibrium constants of the complexation with U(VI), the cup solution containing U(VI) and the ligand in an acidic solution was titrated with NaOH. The nonlinear regression program Hyperquad 2008 was used to fit the data and obtain the equilibrium constants.^[22]

2.3 Spectrophotometry

Spectrophotometry was used to study the complexation of CF_3ABDMA with U(VI), Nd(III) and Np(V). Absorption spectra of U(VI) (350-550 nm, 0.2 nm interval) Nd(III) (560-610 nm, 0.2 nm interval) and Np(V) (950-1065 nm, 0.2 nm interval) were collected on a Cary 6000i UV-Vis-NIR spectrophotometer (Varian Inc.) equipped with sample holders that were maintained at constant temperatures. Quartz cells with 10 mm optical path were used. Solutions containing the metal ions, U(VI), Nd(III) or Np(V), were placed in the cell and titrated with a solution of the ligand. After each addition, the solution was thoroughly mixed for 1-2 minutes

before the spectrum was collected. The mixing time is sufficient because prior studies have shown the absorbance became stable within 30 seconds of mixing. Usually, a set of 16-21 spectra were generated in one titration (with 15-20 additions of the titrant). The commercially available HypSpec program was used to analyze the data.^[22]

2.4 ¹⁹F NMR

To help evaluate the coordination of CF₃ABDMA with UO₂²⁺ and Nd³⁺, ¹⁹F NMR spectra of a few solution samples were acquired on a Bruker AVQ-400 instrument with a quad-channel probe. The samples include aqueous or methanol solutions of CF₃ABDMA in the absence or presence of metal ions (UO₂²⁺ and La³⁺, the latter is used as the analog of Nd³⁺ because it is diamagnetic). To verify the purity of the CF₃ABDMA, ¹H and ¹³C NMR spectra were acquired on a Bruker DRX-500 instrument with a Z-gradient proton/BBO probe.

2.5 X-Ray Crystallography

Pale yellow crystals of the 1:2 (U(VI):L) complex, UO₂L₂(ClO₄)₂, were obtained by slow evaporation from 0.5 mL solution containing 1 mM UO₂²⁺ and 5 mM ligand at pH ~ 6. Crystallographic data were collected on Beamline 11.3.1 (for Small-Crystal Crystallography) at the Advanced Light Source of Lawrence Berkeley National Laboratory (LBNL), with the Bruker

APEX II CCD diffractometer of ω rotation with narrow frames at a wavelength of 0.77490 Å. Intensity data were collected using Bruker Apex 2 software.^[23] The Bruker SAINT software package was used to perform intensity data integration, cell refinement, and data reduction.^[24] The crystal structure was solved using SHELXT and refined with SHELXL-2014.^[25] Hydrogen atoms were placed using a riding model, with freely refined torsion angles where relevant. A high peak of residual density was found to remain proximate to the uranium center, but attempts to detect a second domain or merohedral twin were not successful. Based on the size of this peak (ca. 4 electrons) and the atomic number of uranium (95), any twin component is likely to account for only a small amount of the diffraction, explaining the difficulty in modeling it. Details of the crystallographic data are in Table 1.

Table 1. Crystallographic data and Structural refinement

U(VI) complex	UO ₂ L ₂ (ClO ₄) ₂
Empirical formula	C ₂₀ H ₃₆ F ₆ Cl ₂ N ₆ O ₁₄ U
Formula weight	1007.48
Temperature	100(2) K
Wavelength	0.7749 Å
Crystal system	Monoclinic
Space group	<i>P</i> 2 ₁ /c
Unit cell dimensions	<i>a</i> = 9.037(10) <i>b</i> = 14.372(16) <i>c</i> = 13.640(16) α = 90° β = 107.783(12)° γ = 90°
Volume	1687(3) Å ³

Z	2
Density (calculated)	1.984 Mg/m ³
Absorption coefficient	2.906 mm ⁻¹
F(000)	980
Color, habit	Colorless, needle
Crystal size (mm ³)	0.080 x 0.020 x 0.020
Theta range for data collection	2.31 to 44.12°
Index ranges	-16 ≤ h ≤ 16 -24 ≤ k ≤ 25 -24 ≤ l ≤ 24
Reflections collected	35321
Independent reflections	7622 [R(int) = 0.0363]
Completeness to theta = 27.706°	100%
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	0.7458 and 0.4235
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	10100 / 0 / 227
Goodness-of-fit on F ²	1.057
Final R indices [I > 2sigma(I)]	R1 = 0.0328 wR2 = 0.0886
R indices (all data)	R1 = 0.0446 wR2 = 0.0967
Extinction coefficient	n/a
Largest diff. peak and hole	4.173 and -3.947 e.Å ⁻³

2.6 ESI-MS. Electrospray Ionization Mass spectrophotometry (ESI-MS)

experiments were performed to identify U(VI)/CF₃ABDMA complexes on an Agilent 6340 quadrupole ion trap mass spectrophotometer equipped with a micro electrospray ionization (ESI) source. Spray solutions of water-ethanol (< 5% water) were used in the experiments. The instrument has a detection range of 50 - 2200 m/z with a resolution of ~0.25 m/z in high resolution mode. The instrument parameters used to obtain the MS spectra of the complexes were similar to those previously employed.^[16,17]

2.7 Computational Studies

DFT calculations were carried out on the ligand and complexes using the three-parameter hybrid functional B3LYP.^[26,27] This level of theory has previously been demonstrated to generate reliable structural and energetic results for actinide complexes.^[17,28,29] Stuttgart-Dresden small-core quasi-relativistic effective core potentials have been employed for the metal ions, while other elements have been treated using a 6-31++G(d,p) Gaussian-type basis set. Solvent effect has been taken into account by using the polarizable continuum model (PCM).^[30] Geometry optimizations of the ligand and the complexes were run without symmetry constraints in PCM solvent. The free energies of the complexes were calculated by adding to the electronic energy of each complex the zero-point energy and thermal corrections that comprise electronic, harmonic vibrational, rotational, and translational contributions to the internal energy. For the calculation of the reaction free energies in solution, the correction for the standard state was applied.^[17,31] All calculations were carried out using Gaussian16.^[32]

3. RESULTS AND DISCUSSION

3.1 Protonation of the ligand

Experimental conditions and plots of the protonation titrations are shown in Table S1 and Figure S1 in Supporting Information. The protonation constant of CF₃ABDMA ($\log K$) is calculated to be 4.00 ± 0.09 . This value is lower than those for the three previously studied amine-bridged ligands (Table 2). A trend in $\log K$ is observed as: CF₃ABDMA < BnABDMA < ABDMA < MABDMA. Because the protonation of all these ligands occurs on the central amine N atom, the trend in $\log K$ reflects the order of basicity on N. Among the ligands, CF₃ABDMA has the lowest basicity due to the strong electron-withdrawing ability of the CF₃- group.

Table 2 Protonation of CF₃ABDMA in comparison with BnABDMA, ABDMA, and MABDMA ($H^+ + L = HL^+$) at 25 °C and $I = 1.0$ M NaNO₃, p.w. – present work.

Ligand	$\log K$	ref.
CF ₃ ABDMA	4.00 ± 0.09	p.w.
BnABDMA	6.36 ± 0.09	[16]
ABDMA	7.12 ± 0.09	[16]
MABDMA	7.64 ± 0.09	[16]

3.2 Complexation of U(VI) with CF₃ABDMA

A representative potentiometric titration is shown in Figure 2. The titration curve is best fitted with the model including the formation of 1:1 U(VI) complex, UO₂L²⁺ as shown in reaction 3. In the calculation, the protonation

constant of CF₃ABDMA from this work, the stability constant of UO₂(NO₃)⁺,^[33] and the hydrolysis constants of three U(VI)/OH species, UO₂(OH)⁺, (UO₂)₂(OH)₂²⁺, and (UO₂)₃(OH)₅⁺,^[34] were also included. The calculated equilibrium constant for UO₂L²⁺ is shown in Table 3. The data are compared with those for Nd³⁺ and NpO₂⁺ and with other related diacetamide ligands in subsequent sections.




Figure 2. Potentiometric titration of U(VI)/L complexation at 25 °C and *I* = 1.0 M NaNO₃. Symbols: (•) experimental -log[H⁺]; (dashed line) fitted -log[H⁺]; (solid lines) U(VI) speciation, 1 – free UO₂²⁺, 2 - UO₂(NO₃)⁺, 3 - UO₂L²⁺, 4 - (UO₂)₂(OH)₂²⁺, 5 - (UO₂)₃(OH)₅⁺. This is titration #3, detailed titration conditions of which are shown in Table S2 of Supporting Information.

Table 3 Overall equilibrium constants for the complexation of CF₃ABDMA with UO₂²⁺, Nd³⁺, and NpO₂⁺ at 25 °C and *I* = 1.0 M NaNO₃, method: pot – potentiometry, sp – spectrophotometry.

Metal ion	Reaction	Method	logβ
UO ₂ ²⁺	UO ₂ ²⁺ + L = UO ₂ L ²⁺	pot	(2.05 ±

			0.12)
Nd ³⁺	Nd ³⁺ + L = NdL ³⁺	sp	(1.44 ± 0.09)
	Nd ³⁺ + 2L = NdL ₂ ³⁺		(1.90 ± 0.09)
NpO ₂ ⁺	NpO ₂ ⁺ + L = NpO ₂ L ⁺	sp	(0.46 ± 0.09)

Because a 1:2 U(VI)/CF₃ABDMA complex, UO₂L₂(ClO₄)₂, was obtained in the form of single crystal from aqueous solution (see sections 2.5 and 4.2), efforts were made to identify and determine the stability constant of 1:2 U(VI)/CF₃ABDMA complex in aqueous solution, UO₂L₂²⁺, by potentiometry as well as spectrophotometry. The efforts were not successful due to the following reasons. With potentiometry, higher ratios of [L]/[U] were used to facilitate the formation of UO₂L₂²⁺. Under such conditions, the protonation of the ligand dominates the titration and the change in [H⁺] becomes insensitive to the formation of U(VI)/CF₃ABDMA complexes. With spectrophotometry, a number of hydrolyzed U(VI) species that have significant optical absorption, including UO₂OH⁺, (UO₂)₂(OH)₂²⁺, and (UO₂)₃(OH)₅⁺, must be included in fitting the overall spectra, which proved to be extremely difficult. As a result, in aqueous solution, only the stability constant of the 1:1 U(VI)/CF₃ABDMA complex was obtained in this work.

3.3 Complexation of Nd(III) with CF₃ABDMA

Figure 3 shows a representative set of absorption spectra for the titration

of Nd(III) with CF₃ABDMA. With the addition of CF₃ABDMA, the intensities and positions of the branches of the absorption band varied as the ligand to metal ratio (L/Nd(III)) was increased. Factor analysis of the absorption spectra by the HypSpec program indicates that there are four absorbing species of Nd(III) including the free Nd³⁺ and Nd(NO₃)²⁺ complex that were present in the beginning of the titration. Accordingly, the spectra were best-fitted with the formation of two successive Nd(III)/CF₃ABDMA complexes, as shown by reaction (4):



where $n = 1$ and 2 . The stability constants of the 1:1 and 1:2 complexes between Nd³⁺ and CF₃ABDMA were calculated to be 1.44 ± 0.09 and 1.90 ± 0.09 , respectively. In the calculation, the protonation constant of CF₃ABDMA from this work and the stability constant of Nd(NO₃)²⁺ from the literature^[35] were used.




Figure 3 Spectrophotometric titrations of Nd^{3+} complexation with CF_3ABDMA (upper), and the calculated molar absorptivities of Nd(III) species (lower): Nd^{3+} (blue), NdNO_3^{2+} (black), NdL^{3+} (brown), NdL_2^{3+} (rose). Detailed titration conditions are in Table S3 of Supporting Information.

As shown in Table 3, the CF_3ABDMA complexes with both U(VI) and Nd(III) in aqueous solution are not strong and the U(VI) complex ($\log\beta_1 = 2.05$) is stronger than that of Nd(III) ($\log\beta_1 = 1.44$). These observations are corroborated by the ^{19}F NMR spectra shown in Figure S2 of the Supporting Information. The ^{19}F NMR data also show that the binding of CF_3ABDMA with UO_2^{2+} and La^{3+} becomes stronger in methanol solutions, which can be interpreted as the weaker solvation of metal ions in methanol than in water resulting in less energy required for desolvation. The ^{19}F NMR

spectra and more detailed descriptions are provided in the Supporting Information.

3.4 Complexation of Np(V) with CF₃ABDMA

One representative set of absorption spectra in the spectrophotometric titration of Np(V) with CF₃ABDMA is shown in Figure 4. As the concentration of CF₃ABDMA was increased, the intensity of the absorption band of free NpO₂⁺ around 980.4 nm decreased, and a “shoulder” band appeared at about 988.0 nm. Analysis with the HypSpec program indicates that the spectral changes can be described by the formation of the 1:1 Np(V)/CF₃ABDMA as represented by reaction (5). Using the HypSpec program, the stability constant of the complex was calculated to be 0.46±0.09. In the calculation, the protonation constant of CF₃ABDMA from this work and the stability constant of NpO₂(NO₃) (aq) from the literature^[36] were used.






Figure 4 Spectrophotometric titrations of NpO_2^+ complexation with CF_3ABDMA (upper) and calculated molar absorptivity of Np(V) species (lower): NpO_2^+ (blue), NpO_2NO_3 (brown), NpO_2L^+ (rose). Detailed titration conditions are in Table S3 of Supporting Information.

4. DISCUSSION

4.1 Thermodynamic trends

The stability constants of the complexes of CF_3ABDMA with UO_2^{2+} , Nd^{3+} , and NpO_2^+ from this work can be compared with the data of three structurally related diacetamide ligands, BnABDMA , ABDMA , and MABDMA , in the literature.^[16,17,18] The data are summarized in Table 4. Two trends can be extracted from these data and discussed as follows.

Table 4 Overall equilibrium constants for the complexation of CF₃ABDMA, in comparison with BnABDMA, ABDMA, and MABDMA, with UO₂²⁺, Nd³⁺, and NpO₂⁺ at 25 °C and I = 1.0 M NaNO₃

Metal ion	Ligand	Reaction	log β	Ref.
UO ₂ ²⁺	CF ₃ ABDMA	UO ₂ ²⁺ + L = UO ₂ L ²⁺	2.05 ± 0.12	p.w.
	BnABDMA	UO ₂ ²⁺ + L = UO ₂ L ²⁺	4.33 ± 0.03	[17]
		UO ₂ ²⁺ + 2L = UO ₂ L ₂ ²⁺	7.67 ± 0.06	
	ABDMA	UO ₂ ²⁺ + L = UO ₂ L ²⁺	5.03 ± 0.03	[17]
		UO ₂ ²⁺ + 2L = UO ₂ L ₂ ²⁺	9.08 ± 0.03	
	MABDMA	UO ₂ ²⁺ + L = UO ₂ L ²⁺	5.30 ± 0.03	[17]
UO ₂ ²⁺ + 2L = UO ₂ L ₂ ²⁺		9.66 ± 0.03		
Nd ³⁺	CF ₃ ABDMA	Nd ³⁺ + L = NdL ³⁺	1.44 ± 0.09	p.w.
		Nd ³⁺ + 2L = NdL ₂ ³⁺	1.90 ± 0.09	
	BnABDMA	Nd ³⁺ + L = NdL ³⁺	2.92 ± 0.09	[16]
		Nd ³⁺ + 2L = NdL ₂ ³⁺	5.08 ± 0.09	
		Nd ³⁺ + 3L = NdL ₃ ³⁺	7.13 ± 0.09	
	ABDMA	Nd ³⁺ + L = NdL ³⁺	4.08 ± 0.09	[16]
		Nd ³⁺ + 2L = NdL ₂ ³⁺	6.93 ± 0.09	
		Nd ³⁺ + 3L = NdL ₃ ³⁺	10.02 ± 0.9	
	MABDMA	Nd ³⁺ + L = NdL ³⁺	4.40 ± 0.09	[16]
Nd ³⁺ + 2L = NdL ₂ ³⁺		7.52 ± 0.36		
Nd ³⁺ + 3L = NdL ₃ ³⁺		10.50 ± 0.50		
NpO ₂ ⁺	CF ₃ ABDMA	NpO ₂ ⁺ + L = NpO ₂ L ⁺	0.46 ± 0.09	p.w.
	BnABDMA	NpO ₂ ⁺ + L = NpO ₂ L ⁺	2.90 ± 0.09	[18]
		NpO ₂ ⁺ + 2L = NpO ₂ L ₂ ⁺	4.01 ± 0.09	
	ABDMA	NpO ₂ ⁺ + L = NpO ₂ L ⁺	2.80 ± 0.09	[18]
		NpO ₂ ⁺ + 2L = NpO ₂ L ₂ ⁺	4.00 ± 0.09	
	MABDMA	NpO ₂ ⁺ + L = NpO ₂ L ⁺	3.59 ± 0.09	[18]
		NpO ₂ ⁺ + 2L = NpO ₂ L ₂ ⁺	5.50 ± 0.09	

4.1.1 Comparison between CF₃ABDMA, BnABDMA, ABDMA, and MABDMA

In Figure 5, the stability constants for the 1:1 complexes are plotted. For each metal ion, the binding strength of the four amine-bridged diacetamide ligands follows the trend MABDMA > ABDMA > BnABDMA > CF₃ABDMA, except for the complexation of Np(V) where ABDMA and BnABDMA show very similar binding strength. This trend is in accordance

with the decrease in the basicity of the ligands as reflected by their protonation constants (Table 2). Both the trifluoromethyl and the phenyl groups have the electron-withdrawing ability, which reduces the electron density on the central N atom of CF₃ABDMA and BnABDMA ligands, in comparison with ABDMA. As a result, the basicity of CF₃ABDMA and BnABDMA is less than that of ABDMA, resulting in weaker complexes with CF₃ABDMA and BnABDMA than those with ABDMA. The complexation of CF₃ABDMA with the metal ions is the weakest among the four amine-bridged diacetamides, implying that the electron-withdrawing ability of the trifluoromethyl group is even stronger than that of the phenyl group. On the contrary, the methyl group has the electron-donating ability, which makes MABDMA a more basic ligand. Therefore, the metal complexes with MABDMA are stronger than those with ABDMA, and are the strongest among the complexes with all four diacetamides.




Figure 5. Thermodynamic trends in the complexation of amine-bridged diacetamide ligands with UO_2^{2+} , Nd^{3+} , and NpO_2^+ . Data are for $t = 25\text{ }^\circ\text{C}$ and $I = 1.0\text{ M NaNO}_3$

4.1.2 Comparison between UO_2^{2+} , Nd^{3+} , and NpO_2^+

Figure 5 also shows that, for each ligand, the stability constants of the 1:1 complexes follow the order: $\text{UO}_2^{2+} > \text{Nd}^{3+} > \text{NpO}_2^+$. This agrees with the order of the effective charges on the metal ions: $+3.2 (\text{UO}_2^{2+}) > +3.0 (\text{Nd}^{3+}) > +2.2 (\text{NpO}_2^+)$.^[37] Such agreement implies that the complexation of the amine-bridged diacetamides with actinide and lanthanide ions is predominantly electrostatic in nature.

4.2 Coordination modes

4.2.1 Single crystal structure of $\text{UO}_2\text{L}_2(\text{ClO}_4)_2$

In a previous study^[17] of U(VI) complexation with MABDMA, ABDMA, and

BnABDMA, EXAFS and computations suggested that amine-bridged diacetamides form tridentate complexes with U(VI) via the amine N atom and the two amide O atoms. No crystal structures of the U(VI)/diacetamide complexes were obtained in previous studies.

The structure of $\text{UO}_2\text{L}_2(\text{ClO}_4)_2$ (Figure 6) obtained in this work is the first single crystal structure of a U(VI)/diacetamide complex. The structural data has been deposited as CCDC 1584249 in the Cambridge Crystallographic Data Centre (<https://www.ccdc.cam.ac.uk/>). The perchlorate salt of the dicationic 1:2 U(VI) complex with CF_3ABDMA was found to crystallize in the monoclinic space group $P2_1/c$, with the uranium atom lying directly on a crystallographic inversion center such that one half of the complex generates the other by symmetry. Akin to the other uranyl complexes of this class of diacetamide ligands, CF_3ABDMA binds to the U(VI) center as a neutral species in a tridentate fashion.^[13,17] A direct comparison of the bonding parameters of the U(VI)/ CF_3ABDMA complex with those of the previously reported U(VI) complex of the diacetamide chelator TMDGA, which contains an ether (O) linkage in place of the functionalized amine nitrogen ($\text{N-CH}_2\text{-CF}_3$) found in CF_3ABDMA , can provide a basis for rationalizing the relative donor properties of these two ligands. The relevant crystallographically determined bond lengths for both complexes are given in Table 5. While a comparison of the $\text{U-N}_{\text{amine}}$ and $\text{U-O}_{\text{ether}}$ distances in these two complexes would be imprudent, the shorter U-

O_{amide} and longer $U-O_{\text{uranyl}}$ distances in the CF_3ABDMA complex indicate that despite the diminished basicity of its amine group, CF_3ABDMA is still a stronger ligand for U(VI) than TMDGA. This conclusion, as derived from the crystallographic data, aligns with the measured formation constants for the complexes as presented in this work and in previous work describing the TMDGA ligand.^[13]




Figure 6. X-ray crystal structure of $\text{UO}_2\text{L}_2(\text{ClO}_4)_2$ with 50% probability ellipsoids. Perchlorate ions and hydrogen atoms are omitted for clarity.

Table 5. Selected bond distances (\AA) in $\text{UO}_2\text{L}_2(\text{ClO}_4)_2$ where L stands for CF_3ABDMA in this work or TMDGA. Atom labels align to those given in Figure 4, with the corresponding functionalities specified in parentheses.

Bond	CF_3ABDM	TMDGA ^[13]
	A	
U1-O1 (amide)	2.335(3)	2.416(4)
U1-O2 (amide)	2.386(3)	2.421(3)
U1-O3 (uranyl)	1.766(2)	1.752(3)
U1-N2 (amine) for CF_3ABDMA or U1-O (ether) for TMDGA	2.881(3)	2.614(3)

It needs some explanations why the 1:2 U(VI)/CF₃ABDMA complex was obtained in crystal form but not identified in aqueous solution by potentiometry or spectrophotometry. We believe that the formation of UO₂L₂(ClO₄)₂ is favored at higher pH (it in fact crystalized from a solution of pH ~6, see section 2.5), but complications with the hydrolysis of U(VI) make it very difficult to observe and determine the formation constant of UO₂L₂(ClO₄)₂ in aqueous solution of high pH.



It is known that the hydrolysis of metal ions in a mixture of water/ethanol could be substantially suppressed, due to the much lower dissociation constants ($[H^+][OH^-]$) of such media than that of pure water. Therefore, we hypothesized that the 1:2 U(VI)/CF₃ABDMA complex in solution could probably be observed in a mixture of water/ethanol. To test the validity of this hypothesis, ESI-MS experiments were conducted by spraying the U(VI)/ CF₃ABDMA solution (< 5% water in ethanol) into the instrument. Indeed, the 1:2 U(VI) complex with CF₃ABDMA was observed as a doubly-charged species, UO₂L₂²⁺, by ESI-MS experiments. Detailed results from ESI-MS were provided in Supporting Information (Figure S2).

4.2.2 DFT calculations

It has been previously¹⁷ shown that for the UO₂²⁺/diacetamide complexes, two types of minima of similar energy exist, differing in the conformation of the diacetamide ligand: Type *a* with a “distorted” O-C··C-O dihedral angle ~ 30° and Type *b* with a “planar” O-C··C-O dihedral angle ~ 0° (see

Figure 5 of the reference¹⁷). In this work, for simplicity, only the complexes with “distorted” ligands were computed.

DFT calculations were performed for the 1:1 and 1:2 $\text{UO}_2^{2+}/\text{CF}_3\text{ABDMA}$ complexes by following the same stepwise approach that was detailed in a previous study including optimization of structures, calculation of minimum energy of the complexes in gas phase and water, and calculation of the free energy for theoretical complexation reactions (6) and (7) in water. The calculations also provide the ESP charges on the donor atoms of CF_3ABDMA in the free and complexed ligand that help to explain the binding strength of CF_3ABDMA in comparison with other diacetamide ligands previously studied.



The theoretically calculated free energy for reactions (6) and (7) are shown in Table 6. Other computational results, including structure optimization, minimum energy of the complexes in gas phase and water, and the ESP charges on the donor atoms are provided in Supporting Information as Figure S3, Table S4 and Table S5.

Table 6 Theoretically calculated free energies (ΔG_{water}) for complex formation ($\text{kcal}\cdot\text{mol}^{-1}$) for reactions (6) and (7), in comparison with the experimental equilibrium constants of the formation of U(VI)/diacetamide complexes. $\Delta\log K_{\text{calc}} = \log K_{\text{calc}}(\text{ligand}) - \log K_{\text{calc}}(\text{BnABDMA})$, $\Delta\log K_{\text{exp}} =$

$\log K_{\text{exp}}$ (ligand) - $\log K_{\text{exp}}$ (BnABDMA).

Reaction	Ligand	ΔG_{water}	$\log K_{\text{exp}}$	$\Delta \log K_{\text{calc}}$	$\Delta \log K_{\text{exp}}$	Ref.
(6)	CF ₃ ABDMA	-14.3	2.05	-3.7	-2.3	This work
	BnABDMA	-19.3	4.33	0	0	17
	ABDMA	-19.0	5.03	-0.2	0.7	17
	MABDMA	-21.0	5.30	1.2	1	17
(7)	CF ₃ ABDMA	-17.4	(1.8) ^a	-1.5	(-1.5) ^a	This work
	BnABDMA	-19.5	3.34	0	0	17
	ABDMA	-20.1	4.05	0.4	1.4	17
	MABDMA	-20.0	4.37	0.4	2	17

^aEstimated value from the calculated ΔG_{water} in this work.

As shown in Table 6, the calculated free energy for the formation of 1:1 U(VI)/CF₃ABDMA complex (-14.3 kcal·mol⁻¹) is much less negative than those of the other diacetamide complexes (from -19.0 to -21.0 kcal·mol⁻¹), in good agreement with the order of the experimentally determined equilibrium constants. The values of $\Delta \log K$ in Table 6 are relative values for each ligand with respect to those of $\log K$ for BnABDMA. The correlation between the experimental and calculated $\Delta \log K$ for the 1:1 U(VI)/diacetamide complexes is fairly good as shown in Figure S4 in the Supporting Information.

The ESP charges for free CF₃ABDMA ligand and the U(VI) complexes are shown in Table S5. These data could be compared with those for the other three diacetamide ligands previously calculated at the same level of theory.¹⁷ As observed in the previous study,¹⁷ the calculated results generally indicate that the relative stability of the complexes can't be

predicted in detail by the charge on the central nitrogen, but also solvation and steric effects due to the substituents are the factors affecting the stability of the complex formed.

In brief, DFT calculations confirm that CF₃ABDMA is the weakest ligand among the four studied diacetamides. The calculated free energy for the formation of 1:2 U(VI)/CF₃ABDMA complex (-17.4 kcal·mol⁻¹) is also less than those of the other diacetamide complexes (from -19.5 to -20.1 kcal·mol⁻¹). In the absence of experimentally determined equilibrium constant for the formation of 1:2 U(VI)/CF₃ABDMA complex, we could, using the trends of values for reaction (7) in Table 6, estimate that the logK_{exp} for the stepwise formation of 1:2 U(VI)/CF₃ABDMA complex is about 1.8.

5. CONCLUSION

A new member of the series of amine-bridged diacetamide ligands, 2,2'-(trifluoroazanediy)bis(*N,N'*-dimethylacetamide) (CF₃ABDMA) was synthesized and its complexation with three metal ions, UO₂²⁺, Nd³⁺, and NpO₂⁺, was studied with thermodynamic, structural, and computational techniques. With each of the three metal ions, CF₃ABDMA was found to form the weakest complexes among the four amide-bridged diacetamides, presumably due to the strong electron withdrawing character of the CF₃- group that reduces the basicity of the amine

nitrogen. The binding strength of the four diacetamide ligands with UO_2^{2+} , Nd^{3+} , and NpO_2^+ follows the order of the effective electronic charge on the metal ions, indicating the complexation of amine-bridged diacetamides with these cations is predominantly electrostatic interactions.

Acknowledgement. The thermodynamic measurements, ESI-MS experiments, and X-ray crystallographic work are supported by the U.S. Department of Energy (DOE), Office of Science, Office of Basic Energy Sciences, under Contract No. DE-AC02-05CH11231 at LBNL. Single-crystal X-ray diffraction data were collected and analysed at the Advanced Light Source (ALS) that is supported by the Director, Office of Science, Office of Basic Energy Sciences, U.S. DOE under Contract No. DE-AC02-05CH11231. The collection of NMR data is supported by the UC Berkeley NMR facility, funded in part by NSF grants CHE 9633007, CHE 82-08992, and NIH grants 1S10RR016634-01, RR 02424A-01. Y. Gao acknowledges financial support from the China Scholarship Council for her visit to LBNL. T. D. Lohrey thanks the U.S. DOE Integrated University Program for a graduate research fellowship. A. Melchior acknowledge the support for the computational studies by the “Piano Strategico d’Ateneo 2016-18” Program at University of Udine, Italy.

REFERENCES

- (1) K. L. Nash, G. R. Choppin, Separations Chemistry for Actinide Elements: Recent Developments and Historical Perspective, *Sep. Sci. Technol.* **1997**, 32 (1-4), 255-274.
- (2) D. D. Sood, S. K. Patil, Chemistry of Nuclear Fuel Reprocessing: Current Status, *J. Radioanal. and Nucl. Chem.* **1996**, 203 (2), 547-573.
- (3) *Yucca Mountain Science and Engineering Report Rev. 1*; U. S. Department of Energy: North Las Vegas, 2002.
- (4) H. Suzuki, Y. Sasaki, Y. Sugo, A. Apichaibukol, T. Kimura, Extraction and Separation of Am(III) and Sr(II) by N,N,N',N'-Tetraoctyl-3-Oxapentanediamide (TODGA), *Radiochim. Acta* **2004**, 92 (8), 463-466.
- (5) S. A. Ansari, P. N. Pathak, M. Husain, A. K. Prasad, V. S. Parmar, V. K. Manchanda, Extraction of Actinides Using N,N,N',N'-Tetraoctyl Diglycolamide (TODGA): A Thermodynamic Study, *Radiochim. Acta* **2006**, 94 (6-7), 307-312.
- (6) G. Tian, P. Zhang, J. C. Wang, L. Rao, Extraction of Actinide(III, IV, V, VI) Ions and TcO_4^- by N,N,N',N'-Tetraisobutyl-3-Oxa-Glutaramide, *Solvent Extr. Ion Exch.* **2005**, 23 (5), 631-643.
- (7) Y. Sasaki, G. R. Choppin, Extraction of Np(V) by N,N'-Dimethyl-N,N'-Dihexyl-3-Oxapentanediamide, *Radiochim. Acta* **1998**, 80 (2), 85-88.

- (8) S. A. Ansari, P. Pathak, P. K. Mohapatra, V. K. Manchanda, Chemistry of Diglycolamides: Promising Extractants for Actinide Partitioning, *Chem. Rev.* **2012**, *112* (3), 1751–1772.
- (9) S. Nave, G. Modolo, C. Madic, F. Testard, Aggregation Properties of *N, N, N', N'*-Tetraoctyl-3-oxapentanediamide (TODGA) in *n*-Dodecane, *Solvent Extr. Ion Exch.* **2004**, *22* (4), 527–551.
- (10) H. Narita, T. Yaita, S. Tachimori, Extraction of Lanthanides with *N, N'*-Dimethyl-*N, N'*-diphenyl-malonamide and -3,6-dioxaoctanediamide, *Solvent Extr. Ion Exch.* **2004**, *22* (2), 135–145.
- (11) A. Shimada, T. Yaita, H. Narita, S. Tachimori, K. Okuno, Extraction Studies of Lanthanide(III) Ions with *N, N'*-Dimethyl-*N, N'*-diphenylpyridine-2,6-dicarboxamide (DMDPhPDA) from Nitric Acid Solutions, *Solvent Extr. Ion Exch.* **2004**, *22* (2), 147–161.
- (12) G. Tian, J. Xu, L. Rao, Optical Absorption and Structure of a Highly Symmetrical Neptunium(v) Diamide Complex, *Angew. Chemie - Int. Ed.* **2005**, *44* (38), 6200–6203.
- (13) G. Tian, L. Rao, S. J. Teat, G. Liu, Quest for Environmentally Benign Ligands for Actinide Separations: Thermodynamic, Spectroscopic, and Structural Characterization of U^{VI} complexes with Oxa-Diamide and Related Ligands, *Chem. - A Eur. J.* **2009**, *15* (16), 4172–4181.
- (14) L. Rao, G. Tian, S. J. Teat, Complexation of Np(V) with *N,N*-Dimethyl-3-Oxa-Glutaramic Acid and Related Ligands: Thermodynamics,

- Optical Properties and Structural Aspects, *Dalton Trans.* **2010**, 39 (13), 3326–3330.
- (15) G. Tian, S. J. Teat, L. Rao, Structural and Thermodynamic Study of the Complexes of Nd(III) with N,N,N',N'-Tetramethyl-3-Oxa-Glutaramide and the Acid Analogues, *Inorg. Chem.* **2014**, 53 (18), 9477–9485.
- (16) P. V. Dau, Z. Zhang, P. D. Dau, J. K. Gibson, L. Rao, Thermodynamic Study of the Complexation between Nd³⁺ and Functionalized Diacetamide Ligands in Solution, *Dalton. Trans.* **2016**, 45 (30), 11968–11975.
- (17) P. V. Dau, Z. Zhang, Y. Gao, B. F. Parker, P. D. Dau, J. K. Gibson, J. Arnold, M. Tolazzi, A. Melchior, L. Rao, Thermodynamic, Structural, and Computational Investigation on the Complexation between UO₂²⁺ and Amine-Functionalized Diacetamide Ligands in Aqueous Solution, *Inorg. Chem.* **2018**, 57 (4), 2122–2131.
- (18) Y. Gao, P. V. Dau, B. F. Parker, J. Arnold, A. Melchior, Z. Zhang, L. Rao, Complexation of NpO₂⁺ with Amine-Functionalized Diacetamide Ligands in Aqueous Solution: Thermodynamic, Structural, and Computational Studies, *Inorg. Chem.* **2018**. DOI: 10.1021/acs.inorgchem.8b00654.
- (19) J. A. Dean, Analytical Chemistry Handbook; McGraw-Hill, Inc.: New York, **1995**, 3-108.
- (20) L. Rao, T. G. Srinivasan, A. Yu Garnov, P. Zanonato, P. Di Bernardo, A.

- Bismondo, Hydrolysis of neptunium(V) at variable temperatures (10-85 °C), *Geochim. Cosmochim. Acta* **2004**, 68 (23), 4821-4830.
- (21) G. Gran, Determination of the Equivalence Point in Potentiometric Titrations, Part II, *Analyst* **1952**, 77 (920), 661.
- (22) P. Gans, A. Sabatini, A. Vacca, Investigation of Equilibria in Solution. Determination of Equilibrium Constants with the HYPERQUAD Suite of Programs, *Talanta* **1996**, 43 (10), 1739-1753.
- (23) Apex2. Bruker Analytical X-ray Systems Inc.: Madison, WI, USA 2003.
- (24) SAINT: SAX Area-Detector Integration Program v7.60a, Bruker Analytical X-ray Systems, Inc., Madison, WI, 2010.
- (25) G. M. Sheldrick, A Short History of SHELX. *Acta Cryst. A* **2008**, 64 (1), 112-122.
- (26) A. D. Becke, A New Mixing of Hartree-Fock and Local Density-Functional Theories. *J.Chem.Phys.* **1993**, 98 (2), 1372-1377.
- (27) C. T. Lee, W. T. Yang, R. G. Parr, Development of the Colle-Salvetti Correlation-Energy Formula Into A Functional of the Electron-Density. *Phys.Rev.B* **1988**, 37 (2), 785-789.
- (28) P. Di Bernardo, P. Zanonato, F. Benetollo, A. Melchior, M. Tolazzi, L. Rao, Energetics and Structure of Uranium(VI)-Acetate Complexes in Dimethyl Sulfoxide. *Inorg. Chem.* **2012**, 51 (16), 9045-9055.
- (29) F. Endrizzi, A. Melchior, M. Tolazzi, L. Rao, Complexation of Uranium(VI) with Glutarimidoxime: Thermodynamic and

- Computational Studies. *Dalt. Trans.* **2015**, 44 (31), 13835–13844.
- (30) J. Tomasi, B. Mennucci, R. Cammi, Quantum Mechanical Continuum Solvation Models. *Chem. Rev.* **2005**, 105 (8), 2999–3093.
- (31) S. Vukovic, B. P. Hay, V. S. Bryantsev, Predicting Stability Constants for Uranyl Complexes Using Density Functional Theory. *Inorg. Chem.* **2015**, 54 (8), 3995–4001.
- (32) M. J. G. Frisch, W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, Gaussian 16, Rev. A.03. *Inc.: Wallingford, CT.* 2016.
- (33) L. Rao, G. Tian, Thermodynamic Study of the Complexation of Uranium(VI) with Nitrate at Variable Temperatures, *J. Chem. Thermodynamics* **2008**, 40 (6), 1001–1006.
- (34) P. Zanonato, P. Di Bernardo, A. Bismondo, G. K. Liu, X. Y. Chen, L. Rao, Hydrolysis of Uranium(VI) at Variable Temperatures (10–85 °C). *J. Am. Chem. Soc.* **2004**, 126 (17), 5515–5522.
- (35) L. Rao, G. Tian, Complexation of Lanthanides with Nitrate at Variable Temperatures: Thermodynamics and Coordination Modes. *Inorg. Chem.* **2009**, 48 (3), 964–970.
- (36) P. R. V. Rao, N. M. Gudi, S. V. Bagawde, S. K. Patil, The complexing of Np(V) by some inorganic ligands, *J. Inorg. Nucl. Chem.* **1979**, 41 (2),

235-239.

- (37) G. R. Choppin, L. F. Rao, Complexation of Pentavalent and Hexavalent Actinides by Fluoride. *Radiochim. Acta* **1984**, 37 (3), 143-146.

Graphic Content

CF₃ABDMA ligand coordinates to UO₂²⁺, Nd³⁺, and NpO₂⁺ in a tridentate mode, through the center nitrogen and two amide oxygen atoms.

