UC Davis UC Davis Previously Published Works

Title

Visual and verbal working memory loads interfere with scene-viewing

Permalink https://escholarship.org/uc/item/9st626hc

Journal Attention, Perception, & Psychophysics, 82(6)

ISSN 1943-3921

Authors

Cronin, Deborah A Peacock, Candace E Henderson, John M

Publication Date 2020-08-01

DOI

10.3758/s13414-020-02076-1

Peer reviewed



HHS Public Access

Atten Percept Psychophys. Author manuscript; available in PMC 2021 August 01.

Published in final edited form as:

Atten Percept Psychophys. 2020 August; 82(6): 2814–2820. doi:10.3758/s13414-020-02076-1.

Visual and Verbal Working Memory Loads Interfere with Scene-Viewing

Deborah A. Cronin, Candace E. Peacock

Author manuscript

Center for Mind and Brain, University of California, Davis;

John M. Henderson

Center for Mind and Brain & Department of Psychology, University of California, Davis.

Abstract

Working memory is thought to be divided into distinct visual and verbal subsystems. Studies of visual working memory frequently use verbal working memory tasks as control conditions and/or use articulatory suppression to ensure visual load is not transferred to verbal working memory. Using these verbal tasks relies on the assumption that the verbal working memory load will not interfere with the same processes as visual working memory. In the present study, participants maintained a visual or verbal working memory load as they simultaneously viewed scenes while their eye movements were recorded. Because eye movements and visual working memory are closely linked, we anticipated the visual load would interfere with scene-viewing (and vice versa), while the verbal load would not. Surprisingly, both visual and verbal memory loads interfered with scene-viewing behavior, while eye movements during scene-viewing did not significantly interfere with performance on either memory task. These results suggest that a verbal working memory load can interfere with eye movements in a visual task.

Keywords

eye movements; scene-viewing; visual working memory; verbal working memory; attention

Daily tasks often require attending to the visual world while simultaneously maintaining verbal information in memory. We might mentally rehearse a phone number while searching our surroundings for a pen or maintain a grocery list while scanning shelves at the store. These tasks require us to both move our eyes and engage working memory.

Working memory (WM) is thought to consist of independent, limited-capacity, visual and verbal subsystems (Baddeley & Hitch, 1974; Baddeley, 1986; Luck & Vogel, 1997; Smith, et al., 1996). For this reason, verbal WM tasks are frequently used as control conditions for

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. http://www.springer.com/gb/open-access/authors-rights/aam-terms-v1

Correspondence concerning this article should be addressed to Deborah Cronin, Center for Mind and Brain, University of California Davis, 267 Cousteau Pl, Davis, CA 95618. dacronin@ucdavis.edu.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of a an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

dual-task paradigms investigating visual WM and its related processes (e.g., Cronin & Irwin, 2018; Hollingworth, Richard, & Luck, 2008). Further, articulatory suppression is frequently used in conjunction with visual WM tasks to prevent participants from offloading visual load into verbal memory (e.g., Schmidt, Vogel, Woodman, & Luck, 2002). Using verbal WM tasks as a control relies on the assumption that verbal WM load will minimally interfere with visual tasks.

Contrary to this assumption, there is some evidence that verbal WM can interfere with attention control mechanisms like distractor suppression (Lavie, Hirst, deFockert, & Viding, 2004; de Fockert, 2013) and attentional guidance (Soto & Humphreys, 2007, 2008). Given this cross-modal interference, an interesting and open question is whether a verbal WM load can change eye movements during scene-viewing. Saccadic eye movements are thought to be closely related to visual WM (e.g., Cronin & Irwin, 2018; Hollingworth, Richard, & Luck, 2008) and the contents of visual WM can influence scene-viewing behavior (Bahle, Beck, & Hollingworth, 2018). However, no such relationship has been demonstrated between verbal WM load and scene-viewing behavior. If such a relationship exists, it would add to the evidence cautioning the use of verbal WM tasks and articulatory suppression as control conditions for visual WM tasks.

The Present Study

The present study examined the effects of WM load on eye movements during sceneviewing and the effects of eye movements during scene-viewing on WM performance. Participants maintained a verbal or visual WM load while free-viewing scenes. The same participants also viewed scenes without a memory load (Figure 1). A separate set of participants performed the visual and verbal WM tasks without free-viewing the scenes. Given the literature outlined above, we predicted the visual WM and scene-viewing tasks would interfere with each other under dual-task conditions, while the verbal WM task and scene-viewing task would not. In contrast with previous work, we instead found interference from both the visual and verbal WM tasks on eye movements during scene-viewing and no evidence that eye movements during scene-viewing interfered with either WM task.

Method

Participants

Fifty-eight experimentally naïve participants from the University of California, Davis participated for course credit. Four participants were replaced due to poor eye-tracking (25% data loss) and six were replaced due to below-chance memory task performance. Forty-eight participants' (31 female; mean age = 20.6y, 10.1% mean data loss) data are included in the subsequent analyses.

A separate group of 59 participants recruited from the same pool participated in a memoryonly condition. Eleven subjects were dropped due to below-chance memory task performance. Forty-eight (38 female, mean age = 20.9y) participants' data are included in the analyses.

Apparatus and Stimuli

Dual-Task and Scene-Viewing Conditions.—Participants were seated 85cm from a 24.5" LCD display. Head movements were limited by chin and forehead rests. Scenes were displayed at a resolution of 1024×768 px and subtended $26.5 \times 20^{\circ}$. Eye movements were monitored with a tower-mounted EyeLink 1000+ eye-tracker sampling the right eye at 1000Hz (SR Research, 2010). Experimental stimuli were presented using the SR Research Experiment Builder software (Version 2.1.512). Each participant underwent a 9-point calibration procedure before each block and continued on to the experimental trials once the eye-tracker's average and maximum error were minimized to >0.5° and 0.99°, respectively.

Ninety-six real-world scenes were presented during the scene-viewing task. Scenes were originally collected from online image searches. Scenes were luminance-matched by converting each to LAB color space (0=darkest, 100=brightest), scaling the luminance channel of all scenes from 0 to 1, then adjusting them to the set's average luminance (M= 0.46 L). Each scene was presented once over the course of the experiment, with 24 scenes appearing in a random order in each condition. Scenes appeared equally in each condition across participants.

Working Memory Control.—Participants were seated 85cm from a 24.5" LCD display or a 21" CRT display. Head movements were limited by chin and forehead rests.

Procedure

Dual-Task and Scene-Viewing Conditions.—There were four experimental tasks: visual WM, verbal WM, and visual and verbal control conditions (Figure 1). Participants completed 24 trials in each task. The four tasks were presented in separate blocks in a counterbalanced order across participants.

During the visual WM task, five colored squares were presented for 250ms. Colors were chosen from 9 possible colors ([RGB: 255,0,0], [255,255,0], 255,124,27], [0,255,128], [0,128,0], [0,165,253], [0,0,255], [255,0,255], [128,0,128]). Squares subtended $0.4^{\circ} \times 0.4^{\circ}$ and were presented on an invisible circle of radius 0.4° centered on the screen. After the memory array disappeared, the fixation cross remained for 500ms until scene onset. The scene remained for 3500ms. Participants were instructed to free-view the scenes. After scene offset, five colored squares appeared and remained visible until response. On 50% of trials, one of the five squares' color was changed to a color not used in the original array. Participants responded whether or not a color changed.

During the verbal WM task, seven consonants were sequentially presented at fixation. Each consonant subtended 0.29° and was presented for 300ms. Participants were instructed to verbalize and rehearse the letters throughout the trial. After the last letter offset, a checkerboard mask was presented for 250ms, then a fixation cross for 250ms. Then, a scene appeared for 3500ms. After scene offset, a single letter was presented at central fixation. Participants responded whether or not that letter was one of the seven they held in memory.

The visual and verbal control tasks proceeded in the same way as the visual and verbal WM tasks, but participants were not asked to respond to the memory items.

Working Memory Control.—A second group of participants completed the memory tasks as outlined above without interleaved scenes. The delay interval consisted of a gray screen with a fixation cross at center. The task order was counterbalanced across participants. A previous study found these visual and verbal WM single tasks were matched in difficulty (Cronin & Irwin, 2018).

Data Processing

Fixations and saccades were segmented with EyeLink's standard algorithm using velocity and acceleration thresholds (30° /s and 9500° /s²; SR Research, 2010). Eye-tracking data files were converted to ASCII format using the EDFConverter tool (SR Research, 2010) and imported into Matlab (MathWorks, Inc., 2018a). Off-screen fixations, saccade amplitudes >27°, and the first fixation on the scene, always located at the center of the display due to the pre-scene fixation-marker, were eliminated from analysis. Statistical analyses were performed in R (R Core Team, 2019).

Results

Eye Movement Measures

To determine whether visual and verbal WM loads interfere with participants' scene-viewing behavior, we employed linear mixed effect (LME) models with memory load (load vs. no load) and load type (visual vs. verbal) as fixed effects and subject and scene as random effects. All p-values reported below are adjusted to correct for family-wise error using the Holm-Bonferroni method. The reported effect sizes for LMEs were calculated according to Westfall and colleagues (2014; Brysbaert & Stevens, 2018). LMEs were completed using the R package lme4 (Bates, Maechler, Bolker, & Walker, 2015). Means and standard deviations for each measure are reported in Table 1.

First, we tested whether a WM load influenced fixation-level behavior. Participants with a memory load made fewer fixations than participants without a memory load ($\chi^2(1) = 485.25$, p < 0.001, d = 0.37). The main effect of load-type on the number of fixations was also significant ($\chi^2(1) = 31.71$, p < 0.001, d = 0.33), as was the interaction between load and load-type ($\chi^2(1) = 53.41$, p < 0.001, d = 0.37). Participants made fewer eye movements in the visual load condition than in the verbal load condition but moved their eyes a similar number of times in the two no-load conditions.

There were also effects of load on fixation durations and saccade amplitudes, with participants making longer fixations ($\chi^2(1) = 43.33$, p < 0.001, d = 0.07) and shorter saccades ($\chi^2(1) = 31.71$, p < 0.001, d = 0.33) with a memory load compared to without. The main effect of load-type was not significant for either (fixation duration: $\chi^2(1) = 0.82$, p = 1.00, d = 0.13, $BF_{JZS} = 0.75 + -1.84\%$; saccade amplitude: $\chi^2(1) = 3.33$, p = 1.00, d = 0.26, $BF_{JZS} = 0.17 + -1.97\%$), but the interaction between load and load-type was significant for both fixation duration ($\chi^2(1) = 14.49$, p = 0.002, d = 0.21) and saccade amplitude ($\chi^2(1) = 69.97$, p < 0.001, d = 0.42). Participants' fixations were longer and their saccades were shorter with a visual load than with a verbal load.

Next, we assessed how a memory load influenced scene-viewing over the course of the entire 3.5s viewing period. We examined participants' scan path length, the percent of the scene fixated (Castelhano, Mack, and Henderson, 2009), and the dispersion of fixations from the center of the screen (Anliker, 1976). The main effects of load and load-type were significant for all three measures of fixation spread, as was the interaction between load and load-type (Table 2). Participants' scan paths were shorter, they fixated a smaller percentage of the scene, and their fixations were less dispersed from center with a memory load compared to without a memory load. As evidenced by the interactions, this effect was more pronounced in the visual load condition than in the verbal load condition (Figure 2).

Working Memory Measures

Participants in the dual-task experiment performed well on both the visual (accuracy M = 71.18%, SD = 10.09) and verbal (M = 76.65%, SD = 8.92) memory tasks. The difference in performance was significant (t(94) = 22122.813, p = 0.006, d = 0.57), suggesting the verbal task was easier than the visual task.

To assess whether participants' WM performance was influenced by the concurrent sceneviewing task, a separate group of 48 participants completed just the WM tasks. Their performance on the visual (M= 72.66%, SD= 8.72) and verbal (M= 76.04%, SD= 9.86) tasks were statistically similar to dual-task performance (visual: t(94) = 0.767, p = 0.445, d = 0.16, $BF_{JZS} = 3.59$; verbal: t(94) = 0.316, p = 0.752, d = 0.06, $BF_{JZS} = 4.46$). Whereas the memory tasks influenced eye movements in the scene-viewing task, the scene-viewing task did not significantly affect memory task performance.

General Discussion

In this study, participants viewed scenes while maintaining a WM load or with no WM load. The WM load was visual (remember five colors) or verbal (remember seven letters). We saw profound differences in how overt attention moved through scenes when participants maintained a memory load regardless of load modality. Visual and verbal WM loads influenced when participants moved their eyes (fixation durations) and where they moved them (saccade amplitudes, scan paths, percent fixated, dispersion; Figure 2). We found no differences between dual- and single-task performance on the WM tasks.

Effects of Working Memory Load on Eye Movements

Previous work has shown visual WM and eye movement control are closely linked (e.g., Bahle, et al., 2018; Cronin & Irwin, 2018; Irwin, 1992a, 1992b; Hollingworth & Luck, 2008; Hollingworth, et al., 2008; Tas, Luck, & Hollingworth, 2016). Given this literature, it is unsurprising that a visual WM load influenced eye movements during scene-viewing in the present study. However, it is surprising that a *verbal* memory load also interfered with eye movements during scene-viewing. Our results caution the use of verbal tasks as a control for studies of attention and visual WM: the verbal task may affect the baseline patterns of eye movement behavior despite relying on a different subsystem of WM.

There are many possible reasons why the verbal WM task interfered with overt attention in our study. Recently, the long-held assumption of independent verbal and visual WM

subsystems has been challenged. Studies have found interference between concurrent visual and verbal WM tasks (Allen, Baddeley, & Hitch, 2006; Bae & Luck, 2018; Hardman, Vergauwe, & Ricker, 2017; Makovski, Shim, & Jiang, 2006; Morey & Beiler, 2013; Morey & Cowan, 2004; Ricker, Cowan, & Morey, 2010; Saults & Cowan, 2007; Vergauwe, Barrouillet, & Camos, 2010) and evidence that verbal WM contents can influence visual attention (Soto & Humphreys, 2007, 2008). In the perceptual load literature, a verbal WM load has been shown to interfere with distractor suppression, another function of attention (Lavie, et al., 2004; de Fockert, 2013). In line with these literatures, our WM tasks may have taxed both their modality-specific WM subsystem and a more general resource related to overt attention control such as the central executive (Baddeley & Hitch, 1974).

Because our verbal stimuli were presented visually, it is possible our verbal WM task taxed visual WM to some extent, leading to the interference between the verbal WM task and overt attention. There is evidence that visual information is filtered through visual WM even while it remains on screen (Tsubomi, Fukuda, Watanabe, & Vogel, 2013). We took many precautions to discourage persisting visual representations of our letter stimuli: letters were presented sequentially in the same spatial location and the last letter was followed by a mask. Participants were also instructed to verbalize and rehearse the letter stimuli throughout the trial. While it remains possible that participants used visual WM to support their verbal WM task performance, the verbal WM task we employed in this study is similar to others commonly used in the literature (e.g., Cronin & Irwin, 2018; Lavie & deFockert, 2005). Even articulatory suppression tasks frequently use visually presented stimuli (e.g., Luck & Vogel, 1997;. Peterson et al., 2019; Schmidt et al., 2002). Our results suggest that these verbal WM tasks may interfere with performance on visual tasks.

Finally, participants may have been less engaged in the free-viewing task when they had to simultaneously maintain a WM load. When attention is focused on internal information (i.e., a WM load), attention to external information suffers (Kiyonaga & Egner, 2013; Buetti & Lleras, 2016). Further, when engagement is high in a given task, changes to or onsets of task-unrelated stimuli go unnoticed (e.g., Fougnie & Marois, 2007; Mack & Rock, 1998; Most et al., 2001; Neisser, 1979; Neisser & Becklen, 1975; Simons & Chabris, 1999). Here, participants may have been relatively insensitive to the scene stimuli while maintaining a WM load. The similar WM performance under dual- and single-task conditions fits with this interpretation, as does the difficulty level of our two WM tasks: the verbal task was slightly easier than the visual task, and impeded eye movements less. However, task difficulty does not necessarily equate to task-engagement (Buetti & Lleras, 2016).

Regardless of the mechanism behind the verbal WM task's unexpected interference with overt attention, our results stand: a commonly used verbal WM task interfered with overt attention. Future researchers should take this possibility into account when designing their studies. If a verbal WM task or articulatory suppression will be used, it should be used across all conditions to ensure the effect of the verbal WM load can be accounted for.

Effects of Eye Movements on Working Memory

Aside from the effect of verbal WM load on overt attention, it is also interesting and surprising that visual WM task performance was equivalent under dual- and single-task

conditions. Recent evidence suggests that the target of an eye movement is automatically placed in visual WM, overwriting items already in WM (Cronin & Irwin, 2018; Hollingworth, et al., 2008; Tas et al., 2016). Menneer and colleageus (2019) similarly found visual WM impairment when participants were simultaneously engaged in a search task. The present results contrast with these findings. Participants maintaining a visual WM load during scene-viewing made around seven eye movements during the 3.5s delay period. However, their WM task performance was comparable to participants who made no eye movements in the single-task condition. Therefore, we find no evidence that participants' eye movements under dual-task conditions impaired their WM performance.

Conclusions

Our results suggest visual *and* verbal WM loads impede normal scene-viewing behavior, influencing both the timing and spread of eye movements through a scene. This finding cautions the use of verbal WM tasks and articulatory suppression as an effective control condition for visual WM tasks. Researchers should account for this effect of verbal WM loads when designing experiments. Our results also suggest eye movements may not always tax visual WM, a stark contrast to the standing assumptions about the relationship between visual WM and overt attention. Future work should explore the conditions under which eye movements engage visual WM and the conditions under which they do not.

Acknowledgments

This research was supported by a grant from the National Eye Institute of the National Institutes of Health under award number R01EY027792. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

C.E.P. designed the study. C.E.P. & D.A.C. conducted all analyses. D.A.C. wrote the manuscript. C.E.P. & D.A.C. designed the figures. J.M.H. & C.E.P. edited the manuscript. All authors approved the final version. We thank Conner McDaniels, Praveena Singh, and Jocelyn Huerta for assisting in data collection.

References

- Allen RJ, Baddeley AD, & Hitch GJ (2006). Is the binding of visual features in working memory resource-demanding? Journal of Experimental Psychology: General, 135, 298–313. 10.1037/0096-3445.135.2.298 [PubMed: 16719655]
- Anliker L (1976). Eye movements: On-line measurement, analysis, and control In Monty RA & Senders JW (Eds.), Eye movements and psychological processes (pp. 185–199). Hillsdale, NJ: Erlbaum.
- Baddeley AD (1986). Working Memory. Oxford, UK: Clarendon.
- Baddeley AD & Hitch GJ (1974). Working memory In Bower GH (Ed.), The Psychology of Learning and Motivation (Vol. VIII, pp. 47–90). New York, NY: Academic Press.
- Bae G-Y & Luck SJ (2018). What happens to an individual visual working memory representation when it is interrupted? British Journal of Psychology, 110, 268–287. DOI:10.1111/bjop.12339 [PubMed: 30069870]
- Bahle B, Beck VM, & Hollingworth A (2018). The architecture of interaction between visual working memory and visual attention. Journal of Experimental Psychology: Human Perception and Performance, 44, 992–1011. [PubMed: 29629781]
- Bates D, Maechler M, Bolker B, & Walker S (2015). Fitting linear mixed-effects models using lme4. Journal of Statistical Software, 67(1), 1–48. DOI:10.18637/jss.v067.i01.
- Brysbaert M & Stevens M (2018). Power analysis and effect size in mixed effects models: a tutorial. Journal of Cognition, 1(1), 9 DOI: 10.5334/joc.10 [PubMed: 31517183]

- Castelhano MS, Mack M, & Henderson JM (2009). Viewing task influences eye movement control during active scene perception. Journal of Vision, 9(3):6, 1–15.
- Cronin DA, & Irwin DE (2018). Visual working memory supports perceptual stability across saccadic eye movements. Journal of Experimental Psychology: Human Perception and Performance, 44(11), 1739–1759. 10.1037/xhp0000567 [PubMed: 30091634]
- de Fockert JW (2013). Beyond perceptual laod and dilution: a review of the role of working memory in selective attention. Frontiers in Psychology, 4(287), 1–12. [PubMed: 23382719]
- Fougnie D & Marois R (2007). Executive working memory load induces inattentional blindness. Psychonomic Bulletin, & Review, 14(1), 142–147. [PubMed: 17546744]
- Hardman KO, Vergauwe E, & Ricker TJ (2017). Categorical working memory representations are used in delayed estimation of continuous colors. Journal of Experimental Psychology: Human Perception and Performance, 43, 30–54. 10.1037/xhp0000290 [PubMed: 27797548]
- Hollingworth A & Luck SJ (2009). The role of visual working memory in the control of gaze during visual search. Attention, Perception, & Psychophysics, 71(4), 936–949. DOI: 10.3758/ APP.71.4.936
- Hollingworth A, Richard AM, & Luck SJ (2008). Understanding the function of visual short-term memory: Transsaccadic memory, object correspondence, and gaze correction. Journal of Experimental Psychology: General, 137, 163–181. [PubMed: 18248135]
- Irwin DE (1992a). Memory for position and identity across eye movements. Journal of Experimental Psychology: Learning, Memory, & Cognition,18, 307–317.
- Irwin DE (1992b). Perceiving an integrated visual world In Meyer DE & Kornblum S (Eds.), Attention and performance XIV: Synergies in experimental psychology, artificial intelligence, and cognitive neuroscience (pp. 121–142). Cambridge, MA: MIT Press.
- Kiyonaga A & Egner T (2013). Working memory as internal attention: Toward an integrative account of internal and external processes. Psychonomic Bulletin, & Review, 20(2), 228–242. 10.3758/ s13423-012-0359-y [PubMed: 23233157]
- Lavie N, & de Fockert JW (2005). The role of working memory in attentional capture. *Psychonomic Bulletin &* Review, 12, 669–674. 10.3758/BF03196756
- Lavie N, Hirst A, de Fockert JW, & Viding E (2004). Load theory of selective attention and cognitive control. Journal of Experimental Psychology: General, 133(3), 339–354. DOI: 10.1037/0096-3445.133.3.339 [PubMed: 15355143]
- Luck SJ, & Vogel EK (1997). The capacity of visual working memory for features and conjunctions. Nature, 390, 279–281. 10.1038/36846 [PubMed: 9384378]
- Mack A & Rock I (1998). Innattentional Blindness. Cambridge, MA, US: The MIT Press.
- Menneer T, Cave KR, Kaplan E, Stroud MJ, Chang J, & Donnelly N (2019). The relationship between working memory and the dual-target cost in visual search guidance. Journal of Experimental Psychology: Human Perception and Performance, 45(7), 911–935. 10.1037/xhp0000643 [PubMed: 30985178]
- Miller GA (1956). The magical number seven, plus or minus two: Some limits on our capacity for processing information. Psychological Review, 63, 81–97. [PubMed: 13310704]
- Most SB, Simons DJ, Scholl BJ, Jimenez R, Clifford E, et al. (2001). How not to be seen: The contribution of similarity and selective ignoring to sustained inattentional blindness. Psychological Science, 12(1), 9–17. [PubMed: 11294235]
- Makovski T, Shim WM, & Jiang YV (2006). Interference from filled delays on visual change detection. Journal of Vision, 6, 1459–1470. 10.1167/6.12.11 [PubMed: 17209748]
- Morey CC, & Bieler M (2013). Visual short-term memory always requires general attention. Psychonomic Bulletin & Review, 20, 163–170. 10.3758/s13423-012-0313-z
- Morey CC & Cowan N (2004). When visual and verbal memories compete: Evidence of cross-domain limits in working memory. Psychonomic Bulletin & Review, 11(2), 296–301. 10.3758/ BF03196573 [PubMed: 15260196]

- Neisser U (1979). The control of information pickup in selective looking, in Perception and its Development: A Tribute to Eleanor J Gibson. In: Pick AD, editor. Hillsdale, NJ: Lawrence Erlbaum Associates pp. 201–219.
- Neisser U & Becklen R (1975). Selective looking: Attending to visually specified events. Cognitive Psychology, 7, 480–494. DOI: 10.1016/0010-0285(75)90019-5.
- Peterson DJ, Decker R, Naveh-Benjamin M (2019). Journal of Experimental Psychology: Learning, Memory, & Cognition, 45(1), 56–70. DOI: 10.1037/xlm0000577
- R Core Team (2019). R: A language and environment for statistical computing R Foundation for Statistical Computing, Vienna, Austria URL https://www.R-project.org/.
- Ricker TJ, Cowan N, & Morey CC (2010). Visual working memory is disrupted by covert verbal retrieval. Psychonomic Bulletin & Review, 17, 516–521. 10.3758/PBR.17.4.516 [PubMed: 20702871]
- Saults JS, & Cowan N (2007). A central capacity limit to the simultaneous storage of visual and auditory arrays in working memory. Journal of Experimental Psychology: General, 136, 663–684. 10.1037/0096-3445.136.4.663 [PubMed: 17999578]
- Schmidt BK, Vogel EK, Woodman GF, & Luck SJ (2002). Voluntary and automatic attentional control of visual working memory. Perception & Psychophysics, 64(5), 754–763. [PubMed: 12201334]
- Simons DJ & Chabris CF (1999). Gorillas in our midst: sustained inattentional blindness for dynamic events. Perception, 28(9), 1059–1074. [PubMed: 10694957]
- Smith EE, Jonides J, & Koeppe RA (1996). Dissociating verbal and spatial working memory using PET. Cerebral Cortex, 6(1), 11–20. [PubMed: 8670634]
- Soto D, & Humphreys GW (2007). Automatic guidance of visual attention from verbal working memory. Journal of Experimental Psychology: Human Perception and Performance, 33(3), 730– 757. DOI: 10.1037/0096-1523.33.3730 [PubMed: 17563233]
- Soto D, & Humphreys GW (2008). Stressing the mind: The effect of cognitive load and articulatory suppression on attentional guidance from working memory. Perception & Psychophysics, 70(5), 924–934. [PubMed: 18613638]
- Tas AC, Luck SJ, & Hollingworth A (2016). The relationship between visual attention and visual working memory encoding: A dissociation between covert and overt orienting. Journal of Experimental Psychology: Human Perception and Performance, 42, 1121–1138. [PubMed: 26854532]
- Tsubomi H, Fukuda K, Watanabe K & Vogel EK (2013). Neural limits to representing objects still within view. Journal of Neuroscience, 33(19): 8257–8263. DOI: 10.1523/ JNEUROSCI.5348-12.2013 [PubMed: 23658165]
- Vergauwe E, Barroillet P, & Camos V (2010). Do mental processes share a domain-general resource? Psychological Science, 21(3), 384–390. DOI: 10.1177/0956797610361340 [PubMed: 20424075]
- Westfall J Kenny DA, & Judd CM (2014). Statistical power and optimal design in experiments in which samples of participants respond to samples of stimuli. Journal of Experimental Psychology: General, 143(5): 2020–2045. DOI: 10.1037/xge0000014 [PubMed: 25111580]

Open Practices Statement

Data and materials for this experiment are available upon request. This study was not preregistered.

Visual Load



No Load (Visual)



Verbal Load

Fixation	Encode 7 Letters	Mask	Fixation	Free Viewing	Recognition
+	F N		+		Z
500ms	300ms/letter	250ms	250ms	3500ms	Self-timed
		x24 t	rials		\rightarrow

No Load (Verbal)



Figure 1.

Task procedures for the four conditions.

Cronin et al.



Figure 2.

Distribution of all participants' fixations through all scenes in the four scene-viewing conditions. Blue points represent individual fixations.

Table 1

Mean (SD) eye movement measures across task conditions.

	Visual		Verbal	
	No Load	Load	No Load	Load
Number of Fixations	9.65 (1.95)	7.29 (2.05)	9.46 (2.17)	8.36 (1.84)
Fixation Duration (ms)	327 (121)	422 (170)	360 (172)	375 (140)
Saccade Amplitude (°)	5.01 (1.10)	3.19 (1.42)	4.64 (1.20)	3.70 (1.30)
Scan path length (°)	44.32 (13.10)	21.95 (14.32)	41.71 (15.41)	28.82 (12.51)
Percent of scene fixated	4.43% (0.98)	2.82% (1.04)	4.26% (1.12)	3.34% (0.97)
Dispersion from center (pixels)	275 (37)	173 (47)	299 (36)	211 (40)

Table 2

Summary of Linear Mixed Effects Models for Measures of Fixation Spread

Measure	$\chi^{2}(1)$	р	d	
Scan Path Length				
Load	1250.05	< 0.001	0.66	
Load-Type	19.40	< 0.001	0.33	
Interaction	74.55	< 0.001	0.43	
Percent Scene Fixated				
Load	1118.81	< 0.001	0.63	
Load-Type	22.49	< 0.001	0.33	
Interaction	71.83	< 0.001	0.43	
Dispersion from Center				
Load	5417.33	< 0.001	1.68	
Load-Type	584.19	< 0.001	0.71	
Interaction	22.68	< 0.001	0.23	