Lawrence Berkeley National Laboratory

Recent Work

Title

PSEUDOSCALAR AND VECTOR EXCHANGES IN THE PRODUCTION OF VECTOR MESONS

Permalink

<https://escholarship.org/uc/item/9tm133fd>

Author

Huff, Robert W.

Publication Date

1963-09-05

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

PSEUDOSCALAR AND VECTOR EXCHANGES IN THE
PRODUCTION OF VECTOR MESONS

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

PSEUDOSCALAR AND VECTOR EXCHANGES IN THE
PRODUCTION OF VECTOR MESONS

Robert W. Huff

September 5, 1963

Pseudoscalar and Vector Exchanges in the
Production of Vector Mesons

Robert W. Huff

University of California
Lawrence Radiation Laboratory
Berkeley, California

September 5, 1963

ABSTRACT

The cross section for the general reaction (pseudoscalar meson + nucleon $\rightarrow$ spin- $\frac{1}{2}$ baryon + vector meson) and the decay angular distributions for the final baryon and vector meson are calculated under the assumption that the reaction is dominated by the exchange of pseudoscalar and vector mesons. The results are applied to an analysis of the reaction $\pi^- p \rightarrow \Sigma^0 K^{*0}$.

Pseudoscalar and Vector Exchanges in the Production of Vector Mesons*

Robert W. Huff

University of California
Lawrence Radiation Laboratory
Berkeley, California

September 5, 1963

I. INTRODUCTION

In this paper the cross section for the general reaction (pseudoscalar meson + nucleon $\rightarrow$ spin- $\frac{1}{2}$ baryon + vector meson) is calculated, by assuming that the reaction is dominated by the exchange of pseudoscalar and vector mesons. In Section II, we derive expressions for this cross section, and for the decay angular distributions for the final baryon and vector meson. Section III contains a discussion of the structure of the form factors that appear in these expressions. In Section IV we use the results of the preceding sections in an analysis of the reaction $\pi^- p \rightarrow \Sigma^0 K^{*0}$, which analysis is an extension of one reported earlier.¹

II. CALCULATION OF CROSS SECTIONS

We will use the conventions $\hbar = c = 1$, $g^{\mu\nu} = (1, -1, -1, -1)$, $A_\mu B^\mu = A^0 B^0 - \underline{A} \cdot \underline{B}$, $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$, $\sigma^{\mu\nu} = \frac{1}{2} i[\gamma^\mu, \gamma^\nu]$, and $\gamma_5 = \gamma^0 \gamma^1 \gamma^2 \gamma^3$. Also, $\epsilon_{\mu\nu\lambda\sigma}$ is a completely antisymmetric tensor, which is +1 when $(\mu\nu\lambda\sigma)$ is an even permutation of (0123), -1 when it is odd, and zero otherwise. All spinors will be normalized so that $\sum_r u_r(p) \bar{u}_r(p) = \not{p} + m$.

Let us begin by considering the reaction $K^- p \rightarrow \Lambda \omega \rightarrow (\pi^- p) (\pi^+ \pi^- \pi^0)$. Other reactions of the general form (pseudoscalar meson + nucleon $\rightarrow$ spin- $\frac{1}{2}$ baryon + vector meson) will have the same results, except for a possible overall numerical factor for isotopic spin and a possible

modification for different decay interactions for the final particles. Let p, r, H, Q be the momenta of the target nucleon, incident pseudoscalar meson, final baryon, and vector meson, respectively. Define two additional momenta, $k = H - p = r - Q$ and $s = p + r = H + Q$, so that k^2 and s^2 are the squares of the invariant momentum transfer and of the total center-of-mass energy, respectively. Let $m, \bar{m}$, be the masses of the target nucleon and incident pseudoscalar, $M, \bar{M}$ the masses of the final baryon and vector particle, and v_p and v_v the masses of the exchanged pseudoscalar (K) and vector (K^*) mesons, respectively. Then the most general Feynman amplitudes that can be written for the relevant three-particle vertices are

$$\mathcal{M}(pK\Lambda) = (4\pi)^{1/2} \bar{u}(H) [f_0 \gamma_5] u(p), \quad (1a)$$

$$\mathcal{M}(pK^*\Lambda) = (4\pi)^{1/2} \bar{u}(H) [f_1 \gamma_0 + i f_2 \sigma_{0\lambda} k^\lambda / (M+m)] u(p) e^\delta(k), \quad (1b)$$

$$= (4\pi)^{1/2} \bar{u}(H) [(f_1 + f_2) \gamma_0 - f_2 (p_0 + H_0) / (M+m)] u(p) e^\delta(k), \quad (1c)$$

$$\mathcal{M}(KK\omega) = (4\pi)^{1/2} f_3 r_\mu e^\mu(Q), \quad (1d)$$

and

$$\mathcal{M}(KK^*\omega) = (4\pi)^{1/2} f_4 \epsilon_{\mu\nu\sigma\rho} e^\mu(Q) e^\nu(k) r^\sigma Q^\rho / \bar{M}, \quad (1e)$$

where $e(k)$ and $e(Q)$ are the unit polarization vectors of the exchanged and final vector particles, respectively. The five form factors, f_i , are dimensionless functions of the single variable, k^2 , which may be interpreted as the square of the "effective mass" of the exchanged particle.

The K^- and K^{*-} -exchange contributions then give a $K^+p \rightarrow \Lambda\omega$ Feynman amplitude

$$\mathcal{M}(pK\Lambda\omega) = 4\pi\bar{u}(H) [N_\mu e^\mu(Q)] u(p), \quad (2)$$

where

$$N_\mu = F_0 \gamma_5 r_\mu + [F_1 \gamma_0 - F_2 (p_0 + H_0)/2M] (-g^{\delta\gamma} + k^\delta k^\gamma / v^2) \epsilon_{\mu\gamma\sigma\rho} r^\sigma Q^\rho / \bar{M} \quad (3a)$$

$$= F_0 \gamma_5 r_\mu + [F_1 \gamma_0^\delta - F_2 H_0^\delta / M] \epsilon_{\delta\sigma\rho\mu} r^\sigma Q^\rho / \bar{M} \quad (3b)$$

with

$$F_0 = f_0 f_3 / (k^2 - v_p^2), \quad (4a)$$

$$F_1 = (f_1 + f_2) f_4 / (k^2 - v_v^2), \quad (4b)$$

$$\text{and} \quad F_2 = [2M/(M+m)] f_2 f_4 / (k^2 - v_v^2). \quad (4c)$$

If the ω and Λ polarizations are unobserved and the proton is unpolarized, then the $K^- p \rightarrow \Lambda\omega \rightarrow (\pi^- p)(\pi^+ \pi^- \pi^0)$ differential cross section in the c.m. system is

$$\sigma_H = \frac{\partial\sigma}{\partial\Omega_{H,s}} = \frac{\eta |H_s|}{64\pi^2 s^2 |\underline{r}_s|} \cdot \frac{1}{2} \sum_{p,\Lambda,\omega} |\mathcal{M}(pK\Lambda\omega)|^2 \quad (5)$$

$$= \frac{\eta |H_s|}{2s^2 |\underline{r}_s|} \cdot \frac{1}{4} \text{Tr} [(H_a \gamma^a + M) N_\mu (p_\beta \gamma^\beta + m) \bar{N}_\nu] (-g^{\mu\nu} + Q^\mu Q^\nu / \bar{M}^2). \quad (6)$$

We obtain $\bar{N}$ from N by replacing the F_i by their complex conjugates $\bar{F}_i$. The symbols $A_B^\mu = (A_B^0, \underline{A}_B)$ and $\Omega_{A,B}$ refer to the vector A and the direction of its spatial part, both evaluated in the "rest frame" of the vector B , i.e., the frame in which the spatial part of B vanishes. Thus the subscripts s , Q , and H refer to the c.m. system, the ω rest frame, and the Λ rest frame, respectively. The subscripts to the summation indicate which particles are included in the polarization sum. Note that we have included a factor $\eta \equiv \eta_Q \eta_H$ in the cross section, where $\eta_Q (\eta_H)$ is the

fraction of the final vector particles (baryons) that decay via the observed decay mode.

Rather than evaluate σ_{II} now, we first investigate the decay distributions predicted for the final particles. For the decay $\omega \rightarrow \pi^+ \pi^0 \pi^-$, the most general I-spin-conserving Feynman matrix element is

$$\mathcal{M}(\omega \pi \pi \pi) = f_5 e^\nu(Q) q_+^\alpha q^\beta q_-^\gamma \epsilon_{\nu\alpha\beta\gamma} \quad (7)$$

where q_+ , q , and q_- are the π momenta, and f_5 is a function of two variables [e.g., q^0 and $(\Delta \equiv q_+^0 - q_-^0)$] which is completely symmetric in the three π 's. The $\omega \rightarrow \pi^+ \pi^0 \pi^-$ decay rate for a polarized ω is, in the ω rest frame,

$$\Gamma_\omega = \int \frac{d^3 q_- d^3 q d^3 q_+ \delta^4(Q - q_+ - q - q_-)}{(2\pi)^5 16 \bar{M} q_+^0 q^0 q_-^0} |\mathcal{M}(\omega \pi \pi \pi)|^2 \quad (8)$$

$$= \int \frac{d\Omega \, n_Q \, d\phi \, dq^0 \, d\Delta}{32 (2\pi)^5 M} |\bar{M} f_5 \underline{q}_+ \times \underline{q}_-|^2 (\underline{e}_Q \cdot \underline{n}_Q)^2 \quad (9)$$

where $\underline{n}_Q$ is a unit vector in the direction $\underline{q}_+ \times \underline{q}_-$ (i.e., normal to the decay plane) and ϕ is the azimuthal angle of $\underline{q}$ about $\underline{n}_Q$. Note that throughout this section the vectors $\underline{q}_+$, $\underline{q}$, and $\underline{q}_-$ are to be evaluated in the ω rest frame although the subscript Q has been suppressed.

If the target proton is unpolarized and the Λ polarization is unobserved, then the $K^- p \rightarrow \Lambda \omega \rightarrow (\pi^- p) (\pi^+ \pi^- \pi^0)$ differential cross section is

$$\sigma_{H,n} \equiv \frac{\partial \sigma_H}{\partial \Omega_{n,Q}} = A \frac{\eta |H_s|}{64 \pi^2 s^2 |r_s|} \int \frac{d\phi \, dq^0 \, d\Delta}{32 (2\pi)^5 \bar{M}} \cdot \frac{1}{2} \sum_{p,\Lambda} \left| \sum_{\omega} \mathcal{M}(pK\Lambda\omega) \mathcal{M}(\omega\pi\pi\pi) \right|^2 \quad (10)$$

$$= \frac{\eta |H_s|}{2 s^2 |r_s|} \cdot \frac{1}{4} \text{Tr}[(H_a \gamma^a + M) Q] \cdot A \int \frac{d\phi \, dq^0 \, d\Delta}{32 (2\pi)^5 \bar{M}} |\bar{M} f_5 \underline{q}_+ \times \underline{q}_-|^2 \quad (11)$$

where

$$Q = (N_\mu n^\mu) (p_\beta \gamma^\beta + m) (\bar{N}_\nu n^\nu). \quad (12)$$

Here A is a normalization factor to be chosen such that $\int d\Omega_{n,Q} \sigma_{H,n} = \sigma_H$, and n^μ is a covariant unit vector with the value $n_Q^\mu = (0, \underline{n}_Q)$ in the ω rest frame. From Eq. (11), we see that all azimuthal orientations of the ω decay plane about $\underline{n}_Q$, as measured by ϕ , are equally likely. Note also that the distribution in q^0 and Δ (i.e., the Dalitz-plot density) is independent of n^μ . Both these results are consequences of the form of the matrix element $\mathcal{M}(\omega\pi\pi\pi)$, however, and cannot be used for an experimental check of our K^- and K^* -exchange model. In treating the ω decay in this way, we have assumed only that the ω has a long enough lifetime--and thus decays sufficiently far from the Λ --so that its decay pions do not interact with the Λ . Use of the relation $\int d\Omega_{n,Q} n^\mu n^\nu = \frac{4\pi}{3} (-g^{\mu\nu} + Q^\mu Q^\nu / M^2)$, along with a comparison of Eqs. (6), (9), and (11), shows $A = 1/\Gamma_\omega$ and

$$\sigma_{H,n} = \frac{3\eta |\underline{H}_s|}{8\pi s^2 |\underline{r}_s|} \cdot \frac{1}{4} \text{Tr} [(H_a \gamma^a + M) Q]. \quad (13)$$

Finally, we consider the $\Lambda \rightarrow \pi^- p$ decay, with a Feynman matrix element

$$\mathcal{M}(\Lambda\pi p) = \bar{u}(h) [a + b \gamma_5] u(H). \quad (14)$$

where the constants a and b are the parity-nonconserving and parity-conserving amplitudes, respectively, and h is the decay proton momentum. In the Λ rest frame, a Λ of polarization $\underline{P}$ will correspond to $u(H) \bar{u}(H) = \frac{1}{2} M (1 + \gamma^0) (1 + \underline{P} \cdot \underline{\sigma}) = \frac{1}{2} M (1 + \gamma^0) (1 + i \underline{P} \cdot \underline{\gamma} \gamma_5)$. If the polarization of the decay proton is unobserved, then the $\Lambda \rightarrow \pi^- p$ decay rate in the Λ rest frame is

$$\Gamma_{\Lambda} = \int d\Omega_{\underline{h},H} \sum_P |\mathcal{M}(\Lambda\pi p)|^2 \cdot |\underline{h}_H|/32 \pi^2 M^2 \quad (15)$$

$$= \int d\Omega_{\underline{h},H} (1 + a \hat{\underline{h}}_H \cdot \underline{P}) C |\underline{h}_H|/16 \pi^2 M, \quad (16)$$

where $C = [|a|^2 (h_H^0 + M) + |b|^2 (h_H^0 - M)] , \quad (17)$

$$a = 2 |\underline{h}_H| \operatorname{Im}(ab^*)/C, \quad (18)$$

and $\hat{\underline{h}}_H$ is a unit vector in the direction of $\underline{h}_H$.

With unpolarized target protons and unobserved decay-proton polarization, the $K^-p \rightarrow \Lambda\omega \rightarrow (\pi^-p)(\pi^+\pi^-\pi^0)$ differential cross section is

$$\sigma_{H,n,h} = \frac{\partial \sigma_{H,n}}{\partial \Omega_{\underline{h},H}} = B \frac{3\eta |\underline{H}_s|}{8\pi s^2 |\underline{r}_s|} \cdot \frac{|\underline{h}_H|}{32\pi^2 M^2} \cdot \frac{1}{4} \operatorname{Tr} [\alpha \mathcal{B}], \quad (19)$$

where $\mathcal{B} = (H_a \gamma^a + M) (\bar{a} + \bar{b} \gamma_5) (h_b \gamma^b + m) (a + b \gamma_5) (H_0 \gamma^0 + M) . \quad (20)$

As before, B is a normalization factor to be chosen such that

$\int d\Omega_{\underline{h},H} \sigma_{H,n,h} = \sigma_{H,n}$. A simple calculation gives

$$\mathcal{B} = 2MC (1 + i a \gamma_5 \hat{h}^\mu \gamma_\mu) (H_a \gamma^a + M) = 2MC \alpha \mathcal{O}, \quad (21)$$

where $\hat{h}^\mu$ is a covariant unit vector with the value $\hat{h}_H^\mu = (0, \hat{\underline{h}}_H)$ in the Λ rest frame. A comparison of Eqs. (13), (16), and (19) then shows $B = 1/\Gamma_{\Lambda}$ and

$$\sigma_{H,n,h} = \frac{3\eta |\underline{H}_s|}{32\pi^2 s^2 |\underline{r}_s|} \cdot \frac{1}{4} \operatorname{Tr} [\alpha \mathcal{O}]. \quad (22)$$

The trace calculation is straightforward, and yields

$$\frac{1}{4} \text{Tr} [\mathcal{A}] = D_0 + a \hat{h}_H \cdot (\underline{D}_1 + \underline{D}_2 + \underline{D}_3), \quad (23)$$

where

$$D_0 = \frac{1}{2} [(M - m)^2 - k^2] [|F_0|^2 (\underline{r} \cdot \underline{n})^2 + |F_1|^2 (\underline{r} \times \underline{n})^2] \\ + 2 [|F_1 - \frac{1}{2} (1 + m/M) F_2|^2 - k^2 |F_2|^2 / 4 M^2] (\underline{H} \times \underline{r} \cdot \underline{n})^2, \quad (24)$$

$$\underline{D}_1 = \text{Im} [\overline{F}_0 F_1] [(M - m)^2 - k^2] (\underline{r} \cdot \underline{n}) \underline{R}_H, \quad (25a)$$

$$\underline{D}_2 = 2 \text{Im} [\overline{F}_0 (F_1 - F_2)] (\underline{r} \cdot \underline{n}) (\underline{H} \times \underline{r} \cdot \underline{n}) \underline{P}_H, \quad (25b)$$

and

$$\underline{D}_3 = 2 \text{Im} [F_1 \overline{F}_2] (\underline{H} \times \underline{r} \cdot \underline{n}) (\underline{R}_H \times \underline{P}_H). \quad (25c)$$

Note that $\underline{H}$, $\underline{r}$, and $\underline{n}$ are to be evaluated in the ω rest frame, although the subscript Q has been suppressed for conciseness. $\underline{R}$ is a covariant vector with the value $\underline{R}_Q = (0, \underline{r} \times \underline{n})$ in the ω rest frame. We obtain σ_H and $\sigma_{H,n}$ by simply integrating Eq. (22) with respect to the solid angles $\Omega_{h,H}$ and $\Omega_{n,Q}$.

The "rest frames" of s , Q , and H are related by Lorentz transformations without spatial rotations. In order to avoid any possible ambiguities, those spatial vectors whose values in the Λ rest frame are required in Eqs. (23) and (25) are given below, expressed in terms of the vectors $\underline{H}$, $\underline{r}$, $\underline{n}$, and $\underline{h}_Q$:

$$\underline{P}_H = -\underline{r} - \underline{H} [\overline{M} - r_Q^0 + (\underline{H} \cdot \underline{r}) / (H_Q^0 + M)] / M, \quad (26a)$$

$$\underline{R}_H = (\underline{r} \times \underline{n}) + \underline{H} (\underline{H} \times \underline{r} \cdot \underline{n}) / M (H_Q^0 + M), \quad (26b)$$

and

$$|\underline{h}_H| \hat{h}_H = \underline{h}_Q - \underline{H} [h_H^0 + (\underline{H} \cdot \underline{h}_Q) / (H_Q^0 + M)] / H_Q^0. \quad (26c)$$

Note that h_H^0 and $|h_H|$ are constants, independent of H and h_Q .

For other reactions we may have to modify the above expressions because of isotopic-spin considerations. The isotopic-spin dependence of the matrix elements in Eqs. (1) has not been exhibited because it results in a factor of unity. For the reaction $\pi^- p \rightarrow \Lambda K^{*0}$, however, we would include factors of $\bar{V}_K V_p$, $\bar{V}_{K^*} V_p$, $\bar{V}_{K^*} \tau_\pi V_K$, and $\bar{V}_{K^*} \tau_\pi V_{K^*}$ in Eqs. (1a), (1b), (1c), (1d), and (1e), respectively. Here V and $\bar{V}$ are two-component isotopic spinors, and τ is the isotopic-spin analog of the Pauli spin matrices. Thus the above cross sections would be multiplied by an overall factor $|\bar{V}_{K^*} \tau_\pi V_p|^2 = |(0, 1) \begin{pmatrix} 0 & 0 \\ \sqrt{2} & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}|^2 = 2$. Similarly, there would be a factor 2 for the reaction $\pi^- p \rightarrow \Sigma^0 K^{*0}$, and a factor 4 for a reaction such as $\pi^+ p \rightarrow \Sigma^+ K^{*+}$. For the reaction $K^+ p \rightarrow \bar{K}^0 n$, which involves $\pi^\pm$ and $\rho^\pm$ exchange, there is a factor 4. The factor is unity for the non-charge-exchange reactions, $K^\pm p \rightarrow K^\pm p$, but since η and ω exchange are coherent with π^0 and ρ^0 exchange in these reactions, we must make the substitutions $F_0 \rightarrow F_{0\eta} \pm F_{0\pi}$, $F_1 \rightarrow F_{1\omega} \pm F_{1\rho}$, and $F_2 \rightarrow F_{2\omega} \pm F_{2\rho}$, where the sign $\pm$ is the same as the K charge, and where $F_{0\eta}$, $F_{0\pi}$, $F_{1\omega}$, $F_{1\rho}$, $F_{2\omega}$, and $F_{2\rho}$ each have the form given by Eq. (4). Note that reactions with an incident π will generally not involve both ρ and ω exchange, however, nor both η and π exchange, because this would not conserve G parity.

Finally we consider the modifications to the above expressions that are necessary when particles other than the ω and Λ are produced. The matrix element for decay of the vector particle can always be written in the form $e^\mu(Q) V_\mu$. The amplitude for production and decay would then have the form $\bar{u}(H) [N_\mu v^\mu] u(p)$, where $v^\mu = (-g^{\mu\nu} + Q^\mu Q^\nu / M^2) V_\nu$. Thus $v_Q^\mu = (0, -\underline{V}_Q)$, and we obtain the correctly normalized cross sections simply by redefining $\underline{n}$ as the unit vector in the direction $-\underline{V}_Q$. For $\rho \rightarrow \pi\pi$ or $K^* \rightarrow K\pi$, V^μ is

just the momentum of one of the decay particles, so that $\underline{n}$ is thus a unit vector in the decay direction in the rest frame of the decaying particle.

The results for the Λ decay distribution apply directly to the $\Sigma^\pm$ decay, but with different experimental values for α . The situation is more complicated for spin- $\frac{1}{2}$ baryons that undergo a sequence of decays. The weak decay at the end of the sequence is the only decay that can produce anisotropy in the decay distributions, and is hence the only decay that can give any information on the polarization of the baryon. We will not consider such cases further, except for one comment concerning the $\Sigma^0 \rightarrow \Lambda + \gamma$ decay. It is predicted theoretically² that after integration over the photon direction and polarization, the average Λ polarization is one-third that of the Σ^0 , but in the opposite direction. Therefore our previous results also apply to Σ^0 decay if we use a value $\alpha(\Sigma^0) = -\frac{1}{3} \alpha(\Lambda)$, and still let $\hat{h}_H$ be the direction of the decay proton in the Λ (not Σ) rest frame. Note that with our convention (i.e., protons emitted preferentially in the direction of the baryon spin for $\alpha > 0$), experimental values for α are³ $\alpha(\Lambda) = 0.62 \pm 0.07$, and therefore $\alpha(\Sigma^0) = -0.21 \pm 0.02$.

III. DISCUSSION OF THE MODEL

Again let us consider the reaction $K^- p \rightarrow \Lambda \omega$ as an example of our general reaction. The Feynman diagrams for this reaction may be separated into four sets of diagrams. The first set, represented by Fig. 1a, contains all those diagrams that can be divided into two parts (one connected to the external p and Λ lines, and the other to the external K^- and ω lines) by the cutting of a single internal line. In other words, these are merely the diagrams that contain a single virtual particle at some point in the k channel. Similarly, the second and third sets of diagrams,

represented in Fig. 1b and 1c, are those diagrams containing a single particle at some point in the s and t channels, respectively, where $t = H - r = p - Q$. The fourth set, represented by Fig. 1d, contains all diagrams that cannot be so divided by cutting a single line, i.e., that contain two or more particles in each of the three channels.

Our model includes only those diagrams of the first set that represent the exchange of a vector or pseudoscalar particle. Scalar exchange cannot contribute because a scalar and pseudoscalar cannot be coupled to form a vector. We have not included the exchange of pseudovectors or of particles of spin greater than one, because we have not yet seen such particles experimentally. Diagrams of the other three sets are not included in the model. It is for reasons of mathematical convenience and simplicity, however, that these other diagrams have been ignored, rather than because of any good experimental evidence.

A more detailed look at the structure of the vertices in Fig. 1a will yield information on the form of the functions f_i . As seen in Eqs. (4), the functions F_i have poles at $k^2 = v^2$, where v is the physical mass of the exchanged particle. Similarly, the form factors f_i have singularities corresponding to the physical masses of the various intermediate states in the vertices. Those vertex diagrams of the form of Fig. 2a, where there is a single intermediate particle, would be expected to contribute simple poles. Diagrams of the form of Fig. 2b, with two or more intermediate particles, will contribute branch cuts. If we use the Cauchy theorem and assume the proper convergence at infinity, then the branch-cut contribution to f_i may be written in the form $\int dz h_i(z)/(k^2 - z)$, which is merely an integration over a continuous distribution of poles. This is only to be expected, because a system of two or more particles has a mass that ranges continuously from

some (positive) minimum up to infinity. Knowing the singularities of the form factor f_1 , we may easily write down the form of the functions F_1 . Note that the F_1 have no multiple poles. Multiple poles would correspond to a diagram such as Fig. 3, which merely gives a self-energy correction to the propagator, and does not (after renormalization,⁴ at least) yield a second-order pole. Thus, where $v_0 = v_p$ and $v_{1,2} = v_v$, we may write

$$F_1(k^2) = \frac{G_1}{k^2 - v_1^2} + \int dz \frac{g_1(z)}{k^2 - z}. \quad (27)$$

Note that from Eqs. (4) and (27) we have

$$G_0 = 2\pi i \text{Res} [F_0, k^2 = v_p^2] = f_0(v_p^2) f_3(v_p^2), \quad (28)$$

and similarly for the other G_i .

Since the form factors, f_i , evaluated on the mass shell, are just the corresponding renormalized coupling constants, we see that the G_i are merely products of the coupling constants at the two vertices. Reactions in which more than one vector particle can be exchanged (or more than one pseudoscalar) would have additional pole terms in Eq. (27), but these G_i would still be products of the coupling constants at the two vertices.

Let us briefly mention the qualitative effects to be expected from the other three sets of diagrams. Note that the functions F_1 , from the first set of diagrams, are functions only of k^2 and are independent of s^2 . Similarly, the other diagrams would introduce additional terms into the cross section, involving functions of s^2 only (from the second set of diagrams), of t^2 only (third set), and of s^2 and k^2 both (fourth set). (Note, incidentally, that there are only two independent variables because of the identity $s^2 + k^2 + t^2 = M^2 + \overline{M}^2 + m^2 + \overline{m}^2$.) The functions from the second and third

sets of diagrams would have, except for different variables, the same form as for Eq. (27). However, those from the fourth set have only the branch-cut term, because, by definition, these diagrams do not have any single-particle intermediate states. Now consider those diagrams of Fig. 1d in which the exchanged system of particles is in a $J^P = 0^-$ or $J^P = 1^-$ state. These diagrams will then give a scattering amplitude identical to that for the exchange of a single pseudoscalar or vector particle, except that the functions F_1 would have only the branch-cut term. The branch point will be the same as for single-particle exchange (because the possible intermediate states in Fig. 1d are the same as those in Fig. 2b), but the discontinuity across the branch cut will now be s^2 dependent. Therefore these diagrams can be included in our original calculations merely by replacing the functions $g_1(z)$ in Eq. (27) by $g_1'(z, s^2) = g_1(z) + g_1''(z, s^2)$, where $g_1''(z, s^2)$ is the contribution from the exchange of the pseudoscalar or vector systems of particles. Thus, the only effect of such diagrams is to introduce an s^2 dependence into the branch-cut terms in Eq. (27), and our results will now describe any pseudoscalar or vector exchanges, regardless of whether these are exchanges of single particles or of multiparticle systems.

IV. APPLICATION TO $\pi^-p \rightarrow \Sigma^0 K^0$

In this section, we present details of a previous analysis¹ of the reaction $\pi^-p \rightarrow \Sigma^0 K^0$, and also extend that analysis in order to obtain information about the functions F_1 . As reported earlier,¹ 209 events at incident-pion lab momenta of 2.17 and 2.25 BeV/c were attributed to $\pi^-p \rightarrow \Sigma^0 K^0 \rightarrow \Sigma^0 K^+ \pi^-$, and were found to have decay angular distributions

$$\rho(a) = 1 + (1.59 \pm 0.55) \cos^2 a, \quad (29a)$$

$$\rho(\beta) = 1 - (0.11 \pm 0.21) \cos^2 \beta, \quad (29b)$$

and
$$\rho(\gamma) = 1 - (0.74 \pm 0.11) \cos^2 \gamma. \quad (29c)$$

Here α , β , and γ are the angles, in the K^* rest frame, between the direction of the decay pion and the unit vectors $\hat{N}$, $\hat{r}$, and $\hat{r} \times \hat{N}$, respectively, where $\hat{r}$ is the incident-pion direction and $\hat{N}$ is the normal to the production plane. Because of the small number of events, the effects of a non- K^* background ($< 30\%$) were ignored, and no attempts were made to determine either the Σ^0 decay distribution or the dependence of the functions F_i upon k^2 and s^2 (i.e., upon the π^-p c.m. energy and production angle).

The appropriate cross section, $\sigma_n = \iint \sigma_{H,n,h} d\Omega_{h,H} d\Omega_{H,s}$, is then found from Eqs. (22) to (24) to be

$$\sigma_n = C_0 \cos^2 \beta + C_1 \sin^2 \beta + C_2 \cos^2 \alpha, \quad (30)$$

where
$$C_1 = \int d\mu J_1. \quad (31)$$

$$J_{0,1} = \frac{1}{2} [(M-m)^2 - k^2] |\underline{r}|^2 |F_{0,1}|^2 |\underline{H}_s|^2 / s^2 |\underline{r}_s|. \quad (32a)$$

and
$$J_2 = 2[|F_1 - \frac{1}{2}(1+m/M)F_2|^2 - k^2|F_2|^2/4M^2] |\underline{H}_s|^3 |\underline{r}_s|/\overline{M}^2. \quad (32b)$$

Here μ is the cosine of the angle between the final K^* and the incident π in the c.m. system, and we have used the relation

$$|\underline{H} \times \underline{r}|^2 = (1 - \mu^2) |\underline{H}_s|^2 |\underline{r}_s|^2 s^2 / \overline{M}^2.$$

An overall factor of 2 has been included for isotopic-spin considerations, as discussed in Sec. II. We have set $\eta = 2/3$, because this is the branching fraction predicted by isotopic-spin conservation, for the $K^* \rightarrow K^+ \pi^-$ mode

and because the corrections for the detection efficiency of the Σ^0 will be included in the experimental data.

In our analysis,¹ we fitted Eq. (30) to Eqs. (29b) and (29c) and treated the errors coherently by solving

$$\begin{cases} 2C_0 - 2C_1 - C_2 = -(0.11 + 0.21a)(2C_1 + C_2), \\ -C_0 + C_1 - C_2 = -(0.74 + 0.11b)(C_0 + C_1 + C_2), \end{cases}$$

where $a, b = 0 \pm 1$. (Note that $C_{0,1,2}$ corresponds to $g_{0,1,4}$ of reference 1, but that Eq. (3d) of that reference is written incorrectly.) The solution is then

$$\begin{aligned} \frac{C_2}{C_0} &= \frac{4 - 2(1.89 - 0.21a)(0.26 - 0.11b)}{(0.89 - 0.21a)(2.26 - 0.11b)} \\ &\approx 1.50 + 0.41a + 0.28b = 1.50 \pm 0.50 \end{aligned}$$

and

$$\begin{aligned} \frac{C_1}{C_0} &= \frac{(2.89 - 0.21a)(0.26 - 0.11b)}{(0.89 - 0.21a)(2.26 - 0.11b)} \\ &\approx 0.37 + 0.06a - 0.14b = 0.37 \pm 0.15. \end{aligned}$$

Because our total cross section is $\sigma = \int d\Omega_{n,Q} \sigma_n = \frac{4}{3} \pi (C_0 + 2C_1 + C_2)$, we may use the above solution to obtain

$$C_0 \approx (0.31 - 0.05a) 3\sigma/4\pi,$$

$$C_1 \approx (0.115 - 0.045b) 3\sigma/4\pi,$$

and

$$C_2 \approx (0.46 + 0.05a + 0.09b) 3\sigma/4\pi.$$

The measured total cross section⁵ for the reaction $\pi^+p \rightarrow \Sigma^0 K^+ \pi^-$ is $65 \pm 12 \mu\text{b}$ and $53 \pm 9 \mu\text{b}$ for incident-pion lab momenta of 2.17 and 2.25

BeV/c, respectively. This gives a mean value of $\sigma = 59 \pm 11 \mu\text{b}$ at 2.21 BeV/c, resulting in

$$C_0 = 4.4 \pm 1.1 \mu\text{b}, \quad (33a)$$

$$C_1 = 1.6 \pm 0.7 \mu\text{b}, \quad (33b)$$

and

$$C_2 = 6.5 \pm 1.8 \mu\text{b}. \quad (33c)$$

For convenience, we ignore the errors and use only the mean values for the C_i in the rest of this paper.

To determine the form factors, $f_i(k^2)$, we use Eqs. (4) and evaluate Eqs. (31) at $s = 2.25 \text{ BeV}$ (2.21 BeV/c incident momentum) to obtain

$$J_0 = (11.9 \mu\text{b}) \frac{(1 - 0.438 \mu)^2 (1 - 0.721 \mu)}{(1 - 0.619 \mu)^2} |f_0 f_3|^2. \quad (34a)$$

$$J_1 = (5.92 \mu\text{b}) \frac{(1 - 0.438 \mu)^2 (1 - 0.721 \mu)}{(1 - 0.435 \mu)^2} |(f_1 + f_2) f_4|^2, \quad (34b)$$

and

$$J_2 = \frac{1 - \mu^2}{(1 - 0.435 \mu)^2} [(22.4 \mu\text{b}) |f_1 f_4|^2 + (5.10 \mu\text{b}) (1 - 0.765 \mu) |f_2 f_4|^2], \quad (34c)$$

where $k^2 = (-1.032 + 0.790) \text{ BeV}^2$. By using Eqs. (31), (33), and (34), we find that average values for the form factors in the region of this experiment are $|f_0 f_3|^2 \approx 0.19$, $|(f_1 + f_2) f_4|^2 \approx 0.14$, and $|f_1 f_4|^2 + 0.20 |f_2 f_4|^2 \approx 0.20$, the latter two of which yield roughly $|f_1 f_4|^2 \approx 0.1$ to 0.2 and $|f_2 f_4|^2 \leq 0.2$.

These average values give us very little information about the values of the form factors on the mass shell, however, unless we can determine the k^2 dependence of the f_i from the production angular distribution. In this reaction, the form factors have no single-particle poles, but only the branch

cut from the multiparticle intermediate states, so that

$f_i(k^2) = \int dz h_i(z)/(k^2 - z)$. We will make the approximation $f_i(k^2) \propto 1/(k^2 - \Lambda_i^2)$ in the physical region for this experiment, which gives

$$F_0(k^2) \approx \frac{(G_0 + H_0)(\nu_p^2 - \Lambda_0^2)(\nu_p^2 - \Lambda_3^2)}{(k^2 - \nu_p^2)(k^2 - \Lambda_0^2)(k^2 - \Lambda_3^2)}, \quad (35)$$

and similarly for the other F_i . The parameters Λ_i may be interpreted as average masses of the intermediate states of the form factors. Note that $G_0 + H_0$ is the value of $f_0 f_3$ on the mass shell ($k^2 = \nu_p^2$) in this approximation, which is to be distinguished from the true value G_0 in Eqs. (27) and (28). By rewriting Eq. (35) as a sum of poles, we see that this approximation is equivalent to approximating the branch-cut integral of Eq. (28) by a sum of three poles with mutually related residues,

$$\int dz \frac{g_0(z)}{k^2 - z} \approx \frac{H_0}{k^2 - \nu_p^2} + \frac{(G_0 + H_0)(\Lambda_3^2 - \nu_p^2)}{(\Lambda_0^2 - \Lambda_3^2)(k^2 - \Lambda_0^2)} + \frac{(G_0 + H_0)(\Lambda_0^2 - \nu_p^2)}{(\Lambda_3^2 - \Lambda_0^2)(k^2 - \Lambda_3^2)}. \quad (36)$$

The branch cut in F_0 begins at $(m_K + 2m_\pi)^2 = 0.59 \text{ BeV}^2$ (because $K + \pi$ cannot be in a O^- state), which is well above the pole at $\nu_p^2 = 0.24 \text{ BeV}^2$. Thus in F_0 the $H_0/(k^2 - \nu_p^2)$ term has no direct physical meaning, and is introduced merely to improve the approximation in the physical region. It is expected that $H_0 \ll G_0$, which means that $G_0 + H_0$ is a good approximation to G_0 . For $F_{1,2}$, however, the branch cut begins at $(m_K + m_\pi)^2 = 0.40 \text{ BeV}^2$, which is below the pole at $\nu_v^2 = 0.78 \text{ BeV}^2$. The corresponding $H_{1,2}/(k^2 - \nu_v^2)$ term thus approximates the lower part of the branch cut, so that $H_{1,2}$ is a measure of the coupling to the low-mass multiparticle

intermediate states in $F_{1,2}$, just as $G_{1,2}$ gives the coupling to the K^* . In view of the small number of events in this experiment, let us make the further simplification of setting $\Lambda_0 = \Lambda_3 = \lambda$ and $\Lambda_1 = \Lambda_2 = \Lambda_4 = \Lambda$. Thus our final approximation is

$$f_0 f_3 / \hat{g}_0 = (\nu_p^2 - \lambda^2) / (k^2 - \lambda^2)^2 \quad (37a)$$

$$\text{and} \quad f_1 f_4 / \hat{g}_1 = f_2 f_4 / \hat{g}_2 = (\nu_v^2 - \Lambda^2) / (k^2 - \Lambda^2)^2, \quad (37b)$$

where $\hat{g}_0 = G_0 + H_0$, and similarly for $\hat{g}_{1,2}$.

When we combine Eqs. (31), (34), and (37) and compare the results with the experimental values in Eqs. (33), we obtain equations of the form

$$|\hat{g}_0|^2 = A(\lambda), \quad (38a)$$

$$|\hat{g}_1 + \hat{g}_2|^2 = B(\Lambda), \quad (38b)$$

$$\text{and} \quad C(\Lambda) |\hat{g}_1|^2 + D(\Lambda) |\hat{g}_2|^2 = 1. \quad (38c)$$

Since the relative phase of $\hat{g}_1$ and $\hat{g}_2$ is unknown, Eqs. (38) do not determine the magnitudes of $\hat{g}_1$ and $\hat{g}_2$, but only yield the restriction

$$\left| \frac{C B^{1/2} - (C + D - BCD)^{1/2}}{C + D} \right| \leq |\hat{g}_2| \leq \left| \frac{C B^{1/2} + (C + D - BCD)^{1/2}}{C + D} \right| \quad (39)$$

with a similar restriction on $|\hat{g}_1|$ due to Eq. (38c). The production angular distribution, $\rho(\mu) = (J_0 + 2J_1 + J_2) / (C_0 + 2C_1 + C_2)$, therefore involves the three parameters λ , Λ , and $|\hat{g}_2|^2$. A maximum-likelihood fit to the experimental¹ $\rho(\mu)$ yields $\lambda = \infty$, $\Lambda = 1.19_{-0.33}^{+0.27}$ BeV, and $|\hat{g}_2|^2 = 0.1_{-0}^{+85}$ with a χ^2 of 6.2 for six degrees of freedom. The upper and lower values for Λ are those for which the likelihood function, L , falls to $1/e$ of its

maximum value, $L_{\max}$. All values of $|\hat{g}_2|^2$ in the region $0.1 < |\hat{g}_2|^2 < 85$, as allowed by Eq. (39) for $\Lambda = 1.19$ BeV, correspond to values $L > L_{\max}/e$. The values for the other g_i are $|\hat{g}_0|^2 = 0.185$, $|\hat{g}_1 + \hat{g}_2|^2 = 30$, and $27 \geq |\hat{g}_1|^2 \geq 14$ for $\Lambda = 1.19$ BeV and $0.1 \leq |\hat{g}_2|^2 \leq 85$. The range of values for λ with this fit may be seen by noting that the values $\lambda = 2.06$ (0.95) BeV, $\Lambda = 1.46$ (3.3) BeV, $|\hat{g}_0|^2 = 0.56$ (9.4), $|\hat{g}_1 + \hat{g}_2|^2 = 4.3$ (0.28), $|\hat{g}_1|^2 = 4.2$ (0.35), and $|\hat{g}_2|^2 = 0.001$ (0.004) correspond to $L = L_{\max}/e$ ($L = L_{\max}/e^2$), with $\chi^2 = 8.5$ (10.2). The variations about the mean values in Eqs. (33) do not significantly alter these results.

Note that K and K^* exchanges do not contribute to the related reaction $\pi^- p \rightarrow \Sigma^- K^{*+}$, so that one would expect a considerably smaller cross section for this reaction than for $\pi^- p \rightarrow \Sigma^0 K^{*0}$ if the K^- and K^* -exchange model is valid. Experimentally, at 1.90 and 2.05 BeV/c incident-pion lab momenta, we have⁵ $\sigma(\pi^- p \rightarrow \Sigma^- K^{*+}) \approx 30\% \sigma(\pi^- p \rightarrow \Sigma^0 K^{*0})$. This indicates that other mechanisms are present, but the results are still consistent with the assumption that K and K^* exchange is the dominant mechanism for $\pi^- p \rightarrow \Sigma^0 K^{*0}$.

The above values for $|\hat{g}_0|^2$ will yield values for $|f_0(v_p^2)|^2$, since the experimental K^* width can be used to determine $|f_3(v_p^2)|^2$. In this section, we have used the Feynman matrix element

$$\mathcal{M}(\pi K K^*) = V_{K^*} \tau_\pi V_K \cdot (4\pi)^{1/2} f_3 r_\mu e^\mu(Q).$$

The total $K^{*0} \rightarrow K\pi$ decay rate is then

$$\Gamma = \frac{|\underline{r}|}{8\pi v_p^2} \cdot \frac{1}{3} \sum_{K^*, K, \pi} |\mathcal{M}(\pi K K^*)|^2,$$

where $\underline{r}$ is the decay-pion momentum in the K^* rest frame, and the

summation is over the K^* polarization and the K and π isotopic-spin indices. Thus,

$$\Gamma = |f_3(v_p^2)|^2 |\underline{r}|^3 / 2v_p^2 = (14.4 \text{ MeV}) |f_3(v_p^2)|^2.$$

By setting this decay rate equal to the experimental value of 50 MeV for the full width at half maximum for the K^* , we obtain $|f_3(v_p^2)|^2 = 3.5$. We thus obtain $|f_0(v_p^2)|^2 = |\hat{g}_0|^2 / |f_3(v_p^2)|^2 = 0.053, 0.16, \text{ and } 2.7$ for $\lambda = \infty, 2.06, \text{ and } 0.95 \text{ BeV}$, respectively. These values correspond to values 0.47, 0.45, and 0.29 (or alternatively to 0.53, 0.55, and 0.71) for the f parameter of Martin and Wali.⁶

ACKNOWLEDGMENTS

I would like to thank Professor Richard H. Dalitz and Dr. G. A. Smith for many helpful discussions, and to acknowledge the help of Jerome Friedman in programming the maximum-likelihood fit in Section IV.

FOOTNOTES AND REFERENCES

* Work done under the auspices of the U. S. Atomic Energy Commission

1. Gerald A. Smith, Joseph Schwartz, Donald H. Miller, George R. Kalbfleisch, Robert W. Huff, Orin I. Dahl, and Gideon Alexander, Phys. Rev. Letters 10, 138 (1963).
2. N. Byers and H. Burkhardt, Phys. Rev. 121, 281 (1961).
3. U. W. Cronin and O. E. Overseth, 1962 International Conference on High-Energy Physics at CERN (CERN, Geneva, 1962), p. 453.
4. The question of the renormalizability of field theories with vector particles does not really arise here. The results in this paper are in accord with the S-matrix philosophy, even though some of the terminology in the discussions is reminiscent of field theory.
5. Gerald A. Smith University of California, Lawrence Radiation Laboratory, Berkeley, California (unpublished).
6. A. W. Martin and K. C. Wali, Phys. Rev. 130, 2455 (1963).

FIGURE CAPTIONS

- Fig. 1. Diagrams that contribute to the reaction $\pi p \rightarrow \Lambda \omega$.
- Fig. 2. Typical diagrams that make up a three-particle vertex.
- Fig. 3. This diagram gives a self-energy correction to the K propagator instead of producing a second-order pole in the scattering amplitude.

MU-32002

Fig. 1.

MU-32003

Fig. 2.

MU-32004

Fig. 3.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.