

UCSF

UC San Francisco Previously Published Works

Title

Dependent Lung Opacity at Thin-Section CT: Evaluation by Spirometrically-Gated CT of the Influence of Lung Volume

Permalink

<https://escholarship.org/uc/item/9wn801gd>

Journal

Korean Journal of Radiology, 3(1)

ISSN

1229-6929

Authors

Lee, Ki-Nam
Yoon, Seong Kuk
Sohn, Choon Hee
[et al.](#)

Publication Date

2002

DOI

10.3348/kjr.2002.3.1.24

Peer reviewed

Dependent Lung Opacity at Thin-Section CT: Evaluation by Spirometrically-Gated CT of the Influence of Lung Volume

Ki-Nam Lee, MD¹
Seong Kuk Yoon, MD¹
Choon Hee Sohn, MD²
Pil Jo Choi, MD³
W. Richard Webb, MD⁴

Index terms :

Lung, collapse
Lung, CT
Lung, density
Lung, function
Lung, ventilation

Korean J Radiol 2002;3:24-29

Received August 6, 2001; accepted after revision January 29, 2002.

Department of ¹Diagnostic Radiology, ²Internal Medicine, ³Thoracic Surgery, Dong-A University College of Medicine; Department of ⁴Radiology, University of California, San Francisco

This paper was supported by the Dong-A University Research Fund, 1998.

Address reprint requests to :

Ki-Nam Lee, MD, Department of Diagnostic Radiology, Dong-A University College of Medicine, 3-1 Ga, Dong daesin-dong, Seo-gu, Pusan 602-103, Korea.
Telephone: (8251) 240-5375
Fax: (8251) 253-4931
e-mail: kinamlee@chollian.net

Objective: To evaluate the influence of lung volume on dependent lung opacity seen at thin-section CT.

Materials and Methods: In thirteen healthy volunteers, thin-section CT scans were performed at three levels (upper, mid, and lower portion of the lung) and at different lung volumes (10, 30, 50, and 100% vital capacity), using spirometric gated CT. Using a three-point scale, two radiologists determined whether dependent opacity was present, and estimated its degree. Regional lung attenuation at a level 2 cm above the diaphragm was determined using semiautomatic segmentation, and the diameter of a branch of the right lower posterior basal segmental artery was measured at each different vital capacity.

Results: At all three anatomic levels, dependent opacity occurred significantly more often at lower vital capacities (10, 30%) than at 100% vital capacity ($p = 0.001$). Visually estimated dependent opacity was significantly related to regional lung attenuation ($p < 0.0001$), which in dependent areas progressively increased as vital capacity decreased ($p < 0.0001$). The presence of dependent opacity and regional lung attenuation of a dependent area correlated significantly with increased diameter of a segmental arterial branch ($r = 0.493$ and $p = 0.0002$; $r = 0.486$ and $p = 0.0003$, respectively).

Conclusion: Visual estimation and CT measurements of dependent opacity obtained by semiautomatic segmentation are significantly influenced by lung volume and are related to vascular diameter.

Dependent opacity, an ill-defined increase in subpleural lung attenuation ranging from a few millimeters to a centimeter or more in thickness, and visible only in dependent lung regions, disappearing when the lung is nondependent, is often seen at high-resolution CT (1). By obtaining additional scans with patient prone, it can be readily distinguished from infiltrative lung disease (2–4).

The physical density of a lung depends upon the relative proportions of air, blood, extravascular fluid and lung tissue in the lung volume assessed (5). These vary with lung region and change continuously during normal respiration. It has been shown that dependent opacity arises in areas of passive microatelectasis (3) and has also been attributed to fluid accumulation caused by gravity (5–7). However, the relationship between visible dependent opacity and lung volume, lung levels, and lung attenuation has not been studied in detail. The purpose of this study is to evaluate the influence of lung volume and lung level on the presence of dependent opacity seen at thin-section CT, and to determine the relationship between dependent opacity and regional lung attenuation. In an attempt to estimate the effects of lung volume and pulmonary vascular diameter on dependent opacity, we also assessed the relationship between the

Influence of Lung Volume on the Extent of Dependent Lung Opacity at Thin-Section CT

presence and degree of dependent opacity and segmental arterial diameter at different lung volumes.

MATERIALS AND METHODS

Thirteen healthy volunteers, twelve men and one woman, ranging in age from 27 to 35 (mean, 29.7) years, participated in the study. Their height varied between 165 and 178 (mean, 171.2) cm and their body weight between 52 and 78 (mean, 67.5) Kg. They were non-smokers and had no history of pulmonary disease which might have affected the results of pulmonary function tests. Informed consent was obtained from all volunteers.

Thin-section CT scans of the chest were performed using a Somatom Plus-40 (Siemens, Erlangen, Germany). Images were obtained in the supine position at three different anatomic levels (carina, inferior pulmonary veins, and 2 cm above the right hemidiaphragm), and at lung volumes representing 10%, 30%, 50%, and 100% of vital capacity (determined using a respiratory gating device connected to the CT scanner). Vital capacity was defined as the volume expired after maximal inspiration, or inspired after maximal expiration. Following maximal inspiration, the subjects were instructed to exhale, respiration was interrupted mechanically by closing a valve in the spirometer at the selected lung volume, and a CT scan was obtained. In order to avoid motion artifacts due to straining against a closed

conduit system, the subjects was asked to avoid straining and to suspend respiration when the valve was closed. In order to familiarize each subject with the procedure, a few practice trials preceded actual image acquisition.

At each of the three different anatomic levels at which four lung volumes were determined, scans were obtained using 1-mm collimation. Scan time was 1 sec; tube current, 275 mA; and voltage, 137 kVp. Scans were reconstructed using a high-spatial-frequency algorithm, and window width and level were 1300 and -650 HU, respectively.

Dependent opacity was visually estimated by two radiologists (K.N.L, S.K.Y) who reviewed cases in random order and were blinded to the percent of vital capacity at which the scan was obtained (Table 1). The frequency and degree of dependent opacity seen at the three different anatomic levels at each different lung volume were assessed; at the level of the carina, assessment related to two areas separated by the major fissure, one in the dependent portion of the upper lobe and the other within the apex of the superior segment of the lower lobe. The findings were graded on a three-point scale (0, absent; 1, subtle; 2, obvious), and when two observers disagreed as to the presence or absence of dependent opacity, a decision was reached by consensus. For each different vital capacity, we also measured interobserver agreement as to the degree of dependent opacity at the three anatomic levels.

The relationship between the visually estimated degree

Table 1. Dependent Opacity Data Obtained by Semiautomatic Algorithm and Visual Estimate, and Arterial Diameter at Different Vital Capacities at a Level 2 cm above the Right Diaphragm in Thirteen Healthy Volunteers

Subjects No./Age(y)/ Sex	Attenuation as Determined by Semiautomatic Measurement (HU) in Different Vital Capacities				Visual Score of Dependent Opacity in Different Vital Capacities				Arterial Diameter (mm) in Different Vital Capacities			
	10%	30%	50%	100%	10%	30%	50%	100%	10%	30%	50%	100%
1/29/M	-660.7	-742.3	-824.7	-892	++	+	0	0	7.1	6.8	6.6	6.3
2/28/M	-609.7	-704.7	-772.5	-817.6	++	+	+	+	6.1	6.1	6.1	4.3
3/29/M	-623.3	-669.6	-812.6	-830.6	++	++	+	0	6.6	5.7	5.1	4.9
4/30/M	-552.5	-690.1	-702.8	-827.7	++	++	+	0	6.3	6.0	5.4	4.6
5/27/F	-624.7	-712.4	-784.6	-761.1	++	+	+	+	8.0	7.4	7.4	4.6
6/30/M	-610.8	-724.5	-807.3	-884	++	+	+	0	6.6	5.9	5.7	5.4
7/30/M	-595.4	-698.7	-750	-844.6	++	+	0	0	5.1	4.9	4.6	4.6
8/35/M	-617.4	-693.2	-748.5	-852.9	++	+	+	0	5.1	5.1	5.0	4.9
9/31/M	-573.3	-622.3	-713.4	-742.9	++	++	+	+	6.6	6.6	6.6	6.0
10/29/M	-538.1	-730.7	-792.7	-849.3	++	+	+	0	6.6	5.4	4.9	4.3
11/30/M	-613.7	-712.1	-803.6	-893.6	++	+	0	0	5.4	4.3	4.0	4.0
12/30/M	-667.4	-702.3	-856.3	-903.6	++	+	+	0	5.1	4.7	4.3	4.3
13/28/M	-642.4	-710.5	-757.7	-841.9	++	+	+	0	4.9	4.8	4.3	3.4
Mean	-610.0	-701.0	-779.0	-841.7					6.1	5.7	5.4	4.7
SD	38.1	30.0	44.0	48.5					0.9	0.9	1.0	0.8

Note.— ++: (grade 2) indicates that obvious dependent opacity was recognized by two radiologists, +: (grade 1) denotes subtle dependent opacity, 0: signifies the absence of dependent opacity.

of dependent opacity and regional CT attenuation was determined, being calculated semiautomatically as described by other investigators (8–13). For each scan, the “Pulmo CT” option from Siemens was used to calculate a region of interest (9, 10). For measuring attenuation, a region of interest was selected in the most posterior area at a level 2 cm above the right hemidiaphragm. The region of interest chosen was a zone of the right lung posterior to a horizontal line drawn 1 cm from the most dependent pleural surface. Thus, a crescent-shaped area (average 3.6 cm² ± 0.6) was selected as the most dependent lung region (Fig. 1). We compared regional CT attenuation measured at different lung volumes (10%, 30%, 50%, and 100% of vital capacity) (Table 1).

The diameter of a branch of the right posterior basal segmental artery, chosen in order to avoid artifact-related cardiac pulsation, was measured three times at the different respiratory levels, using a Vernier caliper (millimeters, with two decimal places), and averaged. The relationship between the degree of dependent opacity and arterial diameter at these different lung volumes was then assessed.

For statistical analysis, a microcomputer and SAS software (SAS Institute, Cary, N.C., U.S.A.) were employed, and for statistical comparisons of differences in visually estimated dependent opacity and regional CT attenuation according to vital capacity, we used the Mantel-Haenszel Chi-Square test. For correlation of the degree of visually estimated dependent opacity or attenuation of a dependent area with the diameter of a segmental arterial branch, Pearson’s correlation coefficient was used. A p value of

less than 0.05 was considered statistically significant. To determine interobserver variability, kappa statistics were used: values up to 0.4 indicated positive but poor agreement; values of 0.41–0.75, good agreement; and values greater than 0.75, excellent agreement. A value of 1 demonstrated perfect agreement.

RESULTS

Dependent opacity was found to be present significantly more often at lower vital capacity levels (in particular, 10% or 30%) than at 100% of vital capacity (*p* = 0.001) (Table 2). Both observers noted that at the three different anatomic levels, obvious dependent opacity was 10% of vital capacity in all cases. Interobserver agreement in the visual assessment of dependent opacity was excellent (*K* = 0.82) (Table 3). Although dependent opacity occurred at all three different anatomic levels, it was seen more fre-

Table 2. The Occurrence Rate of Dependent Opacity at Three Different Anatomic Levels according to Vital Capacity

Vital Capacity	Anatomic Level		
	Carina	Inferior Pulmonary Vein	2 cm Above Diaphragm
10%	100	100	100
30%	61.5	84.6	100
50%	23.1	76.9	76.9
100%	15.4	15.4	23.1

Note.—The figures are percentages.

Table 3. K Values for Reader 1 versus Reader 2 at Three Anatomic Levels and Different Vital Capacities

Anatomic Level	Vital Capacity			
	10%	30%	50%	100%
Carina	1	0.693	0.822	0.788
Inferior pulmonary vein	1	0.567	0.693	0.822
2 cm above diaphragm	1	0.788	0.693	0.822

Table 4. The Occurrence of Dependent Opacity in Two Different Dependent Areas at the Level of the Carina

Vital Capacity	Posterior Segment of Right Upper Lobe	Superior Segment of Right Lower Lobe
	10%	13 (100)
30%	8 (61.5)	5 (38.5)
50%	3 (23.1)	1 (7.7)
100%	2 (15.4)	0 (0)

Note.—The figures in parentheses are percentages.




Fig. 1. The region of interest for measurements of regional lung attenuation was determined as a zone of the right lung posterior to a horizontal line drawn 1 cm from the most dependent lung. For the measurement of regional lung attenuation, we selected the same area (3.6 ± 0.6 cm²) as the most dependent area of a crescent shape (arrows).

Influence of Lung Volume on the Extent of Dependent Lung Opacity at Thin-Section CT

quent 2 cm above the diaphragm ($p = 0.001$) (Fig. 2). At the level of the carina, at 100% of vital capacity, dependent opacity was noted in two (15%) of 13 cases in the dependent portion of the right upper lobe, though was not visible at the apex of the superior segment of the lower lobe (Table 4).

When dependent opacity was present, measured regional attenuation at a level 2 cm above the right hemidiaphragm ranged from -538.1 to -856.3 (mean, -695.4 ; SD 76.9)

HU (Table 1). The degree of dependent opacity assessed visually was significantly related to the mean value of the regional attenuation of dependent lung ($p = 0.001$). Furthermore, regional CT attenuation in a dependent lung increased progressively as vital capacity decreased ($p = 0.001$).

The degree of dependent opacity correlated with the diameter of a segmental arterial branch ($r = 0.493$; $p = 0.0002$) (Table 5), and the CT attenuation of a dependent

Table 5. Comparison of Visually Estimated Dependent Opacity with Regional Lung Attenuation and Arterial Diameter in Dependent Areas (n=52) at a level 2 cm above the Right Diaphragm

	Degree of Dependent Opacity		
	Grade 2 (n=16)	Grade 1 (n=23)	Grade 0 (n=13)
Regional lung attenuation [†]	-619.5 ± 41.7 ($-538.1 - -669.6$)	-747.9 ± 46.7 ($-693.2 - -856.3$)	-846 ± 42.1 ($-750 - -903.6$)
Diameter of segmental artery [‡] (mm)	6.1 ± 0.9 (4.9 - 8.0)	5.4 ± 1.0 (4.3 - 7.4)	4.8 ± 0.9 (3.4 - 6.6)

Note.—[†]Mean (\pm SD) HU. [‡]Mean (\pm SD) diameters (mm) of the artery in the right lower posterior basal segment. The figures in parentheses indicate the range.




Fig. 2. Thirty-year-old man who showed dependent opacity (grade 2) at 10% vital capacity.
A. Dependent opacity occurred at the level of the carina. Lung attenuation is greater in the posterior aspect of the upper lobes than in the superior segment of the lower lobes, located more posteriorly. Thus, dependent opacity is influenced not only by gravitational effects.
B. Dependent opacity is seen at the level of the inferior pulmonary vein.
C. Dependent opacity is prominent at a level 2 cm above the diaphragm.




Fig. 3. Comparison of dependent opacity and arterial diameter in a 29-year-old man
A. Dependent opacity was obvious (grade 2) at 10% of vital capacity, and arterial diameter in the right posterior basal segment was 7.1 mm (artery, arrow).
B. At 30% of vital capacity, dependent opacity was graded as subtle (grade 1) and arterial diameter measured 6.8 mm (arrow).
C. Dependent opacity was not apparent at 50% of vital capacity (score 0), at which point arterial diameter was 6.6 mm (arrow).
D. Dependent opacity was absent at 100% vital capacity, and at the same level, arterial diameter was 6.3 mm (arrow).

area correlated with arterial diameter ($r = 0.486$; $p = 0.0003$). As vital capacity decreased, the frequency of visible dependent opacity and arterial diameter both increased (Fig. 3).

DISCUSSION

Dependent opacity, also termed subpleural dependent density, consists of a stripe of ground glass attenuation 2 to 30 mm thick and is usually seen in the most dependent lung regions. It disappears when the lung is nondependent (3). Verschakelen et al. (8) studied the differences in CT attenuation found between dependent and nondependent lung regions, noting that the difference was smallest for lung volumes near total vital capacity. Conversely, when lung volume was reduced, as in full expiration, differences in CT attenuation between dependent and nondependent portions of the lung increased.

The physical density of lung is determined by the relative contribution of its various components: air, blood, extravascular fluid, and lung tissue. Regional differences in lung attenuation are related to lung volume and the effects of gravity (5). In normal subjects, a gradient in CT lung attenuation (attributed primarily to the influence of gravity on blood flow and lung inflation) is observed, with lung attenuation greater in dependent lung (6). Furthermore, largely because of regional changes in lung volume and lung perfusion, lung attenuation changes continuously dur-

ing respiration (8–10, 12–15).

However, variations in scan parameters and sampling methods used in prior studies, and the possible influence of lung pathology on regional lung density measurements, must be considered when assessing the results of these studies (8–10, 12–15). Recently, semiautomatic procedures for the measurement of lung attenuation, as well as spirometric gated CT, have proven useful in the objective assessment of lung attenuation at varying lung volumes (8–11, 14, 15). Spirometric gated CT allows for the selection of a specific point in the respiratory cycle at which flow can be interrupted by closing a valve in the spirometer prior to imaging (8, 9). Also, in the present study normal non-smoking volunteers were selected to avoid the possibility of lung disease influencing the measurement of lung attenuation.

As reported in prior studies, we found that dependent opacity was apparent significantly more often at lower lung volumes (8, 12), and was also more frequent at lung bases. It was visualized in the dependent portions of both right and left lungs at each anatomic level studied (carina, inferior pulmonary veins, and 2 cm above the right diaphragm). However, at the level of the carina, dependent opacity was significantly more frequent in the posterior segment of the upper lobe (anterior to the major fissure) than in the superior segment of the lower lobe (posterior to the major fissure and thus more dependent). This suggests that, at least in this region, factors other than gravity affect

the occurrence of dependent opacity.

We attempted to determine whether the presence and degree of visually estimated dependent opacity was related to regional lung attenuation. A crescent-shaped region of interest measuring $3.6 \pm 0.6 \text{ cm}^2$ was semiautomatically located in the dependent portion of the right posterior basal segment. Visually estimated dependent opacity was present at a wide range of lung attenuation values (range, -856.3 to -538.1 HU; mean, -695.4 ± 76.9 HU), reflecting the varying lung volumes at which assessment was made and the fact that the visibility of dependent opacity is determined not only by the absolute density of dependent lung but also by its relationship to the density of nondependent lung regions. In a study using spirometrically gated CT, mean lung attenuation in normal subjects has been shown to be approximately -760 HU at 20% of vital capacity, -835 HU at 50%, and -860 HU at 80% (9). However, the degree of dependent opacity correlated well with regional attenuation regardless of lung volume.

The gradient in attenuation between dependent and nondependent zones can be explained by gravitational differences in perfusion and inflation. Because lung perfusion is greater in dependent zones, the arteries in these areas are larger. In assessing the relationship between the degree of dependent opacity and the diameter of a segmental artery, we measured arterial diameter in the posterior basal segment of the right lower lobe, a region where dependent opacity is often observed. In our study, there was a significant relationship between increased arterial diameter measured in the right posterior basal segment and the occurrence of dependent opacity and lower lung volumes.

There are several limitations to our study. The subjects were young, normal, healthy volunteers, among whom there may have been interindividual differences in physiologic respiration in the regional cross-section of lung density. All volunteers except one were men, and although this may have introduced a gender bias, it was our desire to study patients with a relatively small range of body habitus. Three female volunteers less than 160 cm and weighing less than 48 kg were excluded.

In conclusion, dependent opacity is significantly influenced by respiratory level, which must be considered when lung attenuation is evaluated, and is related to lung attenuation and vascular diameter. The physiologic phenomenon of varying lung density associated with respiration can be demonstrated using respiratory gated CT.

References

1. Webb WR, Muller NL, Naidich DP. *An illustrated glossary of HRCT*. In: *High-resolution CT of the lung*. 2nd ed. Philadelphia: Lippincott-Raven, 1996; 295-311
2. Primack SL, Remy-Jardin M, Remy J, Muller NL. High-resolution CT of the lungs: pitfalls in the diagnosis of infiltrative lung disease. *AJR* 1996; 167:413-418
3. Aberle DR, Gamsu G, Ray CS, Feuerstein IM. Asbestos-related pleural and parenchymal fibrosis: detection with high-resolution CT. *Radiology* 1988; 166:729-734
4. Volpe J, Storto ML, Lee K, Webb WR. High-resolution CT of the lung: determination of the usefulness of CT scans obtained with the patient prone based on plain radiographic findings. *AJR* 1997; 169:369-374
5. Hedlund LW, Vock P, Effmann EL. Evaluating lung density by computed tomography. *Semin Respir Med* 1983; 5:76-88
6. Rosenblum LJ, Mauceci RA, Wellenstein DE, Bassano DA, Cohen WN, Heitzman ER. Density patterns in the normal lung as determined by computed tomography. *Radiology* 1980; 137:409-416
7. Tokics L, Hedenstierna G, Strandberg A, Brismar B, Lundquist H. Lung collapse and gas exchange during general anesthesia: effects of spontaneous breathing, muscle paralysis and positive end-expiratory pressure. *Anesthesiology* 1987; 66:157-167
8. Verschakelen JA, Van Fraeyenhoven L, Laureys G, Demedts M, Baert AL. Differences in CT density between dependent and nondependent portions of the lung: influence of lung volume. *AJR* 1993; 161:713-717
9. Kalender WA, Rienmuller R, Seissler W, Behr J, Welke M, Fichte H. Measurement of pulmonary parenchymal attenuation: use of spirometric gating with quantitative CT. *Radiology* 1990; 175:265-268
10. Rienmuller RK, Behr J, Kalender WA, et al. Standardized quantitative high-resolution CT in lung diseases. *J Comput Assist Tomogr* 1991; 15:742-749
11. Kalender WA, Fichte H, Bautz W, Skalej M. Semiautomatic evaluation procedures for quantitative CT of the lung. *J Comput Assist Tomogr* 1991; 15:248-255
12. Gevenois PA, Scillia P, de Maertelaer V, Michils A, Vuyst PD, Yernault J. The effects of age, sex, lung size and hyperinflation on CT lung densitometry. *AJR* 1996; 167:1169-1173
13. Wegner OH, Koeppel P, Oeser H. Measurement of lung density by computed tomography. *J Comput Assist Tomogr* 1978; 2:263-273
14. Wandtke JC, Hyde RW, Fahey PJ, et al. Measurement of lung gas volume and regional density by computed tomography in dogs. *Invest Radiol* 1986; 21:108-117
15. Kohz P, Stabler A, Beinert T, et al. Reproducibility of quantitative, spirometrically controlled CT. *Radiology* 1995; 197:539-542
16. Coxon HO, Mayo JR, Behzad H, et al. Measurement of lung expansion with computed tomography and comparison with quantitative histology. *J Appl Physiol* 1995; 79:1525-1530