UC San Diego

UC San Diego Previously Published Works

Title

Behavioral and Physical Activity Interventions for HAND.

Permalink

<https://escholarship.org/uc/item/9x698018>

Authors

Montoya, Jessica L
Henry, Brook
Moore, David J

Publication Date

2021

DOI

10.1007/7854_2018_79

Peer reviewed

Behavioral and Physical Activity Interventions for HAND

Jessica L. Montoya, Brook Henry, and David J. Moore

Contents

- 1 HIV-Associated Neurocognitive Disorders
- 2 Physical Activity as a Behavioral Intervention for HAND
 - 2.1 Physical Activity Levels Among People Living with HIV
 - 2.2 Relationship Between Physical Activity and Neurocognition
 - 2.3 Underlying Biologic Mechanisms Linking Physical Activity and Neurocognition
 - 2.4 Interventions to Increase Physical Activity, Thereby Benefiting Neurocognition
- 3 Diet as a Behavioral Intervention for HAND
 - 3.1 Diet and Neurocognitive Function in the General Population
 - 3.2 Diet and Nutrition Concerns in the Context of HIV
 - 3.3 Diet Interventions Specific to People Living with HIV
- 4 Sleep as a Behavioral Intervention for HAND
 - 4.1 Sleep Disturbance in the Context of HIV
 - 4.2 Effect of Sleep Disturbance on Neurocognitive Function
 - 4.3 Behavioral Strategies to Improve Sleep Quality
- 5 Promotion of Antiretroviral Therapy Adherence as a Behavioral Intervention for HAND
 - 5.1 Antiretroviral Therapy Adherence and Neurocognitive Performance
 - 5.2 Strategies to Promote Antiretroviral Therapy Adherence
- 6 Clinical Implications
- 7 Conclusion
- References

Abstract Approximately 30–50% of persons living with HIV manifest some degree of neurocognitive impairment. Even mild-to-moderate forms of HIV-associated neurocognitive disorders (HAND) can result in difficulties with everyday functioning, such as suboptimal medication adherence and impaired driving. Despite the pervasive presence and consequences of HAND, there is a significant unmet need to develop

J. L. Montoya and B. Henry
Department of Psychiatry, University of California San Diego, La Jolla, CA, USA

D. J. Moore (✉)
Department of Psychiatry, University of California San Diego, La Jolla, CA, USA
HIV Neurobehavioral Research Program, San Diego, CA, USA
e-mail: djmoore@ucsd.edu

effective behavioral strategies to reduce the incidence and consequences of HAND. Although there is an absence of evidence-based behavioral interventions specific to HAND, the literature reviewed in this chapter suggest the following modifiable lifestyle factors as intervention targets: physical activity, diet, sleep, and antiretroviral medication adherence. Adoption and maintenance of these healthy lifestyle factors may reduce inflammation and oxidative stress, which, in turn, may reduce the incidence and/or severity of HAND.

Keywords Antiretroviral therapy adherence · Cognition · Diet · Exercise · Nutrition · Physical activity · Sleep

1 HIV-Associated Neurocognitive Disorders

Approximately 30–50% of persons living with Human Immunodeficiency Virus (HIV) Type-1 manifest some degree of neurocognitive impairment, typically characterized by deficits in executive function, attention, and memory referred to as HIV-associated neurocognitive disorders (HAND) (Heaton et al. 2010). HAND per Frascati diagnostic criteria includes three categories: asymptomatic neurocognitive impairment (ANI), HIV-associated mild neurocognitive disorder (MND), which includes self-reported deficits in everyday functioning, and the most severe of the disorders, HIV-associated dementia (HAD) (Antinori et al. 2007). While HAD prevalence has declined with the use of effective antiretroviral therapy, ANI and MND, marked by impaired performance in at least two neurocognitive domains (greater than one standard deviation below the mean of normative scores), remain a prevalent and significant public health concern. Even mild-to-moderate forms of HAND translate to higher risk of impaired driving, difficulties with instrumental activities of daily living, and poorer antiretroviral therapy adherence (Thames et al. 2011; Vance et al. 2011; Marcotte et al. 2004). Despite the pervasive presence and consequences of HAND, there is a significant unmet need to develop effective behavioral strategies to reduce the incidence and consequences of HAND (Woods et al. 2009b). Among existing behavioral interventions, promotion of physical activity, a healthy diet, improved sleep, and antiretroviral therapy adherence may benefit neurocognitive function.

2 Physical Activity as a Behavioral Intervention for HAND

2.1 *Physical Activity Levels Among People Living with HIV*

Physical activity, traditionally defined as the movement of skeletal muscles that requires energy expenditure, has been recommended as a safe therapeutic strategy to sustain and maintain the health of persons living with HIV (Botros et al. 2012). Physical activity interventions among people living with HIV are reported to

improve body composition, muscle strength, aerobic fitness, and quality of life; however, there are notable barriers to physical activity among people living with HIV, including disease symptoms, antiretroviral therapy side effects, depression, and pain (Henry and Moore 2016; Vancampfort et al. 2018b). Surprisingly, relatively few studies have examined physical activity among people living with HIV. A 2012 review indicated that the diversity and inconsistency of methods used to assess physical activity (typically a variety of self-report questionnaires) precluded any pooled estimate of overall physical activity among people living with HIV, although the range of estimates suggest that 19–73% of people living with HIV are “sedentary,” as defined by various criteria (Schuelter-Trevisol et al. 2012). Webel and colleagues reported that an Ohio cohort of persons living with HIV tended to exercise regularly, but the mean level of physical activity was lower in women than men, and below recommended levels (i.e., 150 min per week of moderate-to-vigorous physical activity) (Webel et al. 2015). A sample of 50 African-American females living with HIV from the southern United States exhibited low levels of physical activity, with only one person exceeding 150 min per week of moderate-to-vigorous physical activity (Rehm and Konkle-Parker 2016). In contrast, two-thirds of a cohort of people living with HIV from Vietnam were rated as physically active on the International Physical Activity Questionnaire, although this group did include rural participants with potentially more endogenous physical activity than urban residents (Dang et al. 2018). A recent meta-analysis reviewed physical activity data from 24 studies of people living with HIV conducted across the world (Vancampfort et al. 2018a). To optimize consistency, the authors included physical activity results obtained only through the International Physical Activity Questionnaire or objective measures, such as an accelerometer. The findings from this seminal study showed that approximately 50% of people living with HIV failed to meet suggested physical activity guidelines (i.e., exhibiting less than 150 min per week of moderate-to-vigorous physical activity). Furthermore, average daily steps, when collected, was about 5,800 per day, which is far below the 10,000 steps recommended for adults and close to a sedentary range (typically defined as below 5,000 steps per day). In summary, it is clear that physical activity habits among people living with HIV may vary significantly depending upon geographic and demographic factors, but studies to date suggest that a large proportion of people living with HIV are not sufficiently physically active. As a result, physical activity interventions may be beneficial for numerous persons living with HIV.

2.2 Relationship Between Physical Activity and Neurocognition

Moderate- and vigorous-intensity physical activity has beneficial effects on neurocognitive function (Engeroff et al. 2018). Among healthy individuals and children, physical activity is shown to improve neurocognitive functioning (Davis et al. 2007; Best et al. 2015; Mura et al. 2015; Guiney and Machado 2013; Gill et al. 2015).

Aerobic exercise is associated with improvement in multiple neurocognitive domains, including attention, processing speed, executive function, and memory (Smith et al. 2010). Physical activity may also benefit global neurocognition in patients with mild neurocognitive impairment (Song et al. 2018) or history of stroke (Vanderbeken and Kerckhofs 2017), and may slow neurocognitive decline in Alzheimer's disease (Farina et al. 2014). In the context of HIV, higher engagement in physical activity is associated with better executive function (Ortega et al. 2015) and reduced likelihood of neurocognitive impairment in cross-sectional studies (Dufour et al. 2013; Fazeli et al. 2015). Longitudinally, persons living with HIV who consistently engage in physical activity begin with, and maintain, significantly better neurocognitive function compared to persons living with HIV who do not engage in physical activity or do so inconsistently (Dufour et al. 2018). In a longitudinal Multicenter AIDS Cohort Study, high engagement in physical activity was associated with lower odds of impairment in the domains of learning, memory, and motor function, and these effects were found to be more pronounced among the sample of men living with HIV compared to the pooled sample of men living with HIV and HIV-uninfected men (Monroe et al. 2017). In a scoping review of the effect of physical activity on neurocognitive function among people living with HIV (Quigley et al. 2018), noninterventional studies reported a positive association between physical activity engagement in neurocognitive function, whereas there was a dearth of positive outcomes of aerobic interventions on neurocognition. Null results of interventional studies potentially may be a result of methodological factors (e.g., low prescribed doses of physical activity, recruitment of relatively young people living with HIV who are at lower risk for neurocognitive impairment compared to older people living with HIV, and reliance on self-reported versus objective measures of neurocognitive function). Thus, future randomized controlled trials should prescribe doses of physical activity consistent with current recommendations (150 min per week of moderate-to-vigorous physical activity), recruit participants most likely to benefit from physical activity interventions, and utilize comprehensive neurocognitive assessments.

2.3 Underlying Biologic Mechanisms Linking Physical Activity and Neurocognition

Chronic inflammation is established early in HIV infection and is postulated to contribute to neurocognitive impairment across the age span of persons living with HIV (Sattler et al. 2015; Hong and Banks 2015; Tavazzi et al. 2014; Kapetanovic et al. 2010, 2014; Ancuta et al. 2008; Gannon et al. 2011). Physical activity is known to exert anti-inflammatory effects, which may be a consequence of reduced visceral fat and decreased release of adipokines (d'Ettorre et al. 2014). In addition to reduced inflammation, some of the underlying biologic mechanisms linking physical activity to neurocognition include promotion of cerebral angiogenesis, improved cerebral and peripheral vascular reactivity (i.e., increased maximal oxygen consumption),

upregulation of neurotrophins, increased neurogenesis, and decreased hippocampus apoptosis (Stimpson et al. 2018). Imaging studies indicate that engagement in physical activity can increase brain volume, and biomarker studies show that physical activity leads to upregulation of brain-derived neurotrophic factor (BDNF) (Firth et al. 2017). A randomized controlled trial showed that among individuals with schizophrenia, those who participated in an aerobic exercise intervention (compared to treatment as usual) showed improvements in physical fitness and neurocognitive function (Kimhy et al. 2015). Furthermore, enhancement of physical fitness and increases in BDNF accounted for much of the variance in neurocognitive improvement, supporting the hypothesis that physical activity-induced upregulation of BDNF may contribute to improved neurocognitive outcomes.

2.4 Interventions to Increase Physical Activity, Thereby Benefiting Neurocognition

Low-to-moderate intensity walking interventions have demonstrated effectiveness in improving neurocognitive performance in older adults (Rosenberg et al. 2012) and neurocognitively impaired populations (Kemoun et al. 2010). Compared to moderate physical activity, high intensity interval training, which involves short periods of exercise performed at high intensity (greater than 80–85% peak oxygen uptake), may have greater benefits for improving cardiometabolic function (e.g., improved insulin and glucose regulation and reduced inflammation markers, such as interleukin-6 and C-reactive protein) (Tjonna et al. 2009; Munk et al. 2011; Ramos et al. 2015). Interventions employing a combination of aerobic and resistance exercise also show evidence of improving body composition and lipid profile in people living with HIV (Hand et al. 2009; O'Brien et al. 2010). However, the effect of a combined aerobic and resistance exercise regimens on neurocognition has not been investigated as an intervention strategy for HAND.

Limitations of many physical activity interventions are issues of feasibility and scalability. Many intervention studies are conducted at gym facilities, which may not be accessible in nonintervention contexts to many persons living with HIV who may face limited financial resources (Montoya et al. 2015). To date, few studies have specifically examined the effect of a physical activity intervention on neurocognition among people living with HIV. One recent 16-week protocol required participants to attend three aerobic exercise sessions at a gym facility each week, but did not observe any effects of the physical activity intervention on neurocognition relative to a control group (McDermott et al. 2017). However, the sample size was extremely small; only 11 people participated in total (6 control and 5 intervention participants completed the study out of 57 persons living with HIV screened), individuals did not attend 40% of the planned physical activity training sessions, and the neurocognitive assessment was limited, including only the Montreal Cognitive Assessment and

Trail-making A and B tests. These issues highlight some of the challenges involved in conducting physical activity interventions to improve neurocognitive outcomes. Although physical activity that is gym-based may not be feasible for many persons living with HIV, increasing walking/step count may be an appropriate treatment target. Walking may reduce sedentary behavior and improve metabolic function (Healy et al. 2008; Manson et al. 1999). For example, an increase in mean daily step counts by approximately 2,000 improves lipid profile (Sugiura et al. 2002).

In the general population, mobile health interventions have been developed to promote engagement in physical activity, such as walking (Fjeldsoe et al. 2009). An ongoing randomized controlled trial is investigating whether a novel and personalized text messaging intervention (iSTEP) can significantly increase moderate physical activity in people living with HIV (R21MH100968) (Henry and Moore 2016). Second, the iSTEP intervention aims to evaluate the effect of physical activity engagement on neurocognitive performance. Preliminary data supports high text message response rates and positive participant feedback (Henry and Moore 2016).

3 Diet as a Behavioral Intervention for HAND

3.1 Diet and Neurocognitive Function in the General Population

Epidemiological research indicates that a healthy diet, such as the Mediterranean diet and consumption of omega-3 fatty acids (e.g., docosahexaenoic acid, DHA), may help prevent neurocognitive decline (Barak and Aizenberg 2010). The Mediterranean diet is characterized by high intake of vegetables, legumes, fruits, nuts, cereals, and olive oil; moderate intake of fish and alcohol; low-to-moderate intake of dairy products; and low intake of saturated lipids and meat (Loughrey et al. 2017). A recent systematic review indicated a positive, concurrent association between the Mediterranean diet and global neurocognition; however, the association between the Mediterranean diet and specific neurocognitive domains is less established (Knight et al. 2017). Methodological differences in relation to neuropsychological assessment are a likely factor contributing to the lack of consensus among studies on the relationship between the Mediterranean diet and neurocognitive function (Knight et al. 2017). Similar to the Mediterranean diet, diets rich in polyphenols – found in fruits, vegetables, tea, wine, juices, plants, and some herbs – may promote better performance in neurocognitive abilities in a dose-dependent manner (Nurk et al. 2009) and slower rates of neurocognitive decline (Devore et al. 2012) in older persons. Results from the Framingham Heart Study (Schaefer et al. 2006) indicate individuals with higher DHA levels had a relative risk of 0.61 of developing Alzheimer’s disease, compared to individuals with lower

DHA levels, after adjustment for relevant covariates (i.e., sex, apolipoprotein E allele, plasma homocysteine concentration, and education level). Higher DHA levels translated to a mean DHA intake of 180 mg/day and a mean fish intake of 3.0 servings per week.

3.2 Diet and Nutrition Concerns in the Context of HIV

The introduction of effective antiretroviral therapy resulted in improvements in nutritional status and weight gain among people living with HIV (Leyes et al. 2008). However, body composition changes (i.e., lipodystrophy) persist even with effective antiretroviral therapy, particularly among people living with HIV on protease inhibitors (Carr et al. 1998). Body composition changes may include intra-abdominal fat accumulation (Miller et al. 1998), which is of relevance to pathogenesis of HAND given that the HIV CHARTER cohort study has found a link between abdominal obesity and neurocognitive function (Sattler et al. 2015). Multiple studies indicate that cardiometabolic dysfunction (e.g., abdominal obesity, hyperglycemia, dyslipidemia, and hypertension) confers risk for HAND and correlates with imaging markers of neurochemical abnormalities and neuroinflammation among persons living with HIV (Cysique et al. 2013; McCutchan et al. 2012; Saylor et al. 2016; Valcour et al. 2005).

The nutritional status of persons living with HIV can influence body composition changes, but the efficacy of diet interventions to treat metabolic conditions, thereby improving neurocognitive outcomes is limited (Leyes et al. 2008). Despite limitations in the literature to be able to tease apart the various contributors of metabolic changes observed among people living with HIV, cross-sectional studies indicate that diets high in cholesterol and saturated and trans fats play a role in metabolic disturbances (Shah et al. 2005; Hadigan et al. 2001). Although diet and nutrient deficiencies have been extensively documented for persons living with HIV in low-income countries, with a focus on treatment with supplements such as vitamins (Duran et al. 2008), less work has been performed to characterize the diet quality for persons living with HIV in developed countries such as the United States. A couple of reports indicate that youth living with HIV demonstrate a lower Healthy Eating Index score, higher fat intake, and lower micronutrient consumption (including vitamins A, E, calcium, and potassium) compared to an age-matched HIV-seronegative comparison group (Kruzich et al. 2004; Ziegler et al. 2014). Pregnant women living with HIV in their third trimester also exhibited low Healthy Eating Index scores, although individuals born outside the United States had better scores, perhaps indicative of more unhealthy diet practices adopted by people living in the United States (Miller et al. 2017). One cohort of adults living with HIV (53% male) also consumed higher levels of saturated fat and lower amounts of polyunsaturated fat and fiber than recommended by the National Cholesterol Education Program (Capili and Anastasi 2008). While more data needs to be obtained,

the existing evidence suggests that interventions that address diet content as well as physical activity may be helpful for people living with HIV in the United States.

3.3 Diet Interventions Specific to People Living with HIV

Current international guidelines recommend dietary intervention as a first-line treatment for HIV-related dyslipidemia (Dube et al. 2003), but the results of clinical trials have been inconsistent. Persons living with HIV who receive macronutrient (protein/carbohydrate) or micronutrient (vitamin) supplements in randomized controlled trials often do not show a significant reduction in morbidity, mortality, or disease progression (e.g., CD4 count and viral load) (Grobler et al. 2013; Visser et al. 2017). Interventions that have promoted low-fat diets are reported to have beneficial but limited effects, such as a reduction in triglycerides but not cholesterol (Stradling et al. 2012). Some nutrition studies have focused on polyunsaturated fatty acids (PUFA) and omega-3 PUFA (n-3 PUFA), which may benefit brain health and function and are emphasized as part of the Mediterranean diet (Poulose et al. 2014). PUFAs may contribute to optimal brain function by reducing oxidative stress and inflammation, maintaining neuronal membrane integrity, and attenuating protein aggregation implicated in neurodegenerative diseases and age-related neurocognitive decline (Poulose et al. 2014). Along these lines, a large 12-month study (PREDIMED) [7 years and 8,000 participants (Zazpe et al. 2008)] indicated that adoption of the Mediterranean diet provided significantly more benefits compared to control or a low-fat diet condition, including a reduction in cardiovascular disease events, decreased markers of oxidative stress, and improved neurocognition (Estruch et al. 2013; Schroder et al. 2014; Fito et al. 2014; Mitjavila et al. 2013; Martinez-Lapiscina et al. 2013). Despite potential neurocognitive benefits, the Mediterranean diet has not been evaluated as a strategy for preventing or treating HAND, although greater Mediterranean diet adherence is associated with improved metabolic function (lower insulin resistance and higher high-density lipoprotein cholesterol) in people living with HIV (Tsiodras et al. 2009). Future studies may examine the effect of the Mediterranean diet on cardiovascular risk and neurocognitive deficits associated with HIV.

4 Sleep as a Behavioral Intervention for HAND

4.1 Sleep Disturbance in the Context of HIV

Sleep disturbance and fatigue are common symptoms reported by persons living with HIV, with up to 75% of people living with HIV experiencing sleep disturbance according to the Pittsburgh Sleep Quality Index (Rubinstein and Selwyn 1998) and up to 88% experience fatigue (Jong et al. 2010). A meta-analysis of sleep disturbances among people living with HIV calculated that the overall prevalence

of self-reported sleep disturbance was 58% (Wu et al. 2015). Furthermore, more than half of a sample population of people living with HIV reported symptoms related to sleep disturbance and fatigue, such as lack of energy (65%), drowsiness (57%), and difficulty sleeping (56%) (Lee et al. 2009). In a Taiwanese study, people living with HIV had a 3.74-fold higher risk of sleep disturbances compared to a general population control group (Chen et al. 2017). Among cohorts of people living with HIV, sleep disturbance and/or fatigue severity are associated with depression and anxiety symptoms (Millikin et al. 2003; Jong et al. 2010); antiretroviral therapy medication types and family and social support (Ren et al. 2018); high levels of HIV-RNA (greater than 1,000 copies/mL) (Womack et al. 2017); and substantial night-to-night variability in bedtime and risetime (Taibi et al. 2013).

4.2 Effect of Sleep Disturbance on Neurocognitive Function

Studies on the association between self-reported sleep disturbance and neurocognition have been mixed but suggest a link between poor sleep quality and worse neurocognitive outcomes (Yaffe et al. 2014). A meta-analysis of the impact of short-term sleep deprivation on neurocognition indicated that sleep deprivation had the largest effect on simple attention tasks (e.g., Psychomotor Vigilance Test and other simple reaction time tests), with tasks of greater complexity being affected to a lesser degree after sleep deprivation (Lim and Dinges 2010). On the other end of the spectrum, longer total sleep time (e.g., greater than or equal to 10 h of sleep/night) is also reported to significantly influence neurocognition, such as lower scores on the Mini-Mental Status Examination (Faubel et al. 2009) and lower performance on a recall test (Xu et al. 2011). Other measures of sleep disturbance (i.e., lower sleep efficiency, higher levels of wake after sleep onset, and a higher number of long wake episodes) have been related to decline in executive function as measured by the Trails B Test over an average study period of 3.4 years in a community-dwelling cohort of older men (mean age 76.0) (Blackwell et al. 2014). In a cohort of people living with HIV, 63% reported poor sleep quality based on a cutoff point of 5 on the Pittsburgh Sleep Quality Index (Byun et al. 2016). In this same sample, poorer subjective sleep quality, shorter or longer total sleep time measured by actigraphy (i.e., fewer than 7 h or greater than 8 h vs 7–8 h), and greater morning fatigue were associated with self-reported problems with neurocognitive function (e.g., difficulties with reasoning, concentration and thinking, confusion, memory, attention, and psychomotor function), even after controlling for covariates including age, gender, education, and sleep medication use. A study of sleep and neurocognition in a cohort of people living with HIV (75% were neurocognitively impaired based on Frascati criteria) that involved a more comprehensive neurocognitive assessment found that better performance on tasks of attention, frontal/executive function, and psychomotor/motor speed were associated with better polysomnogram sleep parameters, including reduced wake-after sleep, greater sleep

efficiency, greater sleep latency, and greater total sleep time (Gamaldo et al. 2013). Thus, these results demonstrate that confounding factors, such as sleep disturbances, might influence the presentation of HAND.

Sleep disturbance may contribute to neurocognitive decline via impaired metabolism and decreased cerebral perfusion. Sleep deprivation may particularly impair metabolism of the prefrontal cortex, a brain region implicated in executive function (Durmer and Dinges 2005). Insufficient sleep duration has been shown to result in decreased cerebral blood flow in the frontal lobes and in worse performance on the Continuous Performance Test and driving performance (i.e., break reaction time in a harsh-braking test) (Miyata et al. 2010). Evidence from epidemiological and experimental studies indicate that sleep disturbance may impair amyloid beta clearance and increase tau phosphorylation, as well as impair synaptic plasticity via disruption of pathways involving gamma-aminobutyric acid (GABA) and cyclin adenosine monophosphate (cAMP) (Yaffe et al. 2014). In a study specific to persons living with HIV, higher inflammation levels (measured by C-reactive protein and interleukin-6) were observed among those with poor sleep characteristics (i.e., later sleep onset and lower total sleep time) and low engagement in moderate-vigorous physical activity (Wirth et al. 2015). These results indicate that disturbances in sleep and low levels of physical activity are associated with inflammation, which is implicated in the pathogenesis of HAND. Thus, improving sleep indices and maintaining regular participation in physical activity may help reduce the risk of inflammation, which, in turn, may reduce the incidence of HAND.

4.3 Behavioral Strategies to Improve Sleep Quality

Cognitive behavioral therapy for insomnia is the first-line treatment for insomnia based on guidelines of the American College of Physicians for management of chronic insomnia (Qaseem et al. 2016). Cognitive behavioral therapy for insomnia is a multimodal cognitive behavioral therapy that can be delivered in individual or group therapy, telephone- or web-based modules, or self-help books, and includes the behavioral strategies of sleep restriction and stimulus control. Although cognitive behavioral therapy for insomnia is associated with robust, long-term improvements in sleep parameters, cognitive behavioral therapy for insomnia has small-to-moderate effects on subjective measures of neurocognitive functioning, and there is insufficient data to determine the effect of cognitive behavioral therapy for insomnia on objective measures of neurocognition in the general population (Herbert et al. 2018).

SystemCHANGE-HIV – a 10-week intervention grounded in a socioecological model that covers different topics of HIV management, including sleep hygiene and behavioral modification strategies – was tested to determine its effectiveness at improving sleep outcomes, including sleep duration, sleep fragmentation index, sleep efficiency, and self-reported sleep quality (Webel et al. 2013). SystemCHANGE-HIV had high levels of engagement (e.g., participants attended 71% of all intervention sessions on average). Although nonsignificant, the pilot

study provided preliminary data indicating that SystemCHANGE-HIV leads to improvement in sleep efficiency and sleep fragmentation. A major limitation of the SystemCHANGE-HIV study in regard to its applicability for the treatment of HAND was the absence of neurocognitive assessments. Given the prevalence and association between sleep disturbances and neurocognitive impairment in the context of HIV, a future research direction is to determine whether strategies for improving sleep (e.g., cognitive behavioral therapy for insomnia, SystemCHANGE-HIV) may lead to improvement in subjective and objective measures of neurocognitive functioning among people living with HIV.

5 Promotion of Antiretroviral Therapy Adherence as a Behavioral Intervention for HAND

5.1 Antiretroviral Therapy Adherence and Neurocognitive Performance

Early detection and initiation of antiretroviral therapy is a strategy for preventing significant immune compromise and protecting against neurocognitive decline (D'Antoni et al. 2018). In addition to initiating antiretroviral therapy during acute HIV infection, antiretroviral therapy adherence is crucial for the prevention of HAND (Martin et al. 1999; Suarez et al. 2001). Adherence is generally defined as the extent to which patients take medications as prescribed and is typically reported as the percentage of prescribed doses taken over a specified period, which may include consideration of specific dose timing (Osterberg and Blaschke 2005). Achieving consistently higher levels of antiretroviral therapy adherence in order to derive therapeutic benefit (e.g., sustained viral suppression and immune reconstitution) is a challenge for many people living with HIV (Kirtane et al. 2016). Generally, adherence rates are lower for chronic compared to acute conditions, and even under the controlled settings of clinical trials, average adherence rates range between 43 and 67% (Osterberg and Blaschke 2005).

Higher levels of antiretroviral therapy adherence are predictive of improvements in neurocognitive performance in the domains of information processing speed, attention, executive function, and motor function (Ettenhofer et al. 2010). Among patients who initiate antiretroviral therapy or change to a more effective regimen, improvements in neurocognitive function have been observed after 6 months for patients who achieved successful viral suppression (Parsons et al. 2006). Although antiretroviral therapy may improve neurocognitive function, a major complication is that people living with HIV with neurocognitive impairment, including deficits in prospective memory, are particularly at risk for antiretroviral therapy non-adherence (Hinkin et al. 2002; Woods et al. 2009a). Thus, behavioral strategies to promote antiretroviral therapy adherence among people living with HIV with neurocognitive deficits are particularly pertinent for the treatment of HAND.

5.2 *Strategies to Promote Antiretroviral Therapy Adherence*

Strategies to promote optimal adherence among people living with HIV include simpler dosing demands (e.g., fewer pills and once-a-day single tablet regimens) (Mohd Salleh et al. 2018). Given the ubiquity of phones, text message interventions have also been developed to prompt people living with HIV to adhere to their antiretroviral therapy regimen (Horvath et al. 2012). Such interventions are particularly effective in promoting antiretroviral therapy adherence in low-resource settings and may be helpful for promoting better antiretroviral therapy dose timing in patients with severe mental illness, such as bipolar disorder (Moore et al. 2015).

Adherence to antiretroviral therapy is influenced by psychosocial stressors, such as the experience of negative life events and depression, as well as individual characteristics like behavioral impulsivity (Salmoirago-Blotcher et al. 2017). Mindfulness training, which may allay distress and lessen impulsivity, is currently being investigated to determine its potential to help persons living with HIV adhere to their antiretroviral therapy regimen (Salmoirago-Blotcher et al. 2017). Given that depression has consistently shown a robust association with poor antiretroviral therapy adherence, several interventions grounded in cognitive behavioral therapy have been developed and evaluated for the treatment of depression and/or to promote medication adherence (Balfour et al. 2006; Safren et al. 2009; Simoni et al. 2007; Olatunji et al. 2006). A small pilot study investigated the effect of one-session behavioral activation treatment for depression designed specifically for people living with HIV (Tull et al. 2018). Although nonsignificant, persons living with HIV who received the one-session of behavioral activation treatment showed improvements in medication adherence of medium effect (Tull et al. 2018). Additional findings indicate that low levels of physical activity were predictive of poor antiretroviral therapy adherence and higher viral load, but importantly, this relationship was mediated by depression symptoms (Blashill et al. 2013). These results highlight the need to develop cost-effective interventions to simultaneously address symptoms of depression and antiretroviral therapy adherence among people living with HIV in order to prevent or treat HAND, and suggest that physical activity may have multifaceted benefits for people living with HIV.

6 Clinical Implications

In this section, intervention approaches to increase engagement in health behaviors (i.e., physical activity, healthy eating, sleep, and antiretroviral therapy adherence) are reviewed. Such approaches have the potential to improve neurocognitive outcomes among people living with HIV by increasing health behaviors that may protect against the development of non-communicable conditions (Lim et al. 2012) that adversely impact neurocognitive outcomes.

Health care providers have a prominent role in integrating health behavior promotion into routine HIV care (Webel et al. 2017). Health care providers can leverage their relationships with patients living with HIV to emphasize a holistic concept of well-being that includes engagement in health behaviors (Webel et al. 2017). Despite evidence supporting the effectiveness of health behavior promotion in primary health care, implementation has been slow (Brotons et al. 2012). One institution-level approach to target health behaviors more effectively and consistently in routine care is to integrate primary care and behavioral health in a single clinic; however, integration requires substantial reengineering of practice (Cifuentes et al. 2015). Another viable solution for targeting health behaviors in routine clinical care has been to train health care providers in Motivational Interviewing, a client-centered, evidence-based behavior change consultation style (Rollnick et al. 2008). Adoption of a Motivational Interviewing approach may be particularly helpful for clinicians providing HIV care, with some evidence indicating the effectiveness of Motivational Interviewing for improving antiretroviral therapy adherence (Golin et al. 2006; Beach et al. 2015).

Beyond the patient–provider relationship, engagement in a health behavior is influenced by a complex system of determinants, including intrapersonal (e.g., cognitive and emotional factors), interpersonal/community, institutional (e.g., access to specialty medical care), environmental (e.g., neighborhood characteristics and employment conditions), cultural, and policy/legislation factors (Dahlgren and Whitehead 2006). For persons living with HIV, readiness to engage in a health behavior is a dynamic and fluctuating construct, which may also be influenced by the episodic nature of HIV and multimorbidity (Simonik et al. 2016). Identification of contextual factors that have a strong relationship with the presence (or absence) of a health behavior is necessary to identify potential modifiable targets of behavioral health interventions (Michie et al. 2011; Wight et al. 2016). Although the field of behavior change research in persons living with HIV acknowledges that interventions need to target more than just factors at the intrapersonal level (Albarracín et al. 2010), there is only a small literature on multi-level models of intervention (Kaufman et al. 2014). Thus, behavior change efforts may be guided by multiple existing theories that, in combination, target various levels of influence (Kaufman et al. 2014).

Ideally, development of behavioral health interventions involves an interdisciplinary group representing relevant expertise (e.g., clinical healthcare, psychology, epidemiology, and policy) and key stakeholders (e.g., patients, caregivers, healthcare professionals, policymakers, and funders) to ensure interventions are evidence-based and acceptable to (1) those for whom the intervention is developed and (2) those involved in the adoption and implementation of the intervention (Eldredge et al. 2016; Witteman et al. 2017; Araújo-Soares et al. 2018). To optimize the effectiveness of a behavioral health intervention, researchers and/or interventionists must consider the appropriateness of various behavior change techniques, mode(s) of intervention delivery, provider(s), location, timing, and dosing of an intervention (Araújo-Soares et al. 2018).

Given the complexity of behavior change interventions, it is important to identify (1) effective behavior change techniques and (2) the processes through which behavior change occurs (i.e., the mechanisms of action) (Connell et al. 2018). For researchers and interventionists, various protocols and taxonomies have been developed to aid in behavior change intervention development. For example, Intervention Mapping describes an iterative process for developing theory- and evidence-based health promotion programs (Kok et al. 2004). In addition, an extensive taxonomy of consensually agreed, distinct behavior change techniques has been developed by a large international network of behavior change experts, which is intended to be used in combination with Intervention Mapping (Michie et al. 2013). Additional research investigating the effectiveness of specific behavior change techniques for promoting the adoption and maintenance of healthy lifestyle factors is needed. Such intervention efforts may reduce inflammation and oxidative stress, which, in turn, may reduce the incidence and/or severity of HAND.

7 Conclusion

Currently, there is no gold-standard treatment for the prevention or treatment of HAND. Although there is an absence of evidence-based neurocognitive interventions for people living with HIV, the literature reviewed in this chapter suggest potential targets for intervention. Modifiable lifestyle factors, such as physical activity, diet, sleep, and antiretroviral therapy adherence, may benefit neurocognitive function among people living with HIV. These healthy lifestyle factors reduce inflammation and oxidative stress, which, in turn, may reduce the incidence of HAND.

References

- Albarracín D, Rothman AJ, di Clemente R, del Rio C (2010) Wanted: a theoretical roadmap to research and practice across individual, interpersonal, and structural levels of analysis. *AIDS Behav* 14(Suppl 2):185–188. <https://doi.org/10.1007/s10461-010-9805-x>
- Ancuta P, Kamat A, Kunstman KJ, Kim EY, Autissier P, Wurcel A et al (2008) Microbial translocation is associated with increased monocyte activation and dementia in AIDS patients. *PLoS One* 3(6):e2516. <https://doi.org/10.1371/journal.pone.0002516>
- Antinori A, Arendt G, Becker JT, Brew BJ, Byrd DA, Cherner M et al (2007) Updated research nosology for HIV-associated neurocognitive disorders. *Neurology* 69(18):1789–1799. <https://doi.org/10.1212/01.WNL.0000287431.88658.8b>
- Araújo-Soares V, Hankonen N, Presseau J, Rodrigues A, Sniehotta F (2018) Developing behavior change interventions for self-management in chronic illness. *Eur Psychol*. <https://doi.org/10.1027/1016-9040/a000330>

- Balfour L, Kowal J, Silverman A, Tasca GA, Angel JB, Macpherson PA et al (2006) A randomized controlled psycho-education intervention trial: improving psychological readiness for successful HIV medication adherence and reducing depression before initiating HAART. *AIDS Care* 18 (7):830–838. <https://doi.org/10.1080/09540120500466820>
- Barak Y, Aizenberg D (2010) Is dementia preventable? Focus on Alzheimer's disease. *Expert Rev Neurother* 10(11):1689–1698. <https://doi.org/10.1586/ern.10.159>
- Beach MC, Roter DL, Saha S, Korthuis PT, Eggly S, Cohn J et al (2015) Impact of a brief patient and provider intervention to improve the quality of communication about medication adherence among HIV patients. *Patient Educ Couns* 98(9):1078–1083. <https://doi.org/10.1016/j.pec.2015.05.011>
- Best JR, Chiu BK, Liang Hsu C, Nagamatsu LS, Liu-Ambrose T (2015) Long-term effects of resistance exercise training on cognition and brain volume in older women: results from a randomized controlled trial. *J Int Neuropsychol Soc* 21(10):745–756. <https://doi.org/10.1017/S1355617715000673>
- Blackwell T, Yaffe K, Laffan A, Ancoli-Israel S, Redline S, Ensrud KE et al (2014) Associations of objectively and subjectively measured sleep quality with subsequent cognitive decline in older community-dwelling men: the MrOS sleep study. *Sleep* 37(4):655–663. <https://doi.org/10.5665/sleep.3562>
- Blashill AJ, Mayer KH, Crane H, Magidson JF, Grasso C, Mathews WC et al (2013) Physical activity and health outcomes among HIV-infected men who have sex with men: a longitudinal mediational analysis. *Ann Behav Med* 46(2):149–156. <https://doi.org/10.1007/s12160-013-9489-3>
- Botros D, Somarriba G, Neri D, Miller TL (2012) Interventions to address chronic disease and HIV: strategies to promote exercise and nutrition among HIV-infected individuals. *Curr HIV/AIDS Rep* 9(4):351–363. <https://doi.org/10.1007/s11904-012-0135-7>
- Brottons C, Bulc M, Sammut MR, Sheehan M, Manuel da Silva Martins C, Bjorkelund C et al (2012) Attitudes toward preventive services and lifestyle: the views of primary care patients in Europe. The EUROPREVIEW patient study. *Fam Pract* 29(Suppl 1):i168–i176. <https://doi.org/10.1093/fampra/cm102>
- Byun E, Gay CL, Lee KA (2016) Sleep, fatigue, and problems with cognitive function in adults living with HIV. *J Assoc Nurses AIDS Care* 27(1):5–16. <https://doi.org/10.1016/j.jana.2015.10.002>
- Capili B, Anastasi JK (2008) Body mass index and nutritional intake in patients with HIV and chronic diarrhea: a secondary analysis. *J Am Acad Nurse Pract* 20(9):463–470. <https://doi.org/10.1111/j.1745-7599.2008.00349.x>
- Carr A, Samaras K, Burton S, Law M, Freund J, Chisholm DJ et al (1998) A syndrome of peripheral lipodystrophy, hyperlipidaemia and insulin resistance in patients receiving HIV protease inhibitors. *AIDS* 12(7):F51–F58
- Chen YC, Lin CY, Strong C, Li CY, Wang JS, Ko WC et al (2017) Sleep disturbances at the time of a new diagnosis: a comparative study of human immunodeficiency virus patients, cancer patients, and general population controls. *Sleep Med* 36:38–43. <https://doi.org/10.1016/j.sleep.2017.04.005>
- Cifuentes M, Davis M, Fernald D, Gunn R, Dickinson P, Cohen DJ (2015) Electronic health record challenges, workarounds, and solutions observed in practices integrating behavioral health and primary care. *J Am Board Fam Med* 28(Suppl 1):S63–S72. <https://doi.org/10.3122/jabfm.2015.S1.150133>
- Connell LE, Carey RN, de Bruin M, Rothman AJ, Johnston M, Kelly MP et al (2018) Links between behavior change techniques and mechanisms of action: an expert consensus study. *Ann Behav Med*. <https://doi.org/10.1093/abm/kay082>
- Cysique LA, Moffat K, Moore DM, Lane TA, Davies NW, Carr A et al (2013) HIV, vascular and aging injuries in the brain of clinically stable HIV-infected adults: a (1)H MRS study. *PLoS One* 8(4):e61738. <https://doi.org/10.1371/journal.pone.0061738>

- D'Antoni ML, Byron MM, Chan P, Sailasuta N, Sacdalan C, Sithinamsuwan P et al (2018) Normalization of soluble CD163 after institution of antiretroviral therapy during acute HIV infection tracks with fewer neurological abnormalities. *J Infect Dis* 218(9):1453–1463. <https://doi.org/10.1093/infdis/jiy337>
- d'Ettore G, Ceccarelli G, Giustini N, Mastroianni CM, Silvestri G, Vullo V (2014) Taming HIV-related inflammation with physical activity: a matter of timing. *AIDS Res Hum Retrovir* 30(10):936–944. <https://doi.org/10.1089/AID.2014.0069>
- Dahlgren G, Whitehead M (2006) European strategies for tackling social inequities in health: levelling up Part 2. World Health Organization, Copenhagen
- Dang AK, Nguyen LH, Nguyen AQ, Tran BX, Tran TT, Latkin CA et al (2018) Physical activity among HIV-positive patients receiving antiretroviral therapy in Hanoi and Nam Dinh, Vietnam: a cross-sectional study. *BMJ Open* 8(5):e020688. <https://doi.org/10.1136/bmjopen-2017-020688>
- Davis CL, Tomporowski PD, Boyle CA, Waller JL, Miller PH, Naglieri JA et al (2007) Effects of aerobic exercise on overweight children's cognitive functioning: a randomized controlled trial. *Res Q Exerc Sport* 78(5):510–519. <https://doi.org/10.1080/02701367.2007.10599450>
- Devore EE, Kang JH, Breteler MM, Grodstein F (2012) Dietary intakes of berries and flavonoids in relation to cognitive decline. *Ann Neurol* 72(1):135–143. <https://doi.org/10.1002/ana.23594>
- Dube MP, Stein JH, Aberg JA, Fichtenbaum CJ, Gerber JG, Tashima KT et al (2003) Guidelines for the evaluation and management of dyslipidemia in human immunodeficiency virus (HIV)-infected adults receiving antiretroviral therapy: recommendations of the HIV Medical Association of the Infectious Disease Society of America and the Adult AIDS Clinical Trials Group. *Clin Infect Dis* 37(5):613–627. <https://doi.org/10.1086/378131>
- Dufour CA, Marquine MJ, Fazeli PL, Henry BL, Ellis RJ, Grant I et al (2013) Physical exercise is associated with less neurocognitive impairment among HIV-infected adults. *J Neurovirol* 19(5):410–417. <https://doi.org/10.1007/s13365-013-0184-8>
- Dufour CA, Marquine MJ, Fazeli PL, Umlauf A, Henry BL, Zlatar Z et al (2018) A longitudinal analysis of the impact of physical activity on neurocognitive functioning among HIV-infected adults. *AIDS Behav* 22(5):1562–1572. <https://doi.org/10.1007/s10461-016-1643-z>
- Duran AC, Almeida LB, Segurado AA, Jaime PC (2008) Diet quality of persons living with HIV/AIDS on highly active antiretroviral therapy. *J Hum Nutr Diet* 21(4):346–350. <https://doi.org/10.1111/j.1365-277X.2008.00886.x>
- Durmer JS, Dinges DF (2005) Neurocognitive consequences of sleep deprivation. *Semin Neurol* 25(1):117–129. <https://doi.org/10.1055/s-2005-867080>
- Eldredge LKB, Markham CM, Ruiter RAC, Kok G, Fernandez ME, Parcel GS (2016) Planning health promotion programs: an intervention mapping approach. Wiley, Hoboken
- Engeroff T, Ingmann T, Banzer W (2018) Physical activity throughout the adult life span and domain-specific cognitive function in old age: a systematic review of cross-sectional and longitudinal data. *Sports Med* 48(6):1405–1436. <https://doi.org/10.1007/s40279-018-0920-6>
- Estruch R, Ros E, Salas-Salvado J, Covas MI, Corella D, Aros F et al (2013) Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 368(14):1279–1290. <https://doi.org/10.1056/NEJMoa1200303>
- Ettenhofer ML, Foley J, Castellon SA, Hinkin CH (2010) Reciprocal prediction of medication adherence and neurocognition in HIV/AIDS. *Neurology* 74(15):1217–1222. <https://doi.org/10.1212/WNL.0b013e3181d8c1ca>
- Farina N, Rusted J, Tabet N (2014) The effect of exercise interventions on cognitive outcome in Alzheimer's disease: a systematic review. *Int Psychogeriatr* 26(1):9–18. <https://doi.org/10.1017/S1041610213001385>
- Faubel R, Lopez-Garcia E, Guallar-Castillon P, Graciani A, Banegas JR, Rodriguez-Artalejo F (2009) Usual sleep duration and cognitive function in older adults in Spain. *J Sleep Res* 18(4):427–435. <https://doi.org/10.1111/j.1365-2869.2009.00759.x>

- Fazeli PL, Marquine MJ, Dufour C, Henry BL, Montoya J, Gouaux B et al (2015) Physical activity is associated with better neurocognitive and everyday functioning among older adults with HIV disease. *AIDS Behav* 19(8):1470–1477. <https://doi.org/10.1007/s10461-015-1024-z>
- Firth J, Cotter J, Carney R, Yung AR (2017) The pro-cognitive mechanisms of physical exercise in people with schizophrenia. *Br J Pharmacol* 174(19):3161–3172. <https://doi.org/10.1111/bph.13772>
- Fito M, Estruch R, Salas-Salvado J, Martinez-Gonzalez MA, Aros F, Vila J et al (2014) Effect of the Mediterranean diet on heart failure biomarkers: a randomized sample from the PREDIMED trial. *Eur J Heart Fail* 16(5):543–550. <https://doi.org/10.1002/ehf.61>
- Fjeldsoe BS, Marshall AL, Miller YD (2009) Behavior change interventions delivered by mobile telephone short-message service. *Am J Prev Med* 36(2):165–173. <https://doi.org/10.1016/j.amepre.2008.09.040>
- Gamaldo CE, Gamaldo A, Creighton J, Salas RE, Selnes OA, David PM et al (2013) Evaluating sleep and cognition in HIV. *J Acquir Immune Defic Syndr* 63(5):609–616. <https://doi.org/10.1097/QAI.0b013e31829d63ab>
- Gannon P, Khan MZ, Kolson DL (2011) Current understanding of HIV-associated neurocognitive disorders pathogenesis. *Curr Opin Neurol* 24(3):275–283. <https://doi.org/10.1097/WCO.0b013e3182834695fb>
- Gill SJ, Friedenreich CM, Sajobi TT, Longman RS, Drogos LL, Davenport MH et al (2015) Association between lifetime physical activity and cognitive functioning in middle-aged and older community dwelling adults: results from the brain in motion study. *J Int Neuropsychol Soc* 21(10):816–830. <https://doi.org/10.1017/S1355617715000880>
- Golin CE, Earp J, Tien HC, Stewart P, Porter C, Howie L (2006) A 2-arm, randomized, controlled trial of a motivational interviewing-based intervention to improve adherence to antiretroviral therapy (ART) among patients failing or initiating ART. *J Acquir Immune Defic Syndr* 42(1):42–51. <https://doi.org/10.1097/01.qai.0000219771.97303.0a>
- Grobler L, Siegfried N, Visser ME, Mahlangu SS, Volmink J (2013) Nutritional interventions for reducing morbidity and mortality in people with HIV. *Cochrane Database Syst Rev* 2:CD004536. <https://doi.org/10.1002/14651858.CD004536.pub3>
- Guiney H, Machado L (2013) Benefits of regular aerobic exercise for executive functioning in healthy populations. *Psychon Bull Rev* 20(1):73–86. <https://doi.org/10.3758/s13423-012-0345-4>
- Hadigan C, Jeste S, Anderson EJ, Tsay R, Cyr H, Grinspoon S (2001) Modifiable dietary habits and their relation to metabolic abnormalities in men and women with human immunodeficiency virus infection and fat redistribution. *Clin Infect Dis* 33(5):710–717. <https://doi.org/10.1086/322680>
- Hand GA, Lyster GW, Jagers JR, Dudgeon WD (2009) Impact of aerobic and resistance exercise on the health of HIV-infected persons. *Am J Lifestyle Med* 3(6):489–499. <https://doi.org/10.1177/1559827609342198>
- Healy GN, Dunstan DW, Salmon J, Cerin E, Shaw JE, Zimmet PZ et al (2008) Breaks in sedentary time: beneficial associations with metabolic risk. *Diabetes Care* 31(4):661–666
- Heaton RK, Clifford DB, Franklin DR Jr, Woods SP, Ake C, Vaida F et al (2010) HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study. *Neurology* 75(23):2087–2096. <https://doi.org/10.1212/WNL.0b013e318200d727>
- Henry BL, Moore DJ (2016) Preliminary findings describing participant experience with iSTEP, an mHealth intervention to increase physical activity and improve neurocognitive function in people living with HIV. *J Assoc Nurses AIDS Care* 27(4):495–511. <https://doi.org/10.1016/j.jana.2016.01.001>
- Herbert V, Kyle SD, Pratt D (2018) Does cognitive behavioural therapy for insomnia improve cognitive performance? A systematic review and narrative synthesis. *Sleep Med Rev* 39:37–51. <https://doi.org/10.1016/j.smrv.2017.07.001>

- Hinkin CH, Castellon SA, Durvasula RS, Hardy DJ, Lam MN, Mason KI et al (2002) Medication adherence among HIV+ adults: effects of cognitive dysfunction and regimen complexity. *Neurology* 59(12):1944–1950
- Hong S, Banks WA (2015) Role of the immune system in HIV-associated neuroinflammation and neurocognitive implications. *Brain Behav Immun* 45:1–12. <https://doi.org/10.1016/j.bbi.2014.10.008>
- Horvath T, Azman H, Kennedy GE, Rutherford GW (2012) Mobile phone text messaging for promoting adherence to antiretroviral therapy in patients with HIV infection. *Cochrane Database Syst Rev* 3:Cd009756. <https://doi.org/10.1002/14651858.Cd009756>
- Jong E, Oudhoff LA, Epskamp C, Wagener MN, van Duijn M, Fischer S et al (2010) Predictors and treatment strategies of HIV-related fatigue in the combined antiretroviral therapy era. *AIDS* 24(10):1387–1405. <https://doi.org/10.1097/QAD.0b013e328339d004>
- Kapetanovic S, Leister E, Nichols S, Miller T, Tassiopoulos K, Hazra R et al (2010) Relationships between markers of vascular dysfunction and neurodevelopmental outcomes in perinatally HIV-infected youth. *AIDS* 24(10):1481–1491. <https://doi.org/10.1097/QAD.0b013e32833a241b>
- Kapetanovic S, Griner R, Zeldow B, Nichols S, Leister E, Gelbard HA et al (2014) Biomarkers and neurodevelopment in perinatally HIV-infected or exposed youth: a structural equation model analysis. *AIDS* 28(3):355–364. <https://doi.org/10.1097/QAD.0000000000000072>
- Kaufman MR, Cornish F, Zimmerman RS, Johnson BT (2014) Health behavior change models for HIV prevention and AIDS care: practical recommendations for a multi-level approach. *J Acquir Immune Defic Syndr* 66(Suppl 3):S250–S258. <https://doi.org/10.1097/QAI.0000000000000236>
- Kemoun G, Thibaud M, Roumagne N, Carette P, Albinet C, Toussaint L et al (2010) Effects of a physical training programme on cognitive function and walking efficiency in elderly persons with dementia. *Dement Geriatr Cogn Disord* 29(2):109–114. <https://doi.org/10.1159/000272435>
- Kimhy D, Vakhrusheva J, Bartels MN, Armstrong HF, Ballon JS, Khan S et al (2015) The impact of aerobic exercise on brain-derived neurotrophic factor and neurocognition in individuals with schizophrenia: a single-blind, randomized clinical trial. *Schizophr Bull* 41(4):859–868. <https://doi.org/10.1093/schbul/sbv022>
- Kirtane AR, Langer R, Traverso G (2016) Past, present, and future drug delivery systems for antiretrovirals. *J Pharm Sci* 105(12):3471–3482. <https://doi.org/10.1016/j.xphs.2016.09.015>
- Knight A, Bryan J, Murphy K (2017) The Mediterranean diet and age-related cognitive functioning: a systematic review of study findings and neuropsychological assessment methodology. *Nutr Neurosci* 20(8):449–468. <https://doi.org/10.1080/1028415X.2016.1183341>
- Kok G, Schaalma H, Ruiter RA, van Empelen P, Brug J (2004) Intervention mapping: protocol for applying health psychology theory to prevention programmes. *J Health Psychol* 9(1):85–98. <https://doi.org/10.1177/1359105304038379>
- Kruzich LA, Marquis GS, Wilson CM, Stephensen CB (2004) HIV-infected US youth are at high risk of obesity and poor diet quality: a challenge for improving short- and long-term health outcomes. *J Am Diet Assoc* 104(10):1554–1560. <https://doi.org/10.1016/j.jada.2004.07.031>
- Lee KA, Gay C, Portillo CJ, Coggins T, Davis H, Pullinger CR et al (2009) Symptom experience in HIV-infected adults: a function of demographic and clinical characteristics. *J Pain Symptom Manag* 38(6):882–893. <https://doi.org/10.1016/j.jpainsymman.2009.05.013>
- Leyes P, Martinez E, Forga Mde T (2008) Use of diet, nutritional supplements and exercise in HIV-infected patients receiving combination antiretroviral therapies: a systematic review. *Antivir Ther* 13(2):149–159
- Lim J, Dinges DF (2010) A meta-analysis of the impact of short-term sleep deprivation on cognitive variables. *Psychol Bull* 136(3):375–389. <https://doi.org/10.1037/a0018883>
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H et al (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380(9859):2224–2260. [https://doi.org/10.1016/s0140-6736\(12\)61766-8](https://doi.org/10.1016/s0140-6736(12)61766-8)

- Loughrey DG, Lavecchia S, Brennan S, Lawlor BA, Kelly ME (2017) The impact of the Mediterranean diet on the cognitive functioning of healthy older adults: a systematic review and meta-analysis. *Adv Nutr* 8(4):571–586. <https://doi.org/10.3945/an.117.015495>
- Manson JE, Hu FB, Rich-Edwards JW, Colditz GA, Stampfer MJ, Willett WC et al (1999) A prospective study of walking as compared with vigorous exercise in the prevention of coronary heart disease in women. *N Engl J Med* 341(9):650–658
- Marcotte TD, Wolfson T, Rosenthal TJ, Heaton RK, Gonzalez R, Ellis RJ et al (2004) A multimodal assessment of driving performance in HIV infection. *Neurology* 63(8):1417–1422
- Martin EM, Pitrak DL, Novak RM, Pursell KJ, Mullane KM (1999) Reaction times are faster in HIV-seropositive patients on antiretroviral therapy: a preliminary report. *J Clin Exp Neuropsychol* 21(5):730–735. <https://doi.org/10.1076/jcen.21.5.730.867>
- Martinez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvado J, San Julian B et al (2013) Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *J Neurol Neurosurg Psychiatry* 84(12):1318–1325. <https://doi.org/10.1136/jnnp-2012-304792>
- McCutchan JA, Marquie-Beck JA, Fitzsimons CA, Letendre SL, Ellis RJ, Heaton RK et al (2012) Role of obesity, metabolic variables, and diabetes in HIV-associated neurocognitive disorder. *Neurology* 78(7):485–492. <https://doi.org/10.1212/WNL.0b013e3182478d64>
- McDermott A, Zaporozjan L, McNamara P, Doherty CP, Redmond J, Forde C et al (2017) The effects of a 16-week aerobic exercise programme on cognitive function in people living with HIV. *AIDS Care* 29(6):667–674. <https://doi.org/10.1080/09540121.2016.1263723>
- Michie S, van Stralen MM, West R (2011) The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 6:42. <https://doi.org/10.1186/1748-5908-6-42>
- Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W et al (2013) The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med* 46(1):81–95. <https://doi.org/10.1007/s12160-013-9486-6>
- Miller KD, Jones E, Yanovski JA, Shankar R, Feuerstein I, Falloon J (1998) Visceral abdominal-fat accumulation associated with use of indinavir. *Lancet* 351(9106):871–875. [https://doi.org/10.1016/S0140-6736\(97\)11518-5](https://doi.org/10.1016/S0140-6736(97)11518-5)
- Miller TL, Jacobson DL, Somarriba G, Neri D, Kurtz-Vraney J, Graham P et al (2017) A multicenter study of diet quality on birth weight and gestational age in infants of HIV-infected women. *Matern Child Nutr* 13(4):e12378. <https://doi.org/10.1111/mcn.12378>
- Millikin CP, Rourke SB, Halman MH, Power C (2003) Fatigue in HIV/AIDS is associated with depression and subjective neurocognitive complaints but not neuropsychological functioning. *J Clin Exp Neuropsychol* 25(2):201–215. <https://doi.org/10.1076/jcen.25.2.201.13644>
- Mitjavila MT, Fandos M, Salas-Salvado J, Covas MI, Borrego S, Estruch R et al (2013) The Mediterranean diet improves the systemic lipid and DNA oxidative damage in metabolic syndrome individuals. A randomized, controlled, trial. *Clin Nutr* 32(2):172–178. <https://doi.org/10.1016/j.clnu.2012.08.002>
- Miyata S, Noda A, Ozaki N, Hara Y, Minoshima M, Iwamoto K et al (2010) Insufficient sleep impairs driving performance and cognitive function. *Neurosci Lett* 469(2):229–233. <https://doi.org/10.1016/j.neulet.2009.12.001>
- Mohd Salleh NA, Richardson L, Kerr T, Shoveller J, Montaner J, Kamarulzaman A et al (2018) A longitudinal analysis of daily pill burden and likelihood of optimal adherence to antiretroviral therapy among people living with HIV who use drugs. *J Addict Med* 12(4):308–314. <https://doi.org/10.1097/ADM.0000000000000403>
- Monroe AK, Zhang L, Jacobson LP, Plankey MW, Brown TT, Miller EN et al (2017) The association between physical activity and cognition in men with and without HIV infection. *HIV Med* 18(8):555–563. <https://doi.org/10.1111/hiv.12490>
- Montoya JL, Wing D, Knight A, Moore DJ, Henry BL (2015) Development of an mHealth intervention (iSTEP) to promote physical activity among people living with HIV. *J Int Assoc Provid AIDS Care* 14(6):471–475. <https://doi.org/10.1177/2325957415601505>

- Moore DJ, Poquette A, Casaletto KB, Gouaux B, Montoya JL, Posada C et al (2015) Individualized texting for adherence building (iTAB): improving antiretroviral dose timing among HIV-infected persons with co-occurring bipolar disorder. *AIDS Behav* 19(3):459–471. <https://doi.org/10.1007/s10461-014-0971-0>
- Munk PS, Breland UM, Aukrust P, Ueland T, Kvaloy JT, Larsen AI (2011) High intensity interval training reduces systemic inflammation in post-PCI patients. *Eur J Cardiovasc Prev Rehabil* 18(6):850–857. <https://doi.org/10.1177/1741826710397600>
- Mura G, Vellante M, Nardi AE, Machado S, Carta MG (2015) Effects of school-based physical activity interventions on cognition and academic achievement: a systematic review. *CNS Neurol Disord Drug Targets* 14(9):1194–1208
- Nurk E, Refsum H, Drevon CA, Tell GS, Nygaard HA, Engedal K et al (2009) Intake of flavonoid-rich wine, tea, and chocolate by elderly men and women is associated with better cognitive test performance. *J Nutr* 139(1):120–127. <https://doi.org/10.3945/jn.108.095182>
- O'Brien K, Nixon S, Tynan AM, Glazier R (2010) Aerobic exercise interventions for adults living with HIV/AIDS. *Cochrane Database Syst Rev* 8:CD001796. <https://doi.org/10.1002/14651858.CD001796.pub3>
- Olatunji BO, Mimiaga MJ, O'Cleirigh C, Safren SA (2006) Review of treatment studies of depression in HIV. *Top HIV Med* 14(3):112–124
- Ortega M, Baker LM, Vaida F, Paul R, Basco B, Ances BM (2015) Physical activity affects brain integrity in HIV+ individuals. *J Int Neuropsychol Soc* 21(10):880–889. <https://doi.org/10.1017/S1355617715000879>
- Osterberg L, Blaschke T (2005) Adherence to medication. *N Engl J Med* 353(5):487–497. <https://doi.org/10.1056/NEJMr050100>
- Parsons TD, Braaten AJ, Hall CD, Robertson KR (2006) Better quality of life with neuropsychological improvement on HAART. *Health Qual Life Outcomes* 4:11. <https://doi.org/10.1186/1477-7525-4-11>
- Poulouse SM, Miller MG, Shukitt-Hale B (2014) Role of walnuts in maintaining brain health with age. *J Nutr* 144(4 Suppl):561S–566S. <https://doi.org/10.3945/jn.113.184838>
- Qaseem A, Kansagara D, Forciea MA, Cooke M, Denberg TD (2016) Management of chronic insomnia disorder in adults: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 165(2):125–133. <https://doi.org/10.7326/M15-2175>
- Quigley A, O'Brien K, Parker R, MacKay-Lyons M (2018) Exercise and cognitive function in people living with HIV: a scoping review. *Disabil Rehabil*:1–12. <https://doi.org/10.1080/09638288.2018.1432079>
- Ramos JS, Dalleck LC, Tjonna AE, Beetham KS, Coombes JS (2015) The impact of high-intensity interval training versus moderate-intensity continuous training on vascular function: a systematic review and meta-analysis. *Sports Med* 45(5):679–692. <https://doi.org/10.1007/s40279-015-0321-z>
- Rehm KE, Konkole-Parker D (2016) Physical activity levels and perceived benefits and barriers to physical activity in HIV-infected women living in the deep south of the United States. *AIDS Care* 28(9):1205–1210. <https://doi.org/10.1080/09540121.2016.1164802>
- Ren J, Zhao M, Liu B, Wu Q, Hao Y, Jiao M et al (2018) Factors associated with sleep quality in HIV. *J Assoc Nurses AIDS Care* 29(6):924–931. <https://doi.org/10.1016/j.jana.2018.04.006>
- Rollnick S, Miller WR, Butler CC (2008) Motivational interviewing in health care. Guilford Press, New York
- Rosenberg DE, Kerr J, Sallis JF, Norman GJ, Calfas K, Patrick K (2012) Promoting walking among older adults living in retirement communities. *J Aging Phys Act* 20(3):379–394. <https://doi.org/10.1123/japa.20.3.379>
- Rubinstein ML, Selwyn PA (1998) High prevalence of insomnia in an outpatient population with HIV infection. *J Acquir Immune Defic Syndr Hum Retrovirol* 19(3):260–265
- Safren SA, O'Cleirigh C, Tan JY, Raminani SR, Reilly LC, Otto MW et al (2009) A randomized controlled trial of cognitive behavioral therapy for adherence and depression (CBT-AD) in HIV-infected individuals. *Health Psychol* 28(1):1–10. <https://doi.org/10.1037/a0012715>

- Salmoirago-Blotcher E, Rich C, Rosen RK, Dunsiger S, Rana A, Carey MP (2017) Phone-delivered mindfulness training to promote medication adherence and reduce sexual risk behavior among persons living with HIV: design and methods. *Contemp Clin Trials* 53:162–170. <https://doi.org/10.1016/j.cct.2016.12.016>
- Sattler FR, He J, Letendre S, Wilson C, Sanders C, Heaton R et al (2015) Abdominal obesity contributes to neurocognitive impairment in HIV-infected patients with increased inflammation and immune activation. *J Acquir Immune Defic Syndr* 68(3):281–288. <https://doi.org/10.1097/QAI.0000000000000458>
- Saylor D, Dickens AM, Sacktor N, Haughey N, Slusher B, Pletnikov M et al (2016) HIV-associated neurocognitive disorder – pathogenesis and prospects for treatment. *Nat Rev Neurol* 12(5):309. <https://doi.org/10.1038/nrneurol.2016.53>
- Schaefer EJ, Bongard V, Beiser AS, Lamon-Fava S, Robins SJ, Au R et al (2006) Plasma phosphatidylcholine docosahexaenoic acid content and risk of dementia and Alzheimer disease: the Framingham Heart Study. *Arch Neurol* 63(11):1545–1550. <https://doi.org/10.1001/archneur.63.11.1545>
- Schroder H, Salas-Salvado J, Martinez-Gonzalez MA, Fito M, Corella D, Estruch R et al (2014) Baseline adherence to the Mediterranean diet and major cardiovascular events: Prevencion con Dieta Mediterranea trial. *JAMA Intern Med* 174(10):1690–1692. <https://doi.org/10.1001/jamainternmed.2014.3463>
- Schuelter-Trevisol F, Wolff FH, Alencastro PR, Grigoletti S, Ikeda ML, Brandao AB et al (2012) Physical activity: do patients infected with HIV practice? How much? A systematic review. *Curr HIV Res* 10(6):487–497
- Shah M, Tierney K, Adams-Huet B, Boonyavarakul A, Jacob K, Quittner C et al (2005) The role of diet, exercise and smoking in dyslipidaemia in HIV-infected patients with lipodystrophy. *HIV Med* 6(4):291–298. <https://doi.org/10.1111/j.1468-1293.2005.00309.x>
- Simoni JM, Pantalone DW, Plummer MD, Huang B (2007) A randomized controlled trial of a peer support intervention targeting antiretroviral medication adherence and depressive symptomatology in HIV-positive men and women. *Health Psychol* 26(4):488–495. <https://doi.org/10.1037/0278-6133.26.4.488>
- Simonik A, Vader K, Ellis D, Kesbrian D, Leung P, Jachyra P et al (2016) Are you ready? Exploring readiness to engage in exercise among people living with HIV and multimorbidity in Toronto, Canada: a qualitative study. *BMJ Open* 6(3):e010029. <https://doi.org/10.1136/bmjopen-2015-010029>
- Smith PJ, Blumenthal JA, Hoffman BM, Cooper H, Strauman TA, Welsh-Bohmer K et al (2010) Aerobic exercise and neurocognitive performance: a meta-analytic review of randomized controlled trials. *Psychosom Med* 72(3):239–252. <https://doi.org/10.1097/PSY.0b013e3181d14633>
- Song D, Yu DSF, Li PWC, Lei Y (2018) The effectiveness of physical exercise on cognitive and psychological outcomes in individuals with mild cognitive impairment: a systematic review and meta-analysis. *Int J Nurs Stud* 79:155–164. <https://doi.org/10.1016/j.ijnurstu.2018.01.002>
- Stimpson NJ, Davison G, Javadi AH (2018) Joggin’ the noggin: towards a physiological understanding of exercise-induced cognitive benefits. *Neurosci Biobehav Rev* 88:177–186. <https://doi.org/10.1016/j.neubiorev.2018.03.018>
- Stradling C, Chen YF, Russell T, Connock M, Thomas GN, Taheri S (2012) The effects of dietary intervention on HIV dyslipidaemia: a systematic review and meta-analysis. *PLoS One* 7(6):e38121. <https://doi.org/10.1371/journal.pone.0038121>
- Suarez S, Baril L, Stankoff B, Khellaf M, Dubois B, Lubetzki C et al (2001) Outcome of patients with HIV-1-related cognitive impairment on highly active antiretroviral therapy. *AIDS* 15(2):195–200
- Sugiura H, Sugiura H, Kajima K, Mirbod SM, Iwata H, Matsuoka T (2002) Effects of long-term moderate exercise and increase in number of daily steps on serum lipids in women: randomised controlled trial [ISRCTN21921919]. *BMC Womens Health* 2(1):3

- Taibi DM, Price C, Voss J (2013) A pilot study of sleep quality and rest-activity patterns in persons living with HIV. *J Assoc Nurses AIDS Care* 24(5):411–421. <https://doi.org/10.1016/j.jana.2012.08.001>
- Tavazzi E, Morrison D, Sullivan P, Morgello S, Fischer T (2014) Brain inflammation is a common feature of HIV-infected patients without HIV encephalitis or productive brain infection. *Curr HIV Res* 12(2):97–110
- Thames AD, Kim MS, Becker BW, Foley JM, Hines LJ, Singer EJ et al (2011) Medication and finance management among HIV-infected adults: the impact of age and cognition. *J Clin Exp Neuropsychol* 33(2):200–209. <https://doi.org/10.1080/13803395.2010.499357>
- Tjonna AE, Stolen TO, Bye A, Volden M, Slordahl SA, Odegard R et al (2009) Aerobic interval training reduces cardiovascular risk factors more than a multitreatment approach in overweight adolescents. *Clin Sci (Lond)* 116(4):317–326. <https://doi.org/10.1042/CS20080249>
- Tsiodras S, Pouliou KA, Yannakoulia M, Chimienti SN, Wadhwa S, Karchmer AW et al (2009) Adherence to Mediterranean diet is favorably associated with metabolic parameters in HIV-positive patients with the highly active antiretroviral therapy-induced metabolic syndrome and lipodystrophy. *Metabolism* 58(6):854–859. <https://doi.org/10.1016/j.metabol.2009.02.012>
- Tull MT, Berghoff CR, Bardeen JR, Schoenleber M, Konkle-Parker DJ (2018) An initial open trial of a brief behavioral activation treatment for depression and medication adherence in HIV-infected patients. *Behav Modif* 42(2):196–209. <https://doi.org/10.1177/0145445517723901>
- Valcour VG, Shikuma CM, Shiramizu BT, Williams AE, Watters MR, Poff PW et al (2005) Diabetes, insulin resistance, and dementia among HIV-1-infected patients. *J Acquir Immune Defic Syndr* 38(1):31–36
- Vancampfort D, Mugisha J, De Hert M, Probst M, Firth J, Gorczynski P et al (2018a) Global physical activity levels among people living with HIV: a systematic review and meta-analysis. *Disabil Rehabil* 40(4):388–397. <https://doi.org/10.1080/09638288.2016.1260645>
- Vancampfort D, Stubbs B, Mugisha J, Firth J, Schuch FB, Koyanagi A (2018b) Correlates of sedentary behavior in 2,375 people with depression from 6 low- and middle-income countries. *J Affect Disord* 234:97–104. <https://doi.org/10.1016/j.jad.2018.02.088>
- Vance DE, Wadley VG, Crowe MG, Raper JL, Ball KK (2011) Cognitive and everyday functioning in older and younger adults with and without HIV. *Clin Gerontol* 34(5):413–426. <https://doi.org/10.1080/07317115.2011.588545>
- Vanderbeken I, Kerckhofs E (2017) A systematic review of the effect of physical exercise on cognition in stroke and traumatic brain injury patients. *NeuroRehabilitation* 40(1):33–48. <https://doi.org/10.3233/NRE-161388>
- Visser ME, Durao S, Sinclair D, Irlam JH, Siegfried N (2017) Micronutrient supplementation in adults with HIV infection. *Cochrane Database Syst Rev* 5:CD003650. <https://doi.org/10.1002/14651858.CD003650.pub4>
- Webel AR, Moore SM, Hanson JE, Patel SR, Schmotzer B, Salata RA (2013) Improving sleep hygiene behavior in adults living with HIV/AIDS: a randomized control pilot study of the SystemCHANGE(TM)-HIV intervention. *Appl Nurs Res* 26(2):85–91. <https://doi.org/10.1016/j.apnr.2012.10.002>
- Webel AR, Barkley J, Longenecker CT, Mittelsteadt A, Gripshover B, Salata RA (2015) A cross-sectional description of age and gender differences in exercise patterns in adults living with HIV. *J Assoc Nurses AIDS Care* 26(2):176–186. <https://doi.org/10.1016/j.jana.2014.06.004>
- Webel AR, Perazzo JD, Dawson-Rose C, Smith C, Nicholas PK, Rivero-Mendez M et al (2017) A multinational qualitative investigation of the perspectives and drivers of exercise and dietary behaviors in people living with HIV. *Appl Nurs Res* 37:13–18. <https://doi.org/10.1016/j.apnr.2017.07.002>
- Wight D, Wimbush E, Jepson R, Doi L (2016) Six steps in quality intervention development (6SQuID). *J Epidemiol Community Health* 70(5):520–525. <https://doi.org/10.1136/jech-2015-205952>

- Wirth MD, Jagers JR, Dudgeon WD, Hebert JR, Youngstedt SD, Blair SN et al (2015) Association of markers of inflammation with sleep and physical activity among people living with HIV or AIDS. *AIDS Behav* 19(6):1098–1107. <https://doi.org/10.1007/s10461-014-0949-y>
- Witteman HO, Presseau J, Nicholas Angl E, Jokhio I, Schwalm JD, Grimshaw JM et al (2017) Negotiating tensions between theory and design in the development of mailings for people recovering from acute coronary syndrome. *JMIR Hum Factors* 4(1):e6. <https://doi.org/10.2196/humanfactors.6502>
- Womack JA, Murphy TE, Bathulapalli H, Akgun KM, Gibert C, Kunisaki KM et al (2017) Trajectories of sleep disturbance severity in HIV-infected and uninfected veterans. *J Assoc Nurses AIDS Care* 28(3):431–437. <https://doi.org/10.1016/j.jana.2017.02.006>
- Woods SP, Dawson MS, Weber E, Gibson S, Grant I, Atkinson JH et al (2009a) Timing is everything: antiretroviral nonadherence is associated with impairment in time-based prospective memory. *J Int Neuropsychol Soc* 15(1):42–52. <https://doi.org/10.1017/S1355617708090012>
- Woods SP, Moore DJ, Weber E, Grant I (2009b) Cognitive neuropsychology of HIV-associated neurocognitive disorders. *Neuropsychol Rev* 19(2):152–168. <https://doi.org/10.1007/s11065-009-9102-5>
- Wu J, Wu H, Lu C, Guo L, Li P (2015) Self-reported sleep disturbances in HIV-infected people: a meta-analysis of prevalence and moderators. *Sleep Med* 16(8):901–907. <https://doi.org/10.1016/j.sleep.2015.03.027>
- Xu L, Jiang CQ, Lam TH, Liu B, Jin YL, Zhu T et al (2011) Short or long sleep duration is associated with memory impairment in older Chinese: the Guangzhou Biobank Cohort Study. *Sleep* 34(5):575–580
- Yaffe K, Falvey CM, Hoang T (2014) Connections between sleep and cognition in older adults. *Lancet Neurol* 13(10):1017–1028. [https://doi.org/10.1016/S1474-4422\(14\)70172-3](https://doi.org/10.1016/S1474-4422(14)70172-3)
- Zazpe I, Sanchez-Tainta A, Estruch R, Lamuela-Raventos RM, Schroder H, Salas-Salvado J et al (2008) A large randomized individual and group intervention conducted by registered dietitians increased adherence to Mediterranean-type diets: the PREDIMED study. *J Am Diet Assoc* 108(7):1134–1144. Discussion 1145. <https://doi.org/10.1016/j.jada.2008.04.011>
- Ziegler TR, McComsey GA, Frediani JK, Millson EC, Tangpricha V, Eckard AR (2014) Habitual nutrient intake in HIV-infected youth and associations with HIV-related factors. *AIDS Res Hum Retrovir* 30(9):888–895. <https://doi.org/10.1089/AID.2013.0282>