UC San Diego

UC San Diego Previously Published Works

Title

Root Inputs Influence Soil Water Holding Capacity and Differentially Influence the Growth of Native versus Exotic Annual Species in an Arid Ecosystem

Permalink

<https://escholarship.org/uc/item/9x91s9fn>

Journal

Restoration Ecology, 22(6)

ISSN

1061-2971

Authors

Mckinney, J
Cleland, EE

Publication Date

2014-11-01

DOI

10.1111/rec.12139

Peer reviewed

1 Title: Root inputs influence soil water holding capacity and differentially influence the
2 growth of native versus exotic annual species in an arid ecosystem

3

4 Submission Category: Research article

5

6 Authors: Jordan McKinney & Elsa E. Cleland*

7

8 Affiliations: Ecology, Behavior & Evolution Section, University of California San Diego

9 * Corresponding author contact:

10

11 Address: 9500 Gilman Dr.#0116, University of California San Diego, La Jolla CA 92093

12 USA

13 Phone: 8586999928

14 Email: ecleland@ucsd.edu

15

16 Keywords: carbon addition, climate change, invasion, nitrogen immobilization, water
17 pulses

Abstract

Invasion by exotic annual species is increasingly impacting Southern California aridlands, altering ecosystem processes and plant community composition. With climate change the Southwestern US is expected to experience increasingly variable rainfall. Larger rainfall events could facilitate invasion by exotic species that can capitalize on high resource conditions. Exotic annual species also have dense shallow root systems that could create positive feedbacks to further invasion by increasing soil organic matter and water holding capacity. Alternatively, fine root inputs could create negative feedbacks to exotic plant growth if they stimulate microbial nutrient immobilization. The dual influences of rainfall regime and fine root inputs on species performance were evaluated in an experiment where native and exotic species were grown individually and in combination under varying watering regimes (large infrequent or small frequent pulses, holding total rainfall constant) and root additions (with or without sterilized exotic roots). Mean soil moisture increased with larger infrequent watering events, and also with root addition. Plant growth (both native and exotic) increased with larger watering events, but declined with root addition. Exotic species growth declined more than native species growth with root additions. Mechanistically, root addition lowered inorganic nitrogen (N) availability, and microbial N immobilization increased with soil moisture content. Together these results show that increased fine root production promotes negative feedbacks to growth of exotic species via microbial N immobilization, especially under conditions of high soil moisture. Further, our results suggest that organic carbon additions are a potentially effective strategy for suppressing growth of problematic desert invaders.

Introduction

Exotic annual species are problematic invaders across western North America, especially in arid and semi-arid regions (Mack 1981, Lovett & Bainbridge 1999, Craig et al. 2010), where they can negatively impact native species (Barrows et al 2009) and alter ecosystem functioning by accelerating the fire regime (Brooks et al. 2004, Schiermeier 2005). Environmental changes such as altered climate and nitrogen (N) deposition may

49 be facilitating these invasion, favoring exotic grasses over native desert species (Rao &
50 Allen 2010). Invasions may become increasingly common in these areas because the
51 Southwestern U.S. is expected to be a climate change “hot spot” in the coming decades,
52 associated with increasingly variable rainfall patterns (Diffenbaugh et al. 2008). The
53 abundance of exotic annual species often increases in high rainfall years (Hobbs et al.
54 2007, Bethany & Mustard 2005), but we know much less about how native and exotic
55 species respond to variation in the magnitude of rainfall events within years, although it
56 has been predicted that more variable rainfall regimes will promote invasion (Knapp et al.
57 2008).

58 Positive plant-soil feedbacks may play an important role in facilitating the spread
59 of problematic invaders (Levine et al. 2006). Exotic species can alter soil microbial
60 communities (Klironomos 2002; Callaway et al. 2004), through inputs of allelopathic
61 chemicals (Bais et al. 2003; but see Lind & Parker 2010), alteration of light availability
62 via the litter layer (Lenz et al. 2003) or by altered nutrient cycling (Ehrenfeld 2003;
63 Evans et al. 2001). One potential feedback mechanism that has been surprisingly untested
64 is the role of root proliferation by exotic annual species, and subsequent additions of
65 organic carbon (C) into soils. Exotic annual grasses often have high below-ground
66 production of dense, fine roots (Holmes & Rice 1996); potentially creating a positive
67 feedback, promoting invasion via by increasing soil water holding capacity.

68 Alternatively, C additions in mesic systems often increase microbial activity and hence
69 immobilization of limiting nutrients, suppressing the growth of exotic species and thus
70 favoring native species (Blumenthal et al. 2003; Perry et al. 2010). The proliferation of
71 fine root biomass associated with invasion by exotic annual species in arid and semi-arid
72 environments could thus lead to a mix of positive (via enhanced soil water holding
73 capacity) and negative feedbacks (via microbial nutrient immobilization) to growth of
74 exotic species. Here we performed a greenhouse experiment to evaluate competition
75 between exotic annual species and native desert forbs under varying soil moisture
76 regimes, both with and without added fine root biomass. We tested the following
77 hypotheses: 1) We expect all species have higher growth with higher soil moisture when
78 grown individually, but exotic species will outperform native species when grown in
79 competition under high soil moisture conditions; 2) Larger, less frequent precipitation

events maintain high soil moisture and disproportionately benefit invasive species; 3) High root inputs from exotic species enhances soil water holding capacity, thus creating a positive feedback to enhance invasion success; 4) Alternatively, C inputs via the roots of exotic species could promote microbial activity, immobilize nutrients, and suppress plant growth. We evaluated these hypotheses for a suite of species commonly found in the aridlands of San Diego County, with the goal of evaluating two potential restoration strategies (C addition, seeding native species) to favor native species over problematic invaders.

Methods

The experiment was conducted in greenhouses at the University of California, San Diego Biological Field Station from January to April 2013. We selected four native focal annual wildflower species which commonly flower in San Diego County each spring, including *Amsinckia tessellata* (bristly fiddleneck), *Malacothrix glabrata* (desert dandelion), *Laesthenia californica* (California goldfields), and *Layia platyglossa* (tidy tips). We selected four problematic exotic species (designated by the California Invasive Plant Council) also common in San Diego County, including *Bromus hordeaceus* (soft brome), *Bromus madritensis* spp. *rubens* (red brome), *Erodium cicutarium* (redstem filaree), and *Schismus barbatus* (common Mediterranean grass). Each species was grown individually as well as together in the eight species mixture (competitive environment) in each watering regime (see treatment description below) and soil organic matter content (five grams sterilized exotic roots versus no roots added) in a factorial block design (n=8, for a total of 288 pots).

Pots were filled with commercially available native topsoil (sandy loam, Agri Service Inc., Vista California). Exotic grass root biomass was collected from greenhouse-grown plants washed free of soil and sterilized by steam autoclaving. These roots were cut into ~ one cm pieces and five grams of root biomass were mixed into the top five cm of soil in each pot in the root addition treatment. No roots were added to the control pots.

Seeds of the focal species were purchased commercially (S & S Seeds, Carpinteria, California) and germination percentages were evaluated on watered field

soils prior to the start of the experiment. Each pot was sown with enough seeds to achieve a target of 24 individuals, (three individuals of each species for the pots where all eight species were grown in competition). Seeds were planted on January 26 and an initial pulse of 500 mL of water was added on February 4. Seedlings of *E. cicutarium* (collected from the UC Elliot Chaparral Reserve) were transplanted on February 7, 2013 due to poor germination success in the greenhouse.

Subsequent watering was initiated on February 9. The small, frequent watering regime added a 60 mL pulse of water every five days, and the large, infrequent watering regime added a 120 mL pulse of water every 10 days. Soil moisture measurements were taken daily during one 10-day watering cycle soon after seeds germinated using a Field Scout TDR 100 Soil Moisture Sensor measuring the volumetric water content. Pots were rotated in blocks every two weeks to account for spatial variation in greenhouse light availability. Watering continued for 47 days, at which time plants were harvested after showing signs of senescence. All above ground biomass was harvested from each plot, dried for 48 hours at 40° C, and weighed.

The top five cm of soil were collected from each pot that contained all eight focal species, and sieved to remove roots and debris. Potassium chloride was used to extract soils for inorganic N (Keeney & Nelson, 1987) and chloroform fumigation and extraction was performed to measure microbial biomass N (Brookes et al., 1985) and C (Beck et al., 1997). Potassium persulfate digestion (Delia et al. 1977) was used to oxidize all organic and inorganic N to nitrate, subsequently absorbance protocols (Doane & Horwath 2003) were used to determine the amount of nitrate in each sample.

Statistical Analysis

Statistical tests were performed with R version 2.15.0 (R Development Core Team 2013). Soil moisture was analyzed using a linear mixed-effect model (nlme R package, Pinheiro et al. 2014) with species origin (native, exotic, or mixed), watering regime (frequent or infrequent), roots (added or not added), and block as factorial fixed factors and individual pot ID as a random factor to account for repeated measures. The growth responses of the focal species were evaluated with a linear mixed effect model

where block, competitive environment (species grown individual versus in combination), species origin (native or exotic), watering regime and roots were included as factorial fixed factors and species identity nested within origin was included as a random factor. Growth responses to the treatments were also assessed for each species individually using a linear model where competitive environment, water and roots were included as factorial fixed factors and block as a random factor. Inorganic N, and microbial N and C were predicted with a linear model where the influence of watering regime and root treatments were factorially assessed with block position as a random factor. We performed type II analysis of deviance tests to assess statistical significance (car R package, Fox and Weisberg 2011). Tukey's Honest Significant Difference tests were performed as posthoc tests when interactions were identified.

When species were grown individually, pot biomass values were divided by eight, because 1/8 of the mass of seeds of each focal species were sown in the eight species competition pots, as compared to the mass planted when each species was grown individually. Hence, in this analysis we compare the observed growth of species when grown in multi-species communities, and compare that value to the expected biomass of that species when grown individually and seeded at the same rate.

Additional information regarding the source and characterization of soils and roots used in this experiment are provided in the Supplementary Material, along with all statistical tables.

Results

Pots supplemented with roots and watered with larger but less frequent events had higher mean (Water $\chi^2_1 = 13.83$, $p < 0.001$; Roots $\chi^2_1 = 19.82$, $p < 0.001$) and minimum (Water $\chi^2_1 = 7.94$, $p < 0.001$; Roots $\chi^2_1 = 19.53$, $p = 0.005$) soil moisture (Fig. 1B; Table S1). Although we performed these measurements soon after germination at a time when we expected minimal impacts of plant growth on soil moisture, species composition had a small but significant effect on mean soil moisture, whereby the pots planted with all 8 species had slightly lower soil moisture (Mixed moisture = $5.79 \text{ cm}^3/\text{cm}^3$, Exotic moisture = $6.42 \text{ cm}^3/\text{cm}^3$, Native moisture = $6.27 \text{ cm}^3/\text{cm}^3$; Origin $\chi^2_1 = 6.21$, $p = 0.045$;

Fig. 1C).

Plant performance, as measured by biomass, was significantly influenced by watering regime, species origin and root additions, with interactions (Fig. 2; Table S1). Watering with large, infrequent events increased plant biomass (Watering regime $\chi^2_1 = 25.60$, $p < 0.001$) and this effect was greater for exotic species (Origin x Water $\chi^2_1 = 4.76$, $p = 0.029$). In contrast, root addition decreased overall plant biomass (Root addition $\chi^2_1 = 3.89$, $p = 0.049$). When roots were added there was little difference between exotic and native plants while exotics produced significantly higher biomass when grown without root additions (Root x Origin $\chi^2_1 = 4.41$, $p = 0.036$). Block was a significant effect in our analysis ($\chi^2_7 = 53.24$, $p < 0.001$), because one block had consistently lower biomass due to unknown factors (data not shown).

Most species had higher biomass (standardized by seeding rate) when grown alone than when grown in combination (significant only for the native *M. glabrata*, and exotics *E. cicutarium*, and *B. hordeaceus*; Fig. 3, Table S2). However, both the native forb *A. tessellata* and the exotic grass *S. barbatus* showed higher biomass when grown in combination with other species (Competitive environment: *A. tessellata* $\chi^2_1 = 5.04$, $p = 0.025$; *S. barbatus* $\chi^2_1 = 27.17$, $p < 0.001$; Fig. 3A & E). Species also displayed variation in their responses to the experimental treatments. Qualitatively, *S. barbatus* and *A. tessellata* showed the most growth under all treatments compared with other species when grown in competition, suggesting that their responses likely drove the exotic and native species responses respectively (Fig. 3, Table S2). Growth of *S. barbatus* was promoted by infrequent watering, and this effect was greater in the pots where all species were grown in combination (Water x Competition $\chi^2_1 = 6.08$, $p = 0.014$; Fig. 3E, Table S2). *S. barbatus* declined with root addition in the combination pots but not when grown alone (Root x Competition $\chi^2_1 = 6.47$, $p = 0.011$). Infrequent, large watering events also increased growth for *B. madritensis*, *B. hordeaceus*, *L. californica*, *M. glabrata* and *E. cicutarium*, although this effect was only significant for the latter two species when grown individually (Fig. 3, Table S2). *E. cicutarium* showed higher growth when roots were added in combination pots, however when grown individually the species had higher growth when no roots were added (Root x Competition $\chi^2_1 = 4.20$, $p = 0.040$).

Inorganic N availability in soils at the end of the experiment was significantly

influenced by an interaction between root addition and watering regime (Root x Water $\chi^2_1 = 5.15$, $p = 0.23$; Table S3); root addition decreased soil inorganic N availability, and this effect was strongest in pots with large but infrequent watering events. Microbial biomass N concentrations were also higher in pots with infrequent watering treatments (Water $\chi^2_1 = 13.97$, $p < 0.001$; Table S3, Fig. 4). Treatments had no significant effects on microbial biomass C (Water $\chi^2_1 = 0.45$, $p = 0.505$; Root $\chi^2_1 = 0.05$, $p = 0.825$; Root x Water $\chi^2_1 = 1.04$, $p = 0.308$; Table 3).

Discussion

We found that larger, infrequent watering events maintained higher average soil moisture than when the same water quantity was applied in small, frequent events, potentially because larger watering events allowed water to reach deeper soil layers, hence minimizing evaporative loss (Fay et al. 2008). Higher soil moisture in turn increased plant biomass; these results are consistent with a prior study in a semi-arid grassland that found larger infrequent rainfall events tended to increase soil moisture and plant growth (Heisler-White et al. 2009). The large, infrequent watering treatment in our experiment disproportionately increased the growth of exotic species, supporting our hypothesis that exotic species would outperform native species under higher soil moisture. While exotic annual species can have dramatically positive responses to years of high rainfall (e.g. Hobbs et al. 2007), some studies have found no impact of soil moisture on direct competition between native and exotic species (Maron and Marler 2008) suggesting that exotic species may not always benefit more than native species from high soil moisture.

We also found that root addition increased mean soil moisture content, consistent with prior studies showing that increasing soil organic matter content can increase soil watering holding capacity (Blumenthal et al. 2003); however, the increase in soil moisture caused by root addition did not increase plant growth. Instead, root addition suppressed plant growth. Our analysis showed root addition suppressed exotic species as a group more than native species (although this was likely driven by the significant response of one species – *S. barbatus*). Microbial biomass N increased with infrequent

watering, suggesting that larger water pulses stimulated microbial activity to a greater degree (e.g. Austin et al. 2004). Inorganic N was reduced under infrequent watering and root addition, consistent with the idea that soil microbes immobilized inorganic N when C was added in the form of fine roots. Adding C to soils in mesic systems often causes soil microbes to immobilize N, favoring the growth of native species (Blumenthal et al. 2003; Alpert & Maron 2000). Thus, root addition in our study may have favored slower growing native species over fast growing exotic species by limiting the available N in the soil (Alpert and Maron 2000, Perry et al. 2010, Bobbink 1991). Our results show that this effect was greatest when soil moisture was high (infrequent but larger watering events). In a study similar to ours, Blumenthal (2009) found that C additions in concert with water addition reduced invasive plant growth in a semi-arid grassland system. Still, the results of this study are consistent with C addition studies in the Mojave desert, showing that C amendments can stimulate increased microbial N immobilization even under xeric conditions (Scheffer et al 2003, Steers et al. 2011).

As a group, we found that exotic species in our experiment tended to competitively suppress the growth of native species, but there was significant variation among our focal species in their responses. In particular *A. tessellata* had the highest growth among the native species when grown in competition (Fig. 3A), and actually had higher biomass when grown in combination than when grown alone, suggesting this species was facilitated by growing in a diverse community. Other studies have also found that *A. tessellata* was robust to exotic competition (Abella et al. 2011; Brooks 2009), potentially due to traits such as large seed size and early-season germination (Salo et al. 2005). Hence, *A. tessellata* could be an effective choice for restoration efforts attempting to establish native cover via seeding, especially when there may be potential competition from exotic species present in the seedbank.

There was also variation among the exotic focal species in their responses to the experimental treatments, with potential implications for management of these problematic invasive species. *E. cicutarium* growth was significantly suppressed when grown in the multi-species competitive environment, suggesting that seeding native species can be an effective strategy for reducing the abundance of this common desert invader. *E. cicutarium* growth was also reduced by low soil moisture (our small, frequent

watering treatment). This species is a common invader of many arid systems but has been shown to decline substantially in drought years (Blackshaw et al. 2000). Our results suggest that restoration efforts employing irrigation to promote germination of native seeds should be cautious to prevent over-watering in areas where *E. cicutarium* could be present in the seedbank, as high soil moisture favors this exotic species over the common native species used in this experiment. Timed herbicide applications have also been found to be effective for managing this particularly early-active species, and are more effective for increasing native cover than management efforts such as weeding that disturb soils (Marushia et al. 2010).

Schismus barbatus is a common exotic grass in Southern California deserts (Brooks 1999) and was the dominant species in pots where our eight focal species were grown in competition. Further, biomass (scaled for seeding rate) was higher for this species when grown with multiple species than when grown in a monoculture; hence, efforts to re-seed native species are unlikely to be successful if this strong competitor is present in the seedbank. However, *S. barbatus* growth was significantly lowered by root addition, suggesting that organic C addition could be a feasible management strategy to combat invasions of *S. barbatus*. This problematic invader has been shown to benefit from N enrichment (Brooks 2003; DeFalco et al. 2001), and a study that used sucrose additions similarly found that labile C additions reduced growth of *S. barbatus* under field conditions (Steers et al. 2011), thus C additions are a logical management strategy with the potential to lower N availability and hence the competitive dominance of this species.

Management efforts focused on reducing soil N availability have generally used sawdust or mulch as sources for C addition (Wilson & Gerry 1995; Zink & Allen 1998; Alpert & Maron 2000; Paschke et al. 2000; Blumenthal et al. 2003). The results of this study suggest that the biomass of exotic annual species could be left in situ as a C source, provided it is harvested before the plants set seed, and there is not a danger of re-sprouting from intact root stocks. However, exotic litter can be a fuel source promoting fire, and so fuel reduction is a major goal for many land managers (Brookes et al. 2004), hence management efforts aimed at maintaining litter with the goal of lowering soil N availability would need to consider associated fire risks of this strategy. Additionally the

timing of C amendments could be important: Early season C amendments are likely to disproportionately suppress exotic annual species in this region (Salo 2004), because they generally have flexible germination cues that allow them to emerge earlier and faster at the start of the growing season (Wainwright & Cleland 2013). Consistent with this idea, Steers et al. (2011) found that C amendment reduced exotic annual grass abundance most in a year when they germinated first, while native species and exotic annual forbs were impacted in a year when the timing of C amendment coincided with their emergence.

Overall, the results of this study demonstrate how inputs of organic C from the fine roots of exotic species can initiate a complex set of positive and negative feedbacks to invasion, and that the outcome of these interactions depends on environmental context (rainfall regime), biotic interactions (interspecific competition), and species identity. In particular, while we found that exotic species benefited more from high soil moisture (resulting from larger, but less frequent watering events), exotic species growth declined more than native species when roots were added to soil, even though root additions also increased soil moisture. This shows that in this case, the negative effect of microbial nutrient immobilization outweighed the positive influence of increased soil moisture for plant growth when exotic roots were added to soil. Although our results highlight general mechanistic feedbacks that may be applicable for understanding the outcome of invasions in many systems, we also found significant variation among species in their responses to our experimental treatments – both within and between groups based on native versus exotic origin – that may aid in identifying effective management strategies for particularly problematic desert invaders.

Implication for Practice

- Carbon additions in arid environments can reduce exotic plant growth via enhanced microbial nutrient immobilization; this effect varies among species and is greatest when soil moisture is high.
- *Schismus barbatus* growth was significantly suppressed by C addition, suggesting this would be a successful strategy to suppress growth of this invasive species.
- *Erodium cicutarium* was suppressed by interspecific competition, suggesting that

seed addition could be a strategy to combat this problematic invader.

- *Amsinckia tessellata* was a strong competitor with exotic plant species; this widespread species is found in a variety of arid and semi-arid California vegetation types, suggesting this species is a prime candidate for seeding in restoration.

Acknowledgements

We thank Edie Allen for constructive comments on an earlier version of this manuscript. We are grateful for help in analysis of soil samples provided by Ellen Esch and Porrachia Magali. JGM was partially supported by NSF DEB #1154082.

References

- Abella, S.R., Craig, D.J., Chiquoine, L.P., Prengaman, K.A., Schmid, S.M., and T.M. Embrey. 2011. Relationships of native desert plants with red brome (*Bromus rubens*): Toward identifying invasion-reducing species. *Invasive Plant Science and Management* 4:115-124.
- Alpert, P., and J.L., Maron. 2000. Carbon addition as a countermeasure against biological invasion by plants. *Biological Invasions* 2:33-40.
- Austin, A.T., Yahdjian, L., Stark, J.M., Belnap, J., Porporato, A., Norton, U., Ravetta, D.A., and S.M. Schaeffer. 2004. Water pulses and biogeochemical cycles in arid and semiarid ecosystems. *Oecologia* 141: 221-235.
- Bais, H.P., Vepachedu, R., Gilroy, S., Callaway, R.M., and J.M. Vivanco. 2003. Allelopathy and exotic plant invasion: from molecules and genes to species interactions. *Science* 301.5638:1377-1380.
- Barrows, C.W., Allen, E.B., Brooks, M.L. & Allen, M.F. 2009. Effects of an invasive plant on a desert sand dune landscape. *Biological Invasions*, 11, 673–686.
- Beck, T., Joergensen, R.G., Kandeler, E., Makeschin, F., Nuss, E., Oberholzer, H.R., and S. Scheu. 1997. An inter-laboratory comparison of ten different ways of measuring soil microbial biomass carbon. *Soil Biology Biochemistry* 29:1023-1032.
- Bradley, B. A., and Mustard, J. F. (2005). Identifying land cover variability distinct from land cover change: cheatgrass in the Great Basin. *Remote Sensing of Environment*, 94,

358 204-213.

359 Blackshaw, R.E., Semach, G.P., and J.T. O'Donovan. 2000. Utilization of wheat seed
 360 rate to manage Redstem Filaree (*Erodium cicutarium*) in a zero-tillage cropping system.
 361 *Weed Technology* 14:389-396.

362 Blumenthal, D., Jordan, N., and M. Russelle. 2003. Soil carbon addition controls weeds
 363 and facilitates prairie restoration. *Ecological Applications* 13:605-615.

364 Blumenthal, D.M. 2009. Carbon addition interacts with water availability to reduce
 365 invasive forb establishment in a semi-arid grassland. *Biological Invasions* 11:1281-1290.

366 Bobink, R. 1991. Effects of nutrient enrichment in Dutch chalk grassland. *Journal of*
 367 *Applied Ecology* 28:28-41.

368 Brooks, M.L. 1999. Habitat invisibility and dominance by alien annual plants in the
 369 western Mojave Desert. *Biological Invasions* 1:325-337.

370 Brooks, M.L. 2003. Effects of increased soil nitrogen on the dominance of alien annuals
 371 plants in the Mojave Desert. *Journal of Applied Ecology* 40:344-353.

372 Brooks, M.L. 2009. Spatial and temporal distribution of non-native plants in upland areas
 373 of the Mojave Desert. Pages 101–124 in R. H. Webb, L. F. Fenstermaker, J. S. Heaton,
 374 D. L. Hughson, E. V. McDonald, and D. M. Miller, eds. The Mojave Desert: Ecosystem
 375 Processes and Sustainability. Reno, NV: University of Nevada Press.

376 Brooks, M.L., D'Antonio, C.M., Richardson, D.M., Grace, J.B., Keeley, J.E., DiTomaso,
 377 J.M., Hobbs, R.J., Pellant, M., and D. Pyke. 2004. Effects of invasive alien plants on fire
 378 regimes. *BioScience* 54:677-688.

379 Brookes, P.C., Landman, A., Pruden, G., and D.S. Jenkinson. 1985. Chloroform
 380 fumigation and the release of soil nitrogen: a rapid direct extraction method to measure
 381 microbial biomass nitrogen in soil. *Soil Biology Biochemistry* 17:837-842.

382 Callaway, R.M., Thelen, G.C., Rodriguez, A. and W.E. Holben. 2004. Soil biota and
 383 exotic plant invasion. *Nature* 427:731–733.

384 Craig, D.J., Craig, J.E., Abella, S.R. & Vanier, C.H. 2010. Factors affecting exotic annual
 385 plant cover and richness along roadsides in the eastern Mojave Desert, USA. *Journal of*
 386 *Arid Environments*, 74, 702–707.

387 DeFalco, L.A., Detling, J.K., Tracy, C.R., and S.D. Warren. 2001. Physiological variation
 388 among native and exotic winter annual plants associated with microbiotic crusts in the

389 Mojave Desert. *Plant and Soil* 234:1-14.

390 Delia, C.T., Steudler, P.A., and N. Corwin. 1977. Determination of total nitrogen in
391 aqueous samples using persulfate digestion. *Limnology and Oceanography* 22:760-764.

392 Diffenbaugh, N.S., Giorgi, F., and J.S. Pal. 2008. Climate change hotspots in the United
393 States. *Geophysical Research Letters* 35:L16709.

394 Doane, T.A. and W.R. Horwath. 2003. Spectrophotometric determination of nitrate with
395 a single reagent. *Analytical Letters* 36:2713-2722.

396 Ehrenfeld, J.G. 2003. Effects of exotic plant invasions on soil nutrient cycling processes.
397 *Ecosystems* 6.6:503-523.

398 Evans, R.D., Rimer, R., Sperry, L., and J. Belnap. 2001. Exotic plant invasion alters
399 nitrogen dynamics in an arid grassland. *Ecological Applications* 11:1301-1310.

400 Fay, P.A., Kaufman, D.M., Nippert, J.B., Carlisle, J.D., and C.W. Harper. 2008. Changes
401 in grassland ecosystem function due to extreme rainfall events: implications for responses
402 to climate change. *Global Change Biology* 14:1600-1608.

403 Fox, J. and S. Weisberg. 2011. *An R Companion to Applied Regression, Second Edition*.
404 Thousand Oaks California, Sage.

405 Heisler-White, J. L., Blair, J. M., Kelly, E. F., Harmony, K. and Knapp, A. K. 2009.
406 Contingent productivity responses to more extreme rainfall regimes across a grassland
407 biome. *Global Change Biology* 15: 2894–2904.

408 Hobbs, R.J., Yates, S., and H.A. Mooney. 2007. Long-term data reveal complex
409 dynamics in grassland in relation to climate and disturbance. *Ecological Monographs*
410 77:545- 568.

411 Holmes, T.H., and K.J. Rice. 1996. Patterns of growth and soil-water utilization in some
412 exotic annuals and native perennial bunchgrasses of California. *Annals of Botany* 78:233-
413 243.

414 Keeney, D.R. and D.W. Nelson. 1987. Nitrogen--Inorganic Forms, sec. 33-3, extraction
415 of exchangeable ammonium, nitrate, and nitrite. pp.648-9. In A. L. Page et al., eds.,
416 Methods of Soil Analysis: Part 2, Chemical and Microbiological Properties. Soil Science
417 Society of America, Madison, Wisconsin USA.

418 Klironomos, J.N. 2002. Feedback with soil biota contributes to plant rarity and
419 invasiveness in communities. *Nature* 417:67– 70.

420 Knapp, A.K., Beier, C., Briske, D.D., Classen, A.T., Luo, Y., Reichstein, M., Smith,
 421 M.D., Smith, S.D., Bell, J.E., Fay, P.A., Heisler, J.L., Leavitt, S.W., Sherry, R., Smith,
 422 B., and Weng, B. 2008. Consequences of more extreme precipitation regimes for
 423 terrestrial ecosystems. *BioScience* 58:811-821.
 424 Lind E.M., and J.D. Parker. 2010. Novel weapons testing: Are invasive plants more
 425 chemically defended than native plants? *PLoS ONE* 5(5): e10429.
 426 Lenz, T.I., Moyle-Croft, J.L. and J.M. Facelli. 2003. Direct and indirect effects of exotic
 427 grasses on species composition of a South African grassland. *Aust. Ecol* 28:23–32.
 428 Lovich, J. E., & Bainbridge, D. 1999. Anthropogenic degradation of the southern
 429 California desert ecosystem and prospects for natural recovery and restoration.
 430 *Environmental Management*, 24, 309-326.
 431 Maron, J. L., & Marler, M. (2008). Field-based competitive impacts between invaders
 432 and natives at varying resource supply. *Journal of Ecology*, 96, 1187-1197.
 433 Mack, R.N. 1981. Invasion of *Bromus tectorum* L. into western North America: an
 434 ecological chronicle. *Agro-Ecosystems* 7:145–165.
 435 Marushia, R. G., Cadotte, M. W. and Holt, J. S. 2010. Phenology as a basis for
 436 management of exotic annual plants in desert invasions. *Journal of Applied Ecology*, 47:
 437 1290–1299.
 438 Paschke, M.W., McLendon, T., and E.F. Redente. 2000. Nitrogen availability and old-
 439 field succession in a short-grass steppe. *Ecosystems* 3:144-158.
 440 Perry, L.G., Blumenthal, D.M., Monaco, T.A., Paschke, M.W., and E.F. Redente. 2010.
 441 Immobilizing nitrogen to control plant invasion. *Oecologia* 163.1:13-24.
 442 Pinheiro J, Bates D, DebRoy S, Sarkar D and R Core Team (2014). *nlme: Linear and*
 443 *Nonlinear Mixed Effects Models*. R package version 3.1-117
 444 R Development Core Team. 2013. *R: A language environment for statistical computing*.
 445 R Foundation for Statistical Computing, Vienna, Austria.
 446 Rao, L.E., and E.B. Allen. 2010. Combined effects of precipitation and nitrogen
 447 deposition on native and invasive winter annual production in California deserts.
 448 *Oecologia* 162:1035-1046.
 449 Salo, L. F. 2004. Population dynamics of red brome (*Bromus madritensis* subsp. *rubens*):
 450 times for concern, opportunities for management. *Journal of Arid Environments* 57:291–

451 296.

452 Salo, L.F., McPherson, G.R., and D.G. Williams. 2005. Sonoran Desert winter annuals
 453 affected by density of red brome and soil nitrogen. *The American Midland Naturalist*
 454 153:95- 109.

455 Schiermeier, Q. 2005. Pall hangs over desert's future as alien weeds fuel wildfires.
 456 *Nature*, 435, 724.

457 Steers, R. J. Funk, J. L., and E. B. Allen. 2011. Can resource-use traits predict native vs.
 458 exotic plant success in carbon amended soils? *Ecological Applications* 21:1211–1224.

459 Wainwright, C. E., and E. E. Cleland. 2013. Exotic species display greater germination
 460 plasticity and higher germination rates than native species across multiple cues.
 461 *Biological invasions* 15:2253-2264.

462 Wilson, S.D., and A.K. Gerry. 1995. Strategies for mixed-grass prairie restoration:
 463 herbicide, tilling, and nitrogen manipulation. *Restoration Ecology* 3:290-298.

464 Zink, T.A., and M.F. Allen. 1998. The effects of organic amendments on the restoration
 465 of a disturbed coastal sage scrub habitat. *Restoration Ecology* 6:52-58.

466

Figure 1. Mean and min soil moisture levels for one watering cycle (10 days), in each of the treatments manipulating watering regime and root additions. In the small frequent watering regime 60 mL of water was added every 5 days, while the infrequent watering regime received a single 120 mL pulse every 10 days.

Figure 2. Above-ground biomass (mean +SE) for native and exotic species grown individually and in competition, for each treatment combination of root addition and watering frequency. Note biomass values are averaged across species to illustrate the statistical significance of the factors in the analysis.

481 Figure 3. Above-ground biomass responses to varying watering regime and root addition
482 treatments for native (A- D) and exotic (E-H) species grown individually (grey) or in
483 interspecific competition with the other focal species (open).

484

Figure 4. Concentrations of soil extractable inorganic nitrogen (A) and microbial biomass nitrogen (B) for combination of watering regime and root addition treatments. Letters distinguish significant differences between means based on post-hoc tests.