

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### Title

Sterility Produced in Mice by Deuterium Oxide

### Permalink

<https://escholarship.org/uc/item/9xx458ff>

### Authors

Hughes, Ann M.  
Calvin, Melvin

### Publication Date

1957-12-06

UCRL 8048

UNIVERSITY OF  
CALIFORNIA

*Radiation  
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UNIVERSITY OF CALIFORNIA

Radiation Laboratory  
Berkeley, California

Contract No. W-7405-eng-48

STERILITY PRODUCED IN MICE BY DEUTERIUM OXIDE

Ann M. Hughes and Melvin Calvin

December 6, 1957

STERILITY PRODUCED IN MICE BY DEUTERIUM OXIDE

Ann M. Hughes and Melvin Calvin

Radiation Laboratory and Department of Chemistry  
University of California, Berkeley 4, California

December 6, 1957

ABSTRACT

Sterility in  $C_{57}$  and Swiss mice has been produced by substituting  $D_2O$  for a part of the drinking water. The effective range lies between 5% and 30%  $D_2O$ . It appears that the effect is greater in  $C_{57}$  males than females, and that the size and viability of the litter is affected.

## STERILITY PRODUCED IN MICE BY DEUTERIUM OXIDE

Ann M. Hughes and Melvin Calvin

Radiation Laboratory and Department of Chemistry,  
University of California, Berkeley 4, California

December 6, 1957

The present availability of  $D_2O$  at a reasonable price has stimulated an increased investigation of its physiological effects. The inhibition of ascites tumor growth and of algal reproduction has recently been reported by this laboratory<sup>1, 2</sup> and others.<sup>3</sup>

We have demonstrated the production of sterility in mice by the substitution of  $D_2O$  for a part of the drinking water. In the first experiment, six female and six male  $C_{57}$  mice that had been maintained on 30%  $D_2O$  in the drinking water for 4 weeks were mated.<sup>1</sup> The animals were housed 3 females and 3 males to a cage.  $D_2O$  administration was continued for 10 weeks. Since there were no pregnancies at the end of this time, the  $D_2O$  was discontinued. At the end of another 8 weeks, when there still were no pregnancies, three of the treated females were mated with three normal males; three of the treated males were mated with three normal females. Although the mating of  $D_2O$ -treated females with normal males resulted in litters at the end of 3 weeks, all offspring died within 24 hours and two of the mothers died. The mating of  $D_2O$ -treated males with normal females did not produce offspring until the end of 13 weeks, at which time one female littered. From the 3  $D_2O$ -treated males and 3  $D_2O$ -treated females remaining together, one female<sup>2</sup> littered in 4 additional weeks, one in 10 additional weeks. This experiment is graphically represented in Fig. 1.

In the second experiment, both  $C_{57}$  and Swiss mice were used. Five mice of each sex of each strain were maintained for 4 weeks on either 30%, 20%, or 5%  $D_2O$  in the drinking water. At the end of this treatment period,  $D_2O$  administration was discontinued and each mouse was individually mated with a normal mouse of the same strain. At the same time, six pairs of normal mice of each strain were mated. Only those litters born during a 30-day period after the beginning of the mating are included in the results.

---

<sup>1</sup>All  $D_2O$  percentages are expressed as volume percent, approximately equal to atoms percent.

<sup>2</sup>As used in this paper, the term mated indicates that males and females were housed together continuously.

<sup>1</sup>Hughes, Tolbert, Lonberg, and Calvin, *Biochim. et Biophys. Acta* (in press).

<sup>2</sup>Holm-Hansen, Moses, and Calvin; *Biochim. et Biophys. Acta* (in press).

<sup>3</sup>A. J. Finkel and D. Czajka; *Proc. Am. Assoc. Cancer Research*, 2, 201 (1957).

Our data, summarized in Table I, indicate that  $D_2O$  causes a sterility in both  $C_{57}$  and Swiss mice, and that the effect is greater in the  $C_{57}$  mice. It also appears that there is a greater effect in  $C_{57}$  males than in females, and that the size and viability of the litter is affected. Further experiments are in progress to verify these latter observations and to investigate some metabolic effects of  $D_2O$  in mice.

The failure of Hansen and Wulfert<sup>4</sup> to observe sterility in mice as a result of  $D_2O$  is probably due to the low  $D_2O$  concentration they used by them.

This work was done under the auspices of the U.S. Atomic Energy Commission.


---

<sup>4</sup>H. Hansen and K. Wulfert, Arch. exptl. Pathol., Pharmakol. Naunyn-Schneidberg's 190, 671 (1938).

Table I

The effect of  $D_2O$  on the fertility of  $C_{57}$  or Swiss Mice, presented as the number of surviving offspring per mated pair.

Sex treated with $D_2O$	$D_2O$ conc. in drinking water (%)	Number surviving (offspring per mated pair)	
		$C_{57}$	Swiss
Male	5	1.2	7.2
Female	5	2.4	7.6
Male	20	1.0	7.4
Female	20	2.5	5.8
Male	30	0.0	2.0
Female	30	2.2	3.2
Controls	0	3.3	8.5



MU-14357

Fig. 1. The production of sterility in  $C_{57}$  mice by the administration of 30%  $D_2O$  in the drinking water. Horizontal arrows indicate length of time animals were given  $D_2O$  or  $H_2O$ . Vertical arrows indicate time of mating or littering. Dotted lines indicate time of cross-mating the treated with the normal animals, as described in the test.