

UC Davis

UC Davis Previously Published Works

Title

Effect of Mootral™ - a garlic and citrus extract based feed additive - on enteric methane emissions in feedlot cattle

Permalink

<https://escholarship.org/uc/item/9z60s7jr>

Journal

Translational Animal Science, 3(4)

ISSN

2573-2102

Authors

Roque, Breanna M
van Lingen, Henk J
Vrancken, Hilde
[et al.](#)

Publication Date

2019-07-01

DOI

10.1093/tas/txz133

Peer reviewed

Effect of Mootral—a garlic- and citrus-extract-based feed additive—on enteric methane emissions in feedlot cattle

Breanna M. Roque,[†] Henk J van Lingen,[†] Hilde Vrancken,[‡] and Ermias Kebreab^{†,1}

[†]Department of Animal Science, University of California Davis, Davis, CA 95616; and [‡]Mootral GmbH, Berlin 13467, Germany

ABSTRACT: Enteric methane (CH₄) production is the main source of greenhouse gas emissions from livestock globally with beef cattle contributing 5.95% of total global greenhouse gas emissions. Various mitigation strategies have been developed to reduce enteric emissions with limited success. In vitro studies have shown a reduction in CH₄ emissions when using garlic and citrus extracts. However, there is paucity of data regarding in vivo studies investigating the effect of garlic and citrus extracts in cattle. The objective of this study was to quantitatively evaluate the response of Angus × Hereford cross steers consuming the feed additive Mootral, which contains extracts of both garlic and citrus, on CH₄ yield (g/kg dry matter intake [DMI]). Twenty steers were randomly assigned to two treatments: control (no additive) and Mootral supplied at 15 g/d in a completely randomized design with a 2-wk covariate and a 12-wk data collection periods. Enteric CH₄ emissions were

measured using the GreenFeed system during the covariate period and experimental weeks 2, 6, 9, and 12. CH₄ yield (g/kg DMI) by steers remained similar in both treatments for weeks 2 to 9. In week 12, there was a significant decrease in CH₄ yield (23.2%) in treatment compared to control steers mainly because the steers were consuming all the pellets containing the additive. However, overall CH₄ yield (g/kg DMI) during the entire experimental period was not significantly different. Carbon dioxide yield (g/kg DMI) and oxygen consumption (g/kg DMI) did not differ between treatments during the entire experimental period. DMI, average daily gain, and feed efficiency also remained similar in control and supplemented steers. The in vivo results showed that Mootral may have a potential to be used as a feed additive to reduce enteric CH₄ production and yield in beef cattle but needs further investigation under various dietary regimens.

Key words: environmental sustainability, greenhouse gas, methane yield, steers

© The Author(s) 2019. Published by Oxford University Press on behalf of the American Society of Animal Science. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Transl. Anim. Sci. 2019.3:1383–1388
doi: 10.1093/tas/txz133

INTRODUCTION

Global emissions of greenhouse gases (GHG) have risen to unprecedented levels despite a growing

number of policies to reduce climate change (International Panel for Climate Change (IPCC), 2014). Anthropogenic sources account for 58% of global GHG emissions (National Academies of Sciences, Engineering, and Medicine, 2018). Methane (CH₄) from enteric fermentation of livestock is the largest contributor with beef cattle contributing 5.95% to global GHG emissions (Gerber et al., 2013).

¹Corresponding author: ekebreab@ucdavis.edu

Received February 20, 2019.

Accepted August 9, 2019.

There are limited strategies to reduce enteric fermentation that can be widely applied to ruminants. Diet manipulation (i.e., changing substantial amount of diet ingredients) and feed additives constitute two strategies with the greatest potential. [Moraes et al. \(2014\)](#) investigated the potential of diet manipulation to reduce enteric CH₄ emissions and concluded that it is possible to reduce emissions through the reduction of neutral detergent fiber (NDF) levels in the diet. However, [Moraes et al. \(2015\)](#) calculated that the cost of implementing this strategy could be up to 49% greater than diets formulated according to national recommendations (e.g., [National Academies of Sciences, Engineering, and Medicine, 2016](#)). Therefore, feed additives may be a better alternative to reduce enteric emissions in a cost-effective manner. Mootral is a feed additive containing allicin obtained from garlic as well as a byproduct of orange processing, referred to citrus extract. In an in vitro study, [Busquet et al. \(2005\)](#) reported that garlic oil (300 mg/L) and allicin (300 mg/L) decreased CH₄ production 73.6% and 19.5%, respectively, compared to control diet consisting of 50:50 forage to concentrate ratio in a 24 h incubation. Further, [Ma et al. \(2016\)](#) investigated the effects of supplementary allicin in sheep diet on CH₄ emissions and reported a 6% decrease compared to control (scaled to metabolic body weight [BW]; L/kg BW^{0.75}) when given at a dose of 2 g/d for 42 d. [Kim et al. \(2012\)](#) incubated serum bottles containing 0.3 g of timothy grass and plant extracts (1% of total volume) for 24 h in vitro. The authors reported that citrus extract reduced CH₄ emissions 16.7% and garlic extract 20% compared to control with no plant extracts.

We hypothesize that supplementing diets with a combination of garlic and citrus extracts will result in reduced enteric CH₄ production. The objective of this study was to quantitatively evaluate the response of Angus × Hereford cross steers consuming the feed additive Mootral, which contains extracts of both garlic and citrus, on CH₄ yield (g/kg dry matter intake [DMI])

MATERIALS AND METHODS

This study was approved by the Institutional Animal Care and Use Committee at the University of California, Davis (Protocol No. 20032).

Study Design, Animals, and Diets

The study consisted of 20 Angus × Hereford cross steers that were blocked by initial BW, to

reduce weight variability, then randomly allocated to one of the following treatments: control (no additive) and Mootral supplied at a dose of 15 g/d for the duration of the trial. The steers were individually housed and were approximately 12 months in age with an average BW of 419 ± 16 kg at the beginning of the trial. The experiment followed a completely randomized design, with a 2-wk covariate and a 12-wk data collection periods. Mootral was first pressed in to an alfalfa pellet then delivered to each of the treatment animals along with their daily total mixed ration (TMR). The control group received “blank” pellets with their daily TMR to ensure Mootral was the only difference between the two treatments. Details on the Mootral supplement formulation has been published by [Eger et al. \(2018\)](#).

Daily intake was calculated as the TMR and alfalfa pellets offered subtracted by individual feed refusal weights. Steers were fed 105% of the previous day's intake twice daily at 0600 and 1800 hours. Steers were fed TMR ([Table 1](#)) formulated to meet or exceed their growth requirement according to the [National Academies of Sciences, Engineering, and Medicine \(2016\)](#) recommendations. Two feedlot diets, a starter and finisher diet were formulated ([Table 1](#)). The starter diet was used as a backgrounding diet to acclimate the animals to a low forage TMR whereas the finisher diet was used as a typical low forage, feedlot diet. Steers were offered water ad libitum.

Sample Collection and Analysis

CH₄, carbon dioxide (CO₂), and oxygen gas emissions from cows were measured using the GreenFeed system (C-Lock, Inc, Rapid City, SD). Gas emissions using the GreenFeed system were measured during the covariate period and experimental weeks 2, 6, 9, and 12. During each measurement period, gas emission data were collected during three consecutive days as follows: starting at 0700, 1300, and 1900 hours (sampling d 1); 0100, 1000, and 1600 hours (sampling d 2); and 2200 and 0400 hours (sampling d 3). Breath gas samples were collected for at least 5 min followed by a 2-min background gas sample collection. The GreenFeed unit was calibrated weekly with a standard gas mixture containing (mol %): CO₂, 0.98, CH₄, 0.151, and the balance being nitrogen gas (Air Liquide America Specialty Gases, Rancho Cucamonga, CA). Recovery rates for both CO₂ and CH₄ observed in this study were between +/- 1% of the known quantities of gas that was released. Alfalfa

pellets were used as bait feed and was offered at each sampling event and was kept below 5% of the total DMI during each sampling period. Alfalfa pellets consumed at the GreenFeed machine contained no Mootral additive, regardless of treatment. The composition of alfalfa pellets is shown in Table 2.

BW was measured once a week to monitor growth rate and the average daily gain (ADG) was calculated from the BW measurements. TMR was sampled once a week and analyzed for dry matter, acid detergent fiber, NDF, lignin, crude

fat, total digestible nutrient and mineral content (Cumberland Valley Analytical Services, Waynesboro, PA).

Statistical Analysis

Statistical analysis was performed using the open-source R statistical software (version 3.1.1; The R Foundation for Statistical Computing, Vienna, Austria.). Statistical analysis was completed using the linear mixed-effects models (lme) procedure, with the steer serving as the experimental unit. GreenFeed emission data were averaged per steer and gas measurement period and the averaged data used in the statistical analysis. Weekly DMI, growth, and gas emission data were analyzed as repeated measure with a rational quadratic spatial correlation structure. The statistical model included treatment, week, and treatment \times week interactions, and the covariate term, with the error term assumed to be normally distributed with mean = 0 and constant variance. Individual animal was used as random effect, whereas all other factors were considered fixed. Statistical significance was established when $P \leq 0.05$ and a trend at $0.05 < P \leq 0.10$.

RESULTS AND DISCUSSION

Although there are several feed additives that have shown potential to reduce CH_4 emissions (e.g., Dijkstra et al. 2018; Roque et al. 2019), currently none are available that can be applied in a commercial setting. This section presents the results of the gas emission analysis and impact on productivity due to supplementation with Mootral.

Gas Production

Enteric CH_4 yield by steers in the control and Mootral-supplemented groups are shown in Fig. 1A. There was no statistical differences in baseline CH_4 yield between the control and supplemented groups. During the subsequent weeks 2, 6 and 9, although the CH_4 yield from supplemented steers remained below the control, the two treatments were not statistically different (week 2 control = 18.0 ± 1.4 g/kg DMI, treatment = 17.5 ± 1.5 g/kg DMI; week 6 control = 10.3 ± 1.4 g/kg DMI, treatment = 9.5 ± 1.5 g/kg DMI and week 9 control = 18.2 ± 1.4 g/kg DMI, treatment = 15.6 ± 1.5 g/kg DMI). CH_4 yield was significantly different between control and treatment in week 12 (19.4 ± 1.4 g/kg DMI vs. 14.9 ± 1.5 g/kg DMI, respectively). The differences in CH_4 yield between treatment and

Table 1. Formulation of starter and finisher rations

Ingredients	Starter (% of dry matter)	Finisher (% of dry matter)
Forage		
Alfalfa hay	15.0	6.00
Wheat hay	12.0	4.00
Dry distillers grain	20.0	5.00
Concentrate		
Rolled corn grain	42.0	
Flaked corn		74.0
Molasses	8.00	5.00
Fat	1.50	3.00
Urea	0.35	0.50
Beef trace salt ¹	0.32	1.00
Magnesium Oxide		0.20
Limestone	0.82	0.70
Phosphate monosodium		0.10
Potassium chloride		0.40

¹Beef Trace Salt sourced from AL Gilbert.

Table 2. Chemical composition of starter diet, finisher diet, and alfalfa pellets (% of dry matter)

Nutrients	Starter TMR	Finisher TMR	Alfalfa pellets
% Dry matter			
Crude protein	14.1	13.5	21.9
Acid detergent fiber	18.8	9.5	30.2
NDF	29.1	17.4	42.1
Lignin	3.41	1.93	7.02
Crude fat	5.29	6.94	2.17
Total digestible nutrients	76.7	85.3	60.9
Ash	6.02	4.78	10.4
Ca	0.49	0.48	1.13
Phosphorus	0.34	0.27	0.3
Magnesium	0.27	0.22	0.59
Potassium	1.44	0.92	2.1
Sodium	0.15	0.36	0.16
Parts per million			
Iron	119	124	1319
Manganese	47	56	55
Zinc	49	59	25
Copper	7	7	12




Figure 1. CH₄ yield (g CH₄/kg DMI) (A), and CO₂ yield (g CO₂/kg DMI) (B) by steers in the control (solid line) and treatment (broken line) groups during the 12-wk experimental period.

control were greater in the last two sampling periods compared to the beginning weeks of the experiment. Visual observation led us to speculate that some of the differences may be attributed to sorting of the feed as some pellets were not consumed. This could be attributed to individual animal preference, or lack thereof, for alfalfa pellets. As the experiment progressed, the steers consumed the daily doses as recommended. Over the course of the trial, the mean enteric CH₄ yield of steers supplemented with Mootral was 13.3% lower compared to control steers but it was not significantly different (16.5 ± 1.0 g/kg DMI, vs. 14.3 ± 1.1 g/kg DMI, for control and supplemented groups, respectively; $P = 0.16$). There were week-to-week variability in CH₄ yield, however, even after removing week 6 from the analysis, which were lower compared to other weeks, the results were not affected.

Although Ma et al. (2016) did not use the same formulation as in this study, allicin was the main ingredient in both studies. The authors used similar concentrations of allicin when scaled to BW and observed reduced CH₄ production by over 160 mL/

kg BW^{0.75}. Acetate production was also reduced while butyrate and iso-butyrate productions were increased. CH₄ production has been positively associated with greater molar proportions of acetate and negatively associated with increased propionate and butyrate production (Alemu et al. 2011). Proportions of acetate, butyrate, and propionate determine the quantity of hydrogen available in the rumen for utilization by CH₄-producing microbes. Pathways resulting in propionate production contribute the least to the quantity of hydrogen, whereas pathways resulting in acetate contribute the most.

The mode of action of allicin is thought to be through reductions in protozoa and methanogenic archaea populations in the rumen (Ma et al. 2016). Miron et al. (2000) showed that allicin is highly permeable through membranes and may contribute to its biological activity. Furthermore, Eger et al. (2018) reported that mixture of garlic and citrus compounds appeared to reduce CH₄ production by altering the archaeal community such that the percentage of *Methanobacteriaceae* was reduced

without exhibiting negative side effects on rumen fermentation.

Compared to stockers, feedlot cattle produce less enteric CH₄ (6.5% vs. 3% Gross Energy intake; IPCC, 2006), mainly because the diet contains highly digestible carbohydrates, which changes the molar proportions of volatile fatty acid production resulting in a greater propionate to acetate ratio. Beauchemin et al. (2010) estimated that 53% of the enteric CH₄ from beef production comes from the cow-calf herd whereas only 10% is attributed to the feedlot cattle. Therefore, feed additives including Mootral have a greater potential to reduce CH₄ production when given to cattle fed forage based diets rather than concentrate diets.

CO₂ production throughout the experiment (Fig. 1B) was similar between the two groups and was not significantly different ($P = 0.48$). Similarly, Hristov et al. (2015) using a feed additive to reduce CH₄ emissions also reported that CO₂ production were not different between control and treatment. The mean oxygen consumption between the two experimental group was also not significantly different ($P = 0.48$; data not shown).

BW Changes

DMI of steers in the control group and treatment groups were also similar; 9.94 vs. 9.47 kg/d respectively (± 0.55 kg, $P = 0.46$). The initial and final BW measurements show that there was no differences in the production efficiency between control and Mootral-supplemented steers (initial 419 vs. 428 kg, respectively [± 16.3 kg, $P = 0.59$]; final 557 vs. 569 kg, respectively [± 19.9 kg, $P = 0.57$]). There was a large day-to-day variability in BW; therefore, a larger number of animals compared to those used in the experiment would have been required to accurately estimate differences in BW due to treatment, if any. ADG was calculated as differences in BW between two consecutive measurements periods. There was no statistical difference in ADG between control and Mootral treatment; 1.65 vs. 1.67 kg/d respectively (± 0.10 , $P = 0.80$). Hence, no significant differences were observed between the feed conversion efficiencies of between the treatments ($P = 0.96$).

In conclusion, steers fed a diet with Mootral added at a rate of 15 g/d had a 23% reduction in CH₄ yield after 12 weeks of supplementation. Despite the average CH₄ production over the entire experimental period not being different between the two dietary treatments, the steers receiving the Mootral

treatment had lower CH₄ yields than the steers receiving the control treatment over time with no effect on DMI, ADG, and feed conversion efficiency.

ACKNOWLEDGMENTS

This research received financial support from Mootral GmbH. We are grateful to Don Harper, David Gall, and several undergraduate interns from Department of Animal Science, UC Davis that participated in the trial.

Conflict of interest statement. None declared.

LITERATURE CITED

- Alemu, A. W., J. Dijkstra, A. Bannink, J. France, and E. Kebreab. 2011. Rumen stoichiometric models and their contribution and challenges in predicting enteric methane production. *Anim. Feed Sci. Tech.* 166–167:761–778. doi:10.1016/j.anifeeds.2011.04.054
- Beauchemin, K. A., H. Janzen, S. M. Little, T. A. McAllister, and S. M. McGinn. 2010. Life cycle assessment of greenhouse gas emissions from beef production in western Canada: a case study. *Agric. Syst.* 103:371–379. doi:10.1016/j.agsy.2010.03.008
- Busquet, M., S. Calsamiglia, A. Ferret, M. D. Carro, and C. Kamel. 2005. Effect of garlic oil and four of its compounds on rumen microbial fermentation. *J. Dairy Sci.* 88:4393–4404. doi:10.3168/jds.S0022-0302(05)73126-X
- Dijkstra, J., A. Bannink, J. France, E. Kebreab, and S. van Gastelen. 2018. Short communication: antimethanogenic effects of 3-nitrooxypropanol depend on supplementation dose, dietary fiber content, and cattle type. *J. Dairy Sci.* 101:9041–9047. doi:10.3168/jds.2018-14456
- Eger, M., M. Graz, S. Riede, and G. Breves. 2018. Application of Mootral™ reduces methane production by altering the archaea community in the rumen simulation technique. *Front. Microbiol.* 9:2094. doi:10.3389/fmicb.2018.02094
- Gerber, P. J., H. Steinfeld, B. Henderson, A. Mottet, C. Opio, J. Dijkman, A. Faluccci, and G. Tempio. 2013. Tackling climate change through livestock—a global assessment of emissions and mitigation opportunities. Food and Agriculture Organization of the United Nations (FAO), Rome.
- Hristov, A. N., J. Oh, F. Giallongo, T. W. Frederick, M. T. Harper, H. L. Weeks, A. F. Branco, P. J. Moate, M. H. Deighton, S. R. Williams, et al. 2015. An inhibitor persistently decreased enteric methane emission from dairy cows with no negative effect on milk production. *Proc. Natl. Acad. Sci. U. S. A.* 112:10663–10668. doi:10.1073/pnas.1504124112
- IPCC. 2006. IPCC Guidelines for National Greenhouse Gas Inventories. Institute for Global Environmental Strategies, Hayama, Japan.
- IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the IPCC. IPCC, Geneva, Switzerland.
- Kim, E. T., C. H. Kim, K. S. Min, and S. S. Lee. 2012. Effects of plant extracts on microbial population, methane emission and ruminal fermentation characteristics in *in vitro*. *Asian-Australas. J. Anim. Sci.* 25:806–811. doi:10.5713/ajas.2011.11447

- Ma, T., D. D. Chen, Y. Tu, N. F. Zhang, B. W. Si, K. D. Deng, and Q. Y. Diao. 2016. Effect of supplementation of allicin on methanogenesis and ruminal microbial flora in Dorper crossbred ewes. *J. Anim. Sci. Biotech.* 7 Article Number: 1. doi: 10.1186/s40104-015-0057-5
- Miron, T., A. Rabinkov, D. Mirelman, M. Wilchek, and L. Weiner. 2000. The mode of action of allicin: its ready permeability through phospholipid membranes may contribute to its biological activity. *Biochim. Biophys. Acta* 1463:20–30. doi:10.1016/s0005-2736(99)00174-1
- Moraes, L. E., A. B. Strathe, J. G. Fadel, D. P. Casper, and E. Kebreab. 2014. Prediction of enteric methane emissions from cattle. *Glob. Chang. Biol.* 20:2140–2148. doi:10.1111/gcb.12471
- Moraes, L. E., J. Fadel, A. Castillo, D. Casper, J. Tricarico, and E. Kebreab. 2015. Modeling the trade-off between diet costs and methane emissions: A goal programming approach. *J. Dairy Sci.* 98:5557–5571.
- National Academies of Sciences, Engineering, and Medicine. 2018. Improving characterization of anthropogenic methane emissions in the United States. Natl. Acad. Press, Washington, DC.
- National Academies of Sciences, Engineering, and Medicine. 2016. Nutrient requirements of beef cattle: eighth revised edition. Natl. Acad. Press, Washington, DC. doi:10.17226/19014.
- Roque, B. M., J. K. Salwen, R. Kinley, and E. Kebreab. 2019. Inclusion of *Asparagopsis armata* in lactating dairy cows' diet reduces enteric methane emission by over 50 percent. *J. Cleaner Prod.* 234:132–138. doi:10.1016/j.jclepro.2019.06.193