

# UCLA

## UCLA Previously Published Works

### Title

Identifying fates of cancer cells exposed to mitotic inhibitors by quantitative phase imaging

### Permalink

<https://escholarship.org/uc/item/9zk5d18s>

### Journal

Analyst, 145(1)

### ISSN

0003-2654

### Authors

Huang, Dian  
Roy, Irena J  
Murray, Graeme F  
et al.

### Publication Date

2019-12-16

### DOI

10.1039/c9an01346f

Peer reviewed



Published in final edited form as:

Analyst. 2019 December 16; 145(1): 97–106. doi:10.1039/c9an01346f.

## Identifying fates of cancer cells exposed to mitotic inhibitors by quantitative phase imaging

Dian Huang<sup>1</sup>, Irena J. Roy<sup>2</sup>, Graeme F. Murray<sup>3</sup>, Jason Reed<sup>3,4</sup>, Thomas A. Zangle<sup>5</sup>, Michael A. Teitell<sup>1,2,6,7</sup>

<sup>1</sup>Department of Bioengineering, University of California at Los Angeles, Los Angeles, CA 90095

<sup>2</sup>Department of Pathology and Laboratory Medicine, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA 90095

<sup>3</sup>Department of Physics, Virginia Commonwealth University, Richmond, VA 23284

<sup>4</sup>Massey Cancer Center, Virginia Commonwealth University, Richmond, VA, USA

<sup>5</sup>Department of Chemical Engineering and Huntsman Cancer Institute, University of Utah, Salt Lake City, UT 84112

<sup>6</sup>Molecular Biology Institute, Broad Center for Regenerative Medicine and Stem Cell Research, and California NanoSystems Institute, University of California at Los Angeles, Los Angeles, CA 90095

<sup>7</sup>Department of Pediatrics and Jonsson Comprehensive Cancer Center, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA 90095

### Abstract

Cell cycle deregulation is a cancer hallmark that has stimulated the development of mitotic inhibitors with differing mechanisms of action. Quantitative phase imaging (QPI) is an emerging approach for determining cancer cell sensitivities to chemotherapies *in vitro*. Cancer cell fates in response to mitotic inhibitors are agent- and dose-dependent. Fates that lead to chromosomal instabilities may result in a survival advantage and drug resistance. Conventional techniques for quantifying cell fates are incompatible with growth inhibition assays that produce binary live/dead results. Therefore, we used QPI to quantify post-mitotic fates of G0/G1 synchronized HeLa cervical adenocarcinoma and M202 melanoma cells during 24 h of escalating-dose exposures to mitotic inhibitors, including microtubule inhibitors paclitaxel and colchicine, and an Aurora kinase A inhibitor, VX-680. QPI determined cell fates by measuring changes in cell biomass, morphology, and mean phase-shift. Cell fates fell into three groups: (1) bipolar division from drug failure; (2) cell death or sustained mitotic arrest; and (3) aberrant endocycling or multipolar division. In this proof-of-concept study, colchicine was most effective in producing desirable outcomes of sustained mitotic arrest or death throughout its dosing range, whereas both paclitaxel


Correspondence: tzangle@chemeng.utah.edu or mteitell@mednet.ucla.edu.

#### CONFLICTS OF INTEREST

M.A.T. is a co-founder, board member, shareholder, and consultant for NanoCav, LLC, a private start-up company working on quantitative phase microscopy techniques and applications. T.A.Z. is consultant for NanoCav, LLC. The other authors do not have any conflicting interests to declare.

and VX-680 yielded dose-dependent multipolar divisions or endocycling, respectively. Furthermore, rapid completion of mitosis associated with bipolar divisions whereas prolonged mitosis associated with multipolar divisions or cell death. Overall, QPI measurement of drug-induced cancer cell fates provides a tool to inform the development of candidate agents by quantifying the dosing ranges over which suboptimal inhibitor choices lead to undesirable, aberrant cancer cell fates.

## Graphical Abstract



A novel analysis method for cell fate responses to mitotic inhibitors using biophysical attributes obtained through quantitative phase imaging.

## INTRODUCTION

Developing effective anti-cancer treatment regimens remains a significant therapeutic challenge. Treatment selection based on available diagnostic data including assessment of histologic tumor subtype, clinical grade and stage, molecular biomarkers, and genome-profiling studies can still lead to variable patient outcomes. This indicates a pressing need to continue developing new agents and regimens.<sup>1</sup> The prediction of treatment outcomes and selection of therapeutic agents typically relies upon drug performance studies from preclinical research and clinical trials. In these settings, drug performance assessments are most commonly by multi-day growth inhibition assays *in vitro* and tumor shrinkage *in vivo*. However, data from these binary analyses may fail to uncover processes within cancer cells that further increase drug resistance and tumor aggressiveness. Cancer cells that persist after therapy may acquire additional genetic or epigenetic changes that make future treatments progressively more difficult.<sup>2</sup> Therefore, a method that captures the full range of cell fates after a specific treatment, for example with a mitotic inhibitor, could reveal unsuspected suboptimal drug regimens with an elevated risk of promoting more aggressive tumors.

Specific mitotic inhibitors have frequent use to treat specific cancers, such as paclitaxel for breast and ovarian cancers. As a group they target the microtubule system or associated cell division kinases with the goal of activating growth checkpoints to induce mitotic arrest and apoptosis of cancer cells.<sup>3</sup> Despite widespread use, most mitotic inhibitors show neurotoxicity, poor *in vivo* efficacy, and are difficult to dose adequately, thereby limiting applications.<sup>4</sup> Preclinical studies also reveal that post-treatment surviving cancer cells may aberrantly exit from mitosis with multipolar cell divisions or endocycling from a weakened mitotic checkpoint caused by suboptimal mitotic inhibitor dosing.<sup>5, 6</sup> These mitotic aberrations may cause aneuploidy, chromosome instability, and increased tumor aggression.<sup>7-9</sup> A rapid method to detect and classify aberrant mitotic outcomes for mitotic inhibitor treated cancers could improve drug development and selection.

Flow cytometry that uses DNA intercalating dyes, confocal microscopy, fluorescence time-lapse microscopy, and multi-day growth inhibition assays are current methods for assessing cellular responses to mitotic inhibitors.<sup>5, 6, 10</sup> Unfortunately these approaches are often laborious, can be cell destructive, are limited to discrete measurement time points that can miss emerging therapy resistance, or require labeling that may interfere with cell behavior. For example, the most commonly practiced multi-day growth inhibition assays only provide total numbers of viable or dead cancer cells in tissue culture at specific treatment time points.  $EC_{50}$  values generated from this type of counting assay only shows population trends and overlooks phenotypic outcomes of individual cancer cells that survive treatment. This approach therefore yields limited insight into drug response kinetics and potential aberrant outcomes.

To overcome limitations in current screening methods and to increase throughput, we deployed a version of quantitative phase imaging (QPI) we refer to as live cell interferometry (LCI) to measure single cell responses to three mitotic inhibitors with different mechanisms of action using dose-escalating drug concentrations. Current state-of-the-art QPI techniques, including digital holographic microscopy and spatial light interference microscopy, accurately quantify optical path length delays caused by cellular contents at submicron resolution, free of labeling agents and phototoxicity effects.<sup>11, 12</sup> Measurements of phase shifts are then processed to produce biophysical cell parameters, such as dry mass and mass transport.<sup>11, 12</sup> These relevant biophysical properties can be analyzed in a wide range of QPI applications, such as measurements of cytotoxicity.<sup>13, 14</sup> Our approach uses quadriwave lateral shearing interferometry (QWLSI) to precisely quantify the phase-shift of incident light interacting with the non-aqueous mass, or biomass, of individual cells.<sup>15</sup> The QWLSI approach has been shown to be accurate over a range of different specimen thickness and insensitive to imaging location, magnification, or degree of spatial light coherence.<sup>16</sup> Conversion of measured phase-shifts in light into biomass uses an experimentally determined cell average specific refractive index, which enables quantifying changes in cell biomass over time.<sup>17, 18</sup> Prior LCI studies revealed breast cancer cell line sensitivities to trastuzumab (Herceptin) within 6 h, a speed compatible for studies of patient biopsy materials, with results replicating multi-day growth inhibition assays.<sup>19, 20</sup> More recently, LCI successfully dissected tumor heterogeneity and drug resistance for melanoma cells in a mixture<sup>21</sup> and could replicate known tumor sensitivities to cisplatin in mouse patient-derived xenograft (PDX) models of breast cancer.<sup>22</sup> These prior LCI studies

validated QPI utility in cancer, but did not evaluate cancer cell outcomes beyond binary growth inhibition results. The use of multi-parametric QPI response profiling data that could further inform preclinical drug development and clinical drug selection is an exciting possibility explored here.

In this proof-of-concept study, we provide a new multi-parametric analytical method to identify different cell fate outcomes to mitotic inhibitors using QPI measurements of cell biomass, morphology, and mean phase-shift of light. Our study provides dynamic data on mitotic inhibitor activities and the frequencies of abnormal and undesirable outcomes during early exposure time points that may make tumors more difficult to treat.<sup>7</sup> Our cell fate identification strategy may also be useful for developing and testing other anticancer agents and regimens.

## MATERIALS AND METHODS

### Cells and cell culture

HeLa human cervical adenocarcinoma cells were from the American Type Culture Collection (ATCC) and M202 human melanoma cells were a gift from Dr. Owen Witte (UCLA). HeLa cells were maintained in 1:1 DME/F-12 media (Thermo Fisher Scientific) and M202 cells were maintained in RPMI 1640 media (Thermo Fisher Scientific), with each media supplemented by 10% FBS (Omega Scientific), 100 U/mL penicillin (Corning), 100 µg/mL streptomycin (Corning) and 2 mmol/l-glutamine (Thermo Fisher Scientific).

### Growth inhibition assay

Twelve-well flat bottom plates (Thermo Fisher Scientific) received  $5 \times 10^4$  cells/well. Paclitaxel (Sigma-Aldrich), colchicine (Sigma-Aldrich), or VX-680 (Selleckchem) small molecule mitotic inhibitors, or DMSO (Sigma-Aldrich) carrier-control, were added to cell culture media at the indicated doses and durations (Fig. S1, S2). Cells from three replicate wells per treatment condition were harvested each day, stained with trypan blue, and counted using an automated cell counter (Countess; Invitrogen).

### Cell preparation for QPI

ibidi 4-well <sup>Ph+</sup> µ-slides received  $1.5 \times 10^4$  cells/mL that were then grown for 7–10 h to homeostasis. Media containing 2mM thymidine (Sigma-Aldrich) was added to arrest HeLa cells in G0/G1 phase for 18 h and M202 cells for 20 h. Synchronized cells were released from cell cycle block by media washing three times. Fresh media with the indicated doses of paclitaxel, colchicine, VX-680, or DMSO were then added to the 4-well µ-slide and then sealed with anti-evaporation oil (ibidi) before QPI on the microscope stage.

### Live cell interferometry

QPI of HeLa and M202 cells was performed on an Axio Observer A1 inverted microscope (Zeiss) with a SID4Bio quadriwave lateral shearing interferometry (QWLSI) camera (Phasics). A temperature and CO<sub>2</sub> regulated stage-top cell incubation chamber (ibidi) was fit to a motorized xy stage (Thorlabs) to maintain environmental homeostasis and enable QPI at multiple locations. A Zeiss LD Plan Neofluar 20x NA 0.4 objective was heated and

maintained at 37°C with a custom built copper objective heater driven by a heat controller (Thorlabs). Trans-illumination was by a 660 nm center wavelength collimated LED (Thorlabs). Image collection occurred every 10 min for 24 h at 15 randomly selected imaging locations per well containing cells plated with sufficient spacing to enable automated image processing and biomass segmentation. Imaging locations were selected within the central part of each well, where the highest quality quantitative phase information can be obtained without aberrations caused by the cell culture apparatus or optical hardware. The selected imaging fields cover a range of locations within each imaged well, to obtain a sample that is representative of the cell population (Fig. S3).

### QPI data analysis

Interferograms captured by the SID4Bio QWLSI camera were converted to phase-shift and intensity images using the Phasics MATLAB software development kit. These images were analyzed using custom MATLAB (MathWorks) scripts that eliminated background aberrations by fitting a 4<sup>th</sup> order Zernike surface to cell-free regions and subtracting the fitted surface from the image. Cell biomass data was extracted from background corrected images by integrating light phase-shift in segmented cell projected areas and multiplying by the experimentally determined specific refractive increment of 0.0018<sup>17, 23, 24</sup>. Data analysis of cell fates and time spent in mitosis are performed using morphology as discussed in the following section. The entire procedure requires 3 – 4 hours for image processing and data analyses, depending on the number of locations imaged in each experiment.

### Morphology metrics

**Mean phase shift:** The mean phase-shift of a cell is obtained by dividing the total integrated phase- shift for that particular cell by its projected area. Interphase flat and spread-out versus round and mitotic phase HeLa cells were sampled to establish average mean phase-shift values for both morphologies. Since HeLa and M202 cells share comparable sizes, morphologies, optical densities and the same tracking criteria for the MATLAB tracking algorithm, the same average mean phase- shift values are representative for both cell types (Fig. S4).

**Shape factor:** The shape factor of a cell, also called circularity or isoperimetric quotient, is calculated by dividing the projected area,  $A$ , of a cell by its circumference or the length of its perimeter,  $P$ , ( $4\pi A/P^2$ ). Random flat HeLa and M202 cells and mitotic HeLa cells were sampled to establish average shape factor values for both morphologies (Fig. S4). A perfect circle has a shape factor of 1.0 and an irregular shape, such as an interphase adherent cell, has a shape factor of approximately 0.5 (Fig. S4).

### Flow cytometry

Cells were collected from T25 flasks after 24 and 48 h of drug exposure, washed once with 1x PBS, pH 7.4, and then re-suspended in 500 $\mu$ L of FxCycle™ PI/RNase Staining Solution (Thermo Fisher Scientific). Flow cytometry was performed on FACS BD LSRII and FACS BD Fortessa flow cytometers (BD Biosciences). DNA content analysis was by FlowJo software.

## Statistical analysis

Chi-square tests of independence were performed on contingency tables assessing the observed cell fate counts per inhibitor dosage for significant association relative to expected counts in each drug treatment panel with a 95% confidence interval (Prism 6, GraphPad, Inc.). Dose response curve-fitting was performed using the curve fitting toolbox in MATLAB (MathWorks). Two-sample Kolmogorov-Smirnov (KS) tests were performed between bipolar cell division versus prolonged mitotic arrest and cell death fate distributions for each drug treatment panel in MATLAB with a 95% confidence interval. One-way ANOVA with unbalanced sample groups was performed on duration of mitosis datasets between cell fate groups in MATLAB with a 95% confidence interval. Statistical significance required  $p < 0.05$ .

## RESULTS

### Identifying post-mitotic cell fate outcomes using QPI

We deployed QPI to measure synchronized HeLa and M202 single cancer cell biomass and morphology responses after mitotic entry, in the presence of escalating doses of several mitotic inhibitors (Fig. 1, Table 1). We chose paclitaxel, a microtubule stabilizing agent, colchicine, a microtubule destabilizing drug, and VX-680, an Aurora kinase A inhibitor that represent a range of mitotic inhibitor modes of action. Cells were treated with each drug for 5 h before QPI of randomly selected locations at 10 min imaging intervals over the next 24 h (Fig. 1A). Changes in biophysical and morphological parameters over the imaging period provided data to determine cell fate outcomes of single cancer cells in each condition (Fig. 1B-C). Two  $EC_{50}$  values, one for successful bipolar divisions representing drug failure and one for cell death/arrest, were calculated based on the distributions of cell fate outcomes in the sampled cell populations (Fig. 1D).  $EC_{50}$ , the difference between these two  $EC_{50}$  values, therefore describes the concentration range over which cancer cells display aberrant mitotic exits at suboptimal dosages for each type of drug treatment (Fig. 1D).

We classified the mitotic fate of each tracked cancer cell into one of five categories: (1) successful bipolar division, (2) multipolar division, (3) endocycling, (4) cell death, and (5) prolonged mitotic arrest. (Fig. 2A). We then subjectively divided these five outcomes into three groups. Group 1 is successful bipolar divisions, representing the failure of a drug to block mitosis at the surveyed doses. Group 2 includes prolonged mitotic arrest and cell death, which are desirable anticancer cell fates. Group 3 fates are hidden in growth inhibition assays and include multipolar divisions and endocycling. Group 3 fates can yield chromosomal aberrations and make tumor cells increasingly aggressive and difficult to eradicate.<sup>7-9</sup> Importantly, conventional multi-day growth inhibition counting assays combine Group 1, Group 2 mitotic arrest, and Group 3 outcomes to report on the cumulative number of live cancer cells at specific time points during and post-drug treatment without discriminating Group 3 unfavorable and potentially dangerous outcomes.

Criteria for classifying cell fates relies upon specific biophysical and morphological QPI measurements. A sharp surge in mean phase-shift above 140 nm and shape factor, a measurement of roundness, above 0.7 defines the time point of mitotic entry as cells 'round

up' (Fig. S4).<sup>25</sup> During a successful bipolar division, a single cell separates into two cells and then flattens to resume growth during interphase. The mitotic interval with cytokinesis is complete when two cells emerge at or near the former parent cell location with mean phase-shift and shape factor below 140 nm and 0.7 thresholds (Fig. 2A-B).<sup>25</sup> A cell death fate occurs when cell biomass shows either a sudden or a slow decline following mitotic entry (Fig. 2A, 2C).<sup>26</sup> During cell death the mean phase-shift through the cell decreases but stays above the 140nm threshold due to pyknosis, karyorrhexis, and cytoplasm content condensation.<sup>27</sup> The shape factor value also decreases when severe membrane blebbing or cell disintegration occurs (Fig. 2C). A prolonged mitotic arrest is identified by stagnation of changes in cell biomass, with mean phase-shift and shape factor remaining above 140nm and 0.7 thresholds, and no cell division or other morphological changes occurring (Fig. 2A, 2D). A multipolar cell division fate shows the same time-dependent tracing pattern in mean phase-shift, shape factor, and biomass accumulation as bipolar divisions with three or more cells arising at or near the previous parent cell location (Fig. 2A, 2E). Following multipolar divisions, not all daughter cells grow or thrive, likely due to aberrant chromosome partitioning.<sup>28</sup> Finally, QPI identifies endocycling when a cancer cell shows a sudden surge in mean phase-shift and shape factor, indicative of mitotic 'rounding', but returns to a G0/G1 flat morphology, mean phase-shift under 140nm, and shape factor less than 0.7 without cell division (Fig 2A, 2F).<sup>29</sup> In cases when single cells are hard to dissect from neighbors or are difficult to track over time, we apply manual screening corrections to the results of the cell fate algorithm.

### Calculating $EC_{50, \text{growth}}$ and $EC_{50, \text{death}}$ using QPI quantified cell fate outcomes

In growth inhibition assays, cell fate outcomes are binary, meaning that a drug does or does not alter live cell numbers compared to control conditions over time from the sum of cell divisions, arrests, and deaths. Thus, conventional preclinical drug development assays generate one  $EC_{50}$  curve to fit a binary outcome. By contrast, QPI yields three cell fate groups (Groups 1 – 3) with qualitatively distinct outcomes available for quantification. Therefore, instead of analyzing total live cell numbers, we generated two dose-response curves from QPI data. One dose-response curve corresponds to successful bipolar divisions from drug insensitivity (Fig. 3, solid line). The second dose-response curve describes desirable outcomes with prolonged mitotic arrest or cell death (Fig. 3, dotted line). We characterized these curves based on a standard log-scale normalized response equation:

$$f(x) = \frac{1}{(1 + 10^{\frac{x - EC_{50}}{EC_{50}}})} \quad (1)$$

We next compared calculated  $EC_{50}$  values between drug treatment panels (Table S1). In particular, the ability of QPI to identify multiple cancer cell fates enables the characterization of mitotic inhibitors for both sensitivity, represented by a low average  $EC_{50}$  value, as well as avoidance of undesirable fates. The concentration window in which undesirable fates occur is described by  $EC_{50} = EC_{50, \text{death}} - EC_{50, \text{growth}}$ . Therefore, QPI also enables the identification of mitotic inhibitors with a low  $EC_{50}$ , indicating a small concentration window that results in undesirable cell fates. When comparing  $EC_{50}$  between



inhibitors and cell lines, we used normalized value of  $EC_{50}/EC_{50,growth}$  because relative  $EC_{50}$  shows the impact of concentration variance relative to the target dose of a specific inhibitor (Table S1). Colchicine had the smallest relative  $EC_{50}$ , for both HeLa cells (0.35) and M202 cells (0.72) (Fig. 3B, 3E). This result is consistent with previous results showing that colchicine induces negligible or low amounts of aneuploidy.<sup>6</sup> Paclitaxel showed a slightly larger relative  $EC_{50}$  for both HeLa cells (1.1) and M202 cells (2.0), due to the increase in multipolar divisions and endocycling events compared to colchicine (Fig. 3A, 3D). VX-680 presented by far the largest relative  $EC_{50}$  for HeLa cells (110) and M202 cells (17), due to a large proportion of endocycling events (Fig. 3C, 3F). Lastly, Kolmogorov-Smirnov test results show that bipolar divisions and prolonged mitotic arrest or cell death dose response distributions statistically differ (Table S1), with the exception of M202 cells under increasing paclitaxel doses ( $p = 0.052$ ).

To confirm that a wide range of VX-680 dosing generates QPI detected endocycling, as predicted by a large  $EC_{50}$ , we analyzed the DNA content of HeLa and M202 cells exposed to VX-680 using flow cytometry (Fig. S5). As an example, there was a large increase in the number of cells with 4n DNA compared to those with 2n DNA in cells exposed to 300nM and 600nM VX-680. A small population of HeLa cells with 8n DNA content emerged at 300nM VX-680 that dramatically increased with 600nM VX-680 (Fig. S5C-F), in agreement with the features of endocycling.<sup>29</sup>

### Comparing $EC_{50,growth}$ and $EC_{50,death}$ to conventional $EC_{50}$ from live cell counting assays

In parallel, we performed live cell counting assays for HeLa and M202 cancer cells exposed to paclitaxel, colchicine, and VX-680 at 24 hours and multi-day timepoints (Fig. S1, S2). As anticipated, these agents markedly reduced the accumulation of cancer cells at 24 hours and over 6 days (Fig. S1, S2). Growth inhibition studies are the standard for drug development, although drug effects beyond changes in total live cell numbers were indiscernible. It is important to note that  $EC_{50}$  values from live cell counting assays are defined by concentrations at which the total number of live cells present in the treatment well is 50% of the total number of live cells present in the control well. Whereas,  $EC_{50}$  values from QPI analysis are defined by concentrations at which 50% of the sampled cells in a specific treatment undergo a certain cell fate within the first division after treatment. When compared to QPI measured  $EC_{50}$  values, live cell counting  $EC_{50}$  values are therefore on the same or nearest log scale, but greater than both  $EC_{50,Growth}$  and  $EC_{50,Death}$  for paclitaxel and colchicine treatments. However, in the case of VX-680, many endocycling events occur that do not lead to immediate cell deaths. This leads to large differences between  $EC_{50,Growth}$  and  $EC_{50,Death}$  values measured by QPI. The  $EC_{50}$  measured by live cell counting assay falls in the middle of the two on the log scale. This result therefore indicates that the QPI  $EC_{50,Growth}$  and  $EC_{50,Death}$  correspond differently to the standard live cell counting  $EC_{50}$  for different mitotic inhibition mechanisms.

We also assessed whether inhibitor dosage statistically associates with outcome by applying chi-square tests of independence for each cancer cell type and drug combination (Fig. 3). Our analyses show that within each of the six cancer cell-drug treatment pairings, cell fate outcomes were dose-dependent. Increasing dose shifts cell fate distributions from mostly

successful bipolar divisions to mostly mitotic arrests and deaths ( $p < 0.0001$  for each condition) (Table S2). For example, HeLa cells showed a range of outcomes at lower doses of paclitaxel and VX-680 (Fig. 3A, 3C). At 10 nM of paclitaxel, a dose between the two observed QPI EC<sub>50</sub> values, all outcomes except endocycling occurred, with only a small amount of endocycling appearing at slightly higher drug concentrations (Fig. 3A). By contrast, the majority of HeLa cells endocycle at all VX-680 concentrations within the first 24 hours of treatment, whereas prolonged mitotic arrest or cell death increased in outcome frequency only at 600nM VX-680. This concentration is 4–12 times higher than the recommended effective dose (50 – 150nM) *in vitro* against thyroid and blood-cancer cell lines (Fig. 3C),<sup>30, 31</sup> yet well within the range of mean plasma concentration of the maximum-tolerated dose (64 mg/m<sup>2</sup>-h) in patients, determined in phase I clinical trial.<sup>32</sup>

### Analyzing correlation between mitosis durations and cell fate outcomes

Finally, there has been debate over whether the duration of mitosis during anticancer drug exposure affects cancer cell outcomes.<sup>10, 33</sup> This data is quantifiable from QPI data showing the period from onset of mitosis to scored cell outcomes.<sup>3, 10</sup> Single HeLa cell tracking data show that time spent in mitosis statistically differ between cell fate outcomes ( $p = 1.8e-54$ ). Multipolar divisions, cell death, or arrest are frequent outcomes of extended mitosis time, whereas endocycling usually results from shorter periods of mitosis (Fig. 4). For M202 cells, even though the mean values of time spent in mitosis for cell fate outcomes are statistically different ( $p = 2.3e-4$ ), there is no significant trend for mitosis durations between outcomes categories (Fig. S6). Mitosis durations less than 100 minutes more frequently (45% for HeLa and 34% for M202) resolve as drug insensitivity with successful bipolar cell divisions compared to other fates in both cell lines (Fig. 4D, S6D). These data clearly reveal variability in responses to the same mitotic inhibitor treatments from different cancer cell lines. Thus, QPI-derived outcomes classifications reinforce the importance of studying responses in multiple cancer cell lines and types during preclinical drug development that may correspond to differences in personalized responses to different treatment agents and regimens for individuals with cancer.

## DISCUSSION

Changes in cell biomass as a response indicator for screening cancer drugs has gained traction in recent years because of multiple technological breakthroughs.<sup>19, 21, 22, 34-36</sup> A common thread in this emerging area is the increasing linkage between biomass accumulation rates with traditional measures of drug efficacy and preclinical outcomes of growth inhibitors in many types of modeled malignancies. QPI methods in particular are providing additional biophysical insights based on changes in cell morphology during studies of cell division and cancer-immune cell interactions.<sup>25, 37</sup> Here, we expanded the application spectrum for quantitative phase methods by combining biomass and morphological analyses in studies of the pharmacodynamic characteristics of small molecule mitotic inhibitors against two types of cancer.

We showed that during exposure to mitotic inhibitors QPI classification into three outcome Groups yields useful concentration windows in which undesirable fates that include

multipolar cell divisions and endocycling occur, as described by  $EC_{50}$ . Cancer cells exposed to inadequate mitotic inhibitor concentrations resolve transient cell cycle arrest by apoptosis or ‘mitotic slippage’ to resume growth and cycling<sup>38</sup>, which can result in chromosome abnormalities such as aneuploidy and tetraploidy.<sup>39</sup> Numerical and structural chromosome aberrations, which often occur with multipolar cell divisions and endocycling, may contribute to increased therapy resistance<sup>40</sup> through a range of molecular mechanisms.<sup>7, 8</sup> Aneuploidy paradoxically can promote or suppress tumor growth and may cause an elevated rate of tumor recurrence by generating drug-resistant heterogeneity with evolving growth and survival advantages.<sup>40, 41</sup> Tetraploid cells and cells with certain chromosomal rearrangements act as intermediates to further chromosome instability and the development of aneuploidy.<sup>7</sup> Unfortunately, tetraploidy and chromosomal aberrations that can contribute to therapy resistance have also been linked to mitotic inhibitor exposure at sub-lethal concentrations.<sup>6</sup> It has been shown that mitotic slippage occurs far more frequently *in vivo* than *in vitro*, suggesting possible underestimation of Group 3 outcomes with QPI compared to inefficacious agents and malignancy augmentation that may occur in patients.<sup>42</sup> Interestingly, proteotoxic and metabolic stress are reported for aneuploid cells,<sup>43</sup> which may result from multipolar divisions and endocycling events identified using our QPI methods.

Inhibitor mechanisms, concentrations, and mitotic arrest duration are major determinants of cancer cell outcomes following escape from mitotic arrest.<sup>33, 44, 45</sup> Our QPI approach analyzed these key characteristics for different mitotic inhibitors. Because of off-target cytotoxicity, mitotic inhibitors typically have low therapeutic indices that can limit clinical dosages<sup>43</sup> and cause abnormal exits with viable cells containing chromosomal abnormalities.<sup>6, 44</sup> *In vivo*, this undesirable outcome can occur in under-dosed or poorly perfused tumors and potentially jeopardizes clinical outcomes by adding to therapy resistance and tumor aggressiveness.

*In vitro* multi-day growth inhibition binary assays yield a single  $EC_{50}$  curve that overlooks a range of potential mitotic slippage events that add to chromosomal aberrations. Advances in non-QPI techniques such as *in vivo* microscopy and real time FUCCI imaging enable quantifying mitotic exits for a small number of cells.<sup>46, 47</sup> Additional methods, such as microchannel resonators, have limitations in measurement longevity and are low throughput. Recently, QPI methods were used to quantify cytotoxicity using area and morphological information.<sup>13, 14</sup> Although previous QPI applications showed results comparable to automatic cell profilers, the analysis did not fully leverage the wealth of biophysical information captured by QPI techniques. In contrast, the QPI analysis method presented here exposed differences in outcomes between a microtubule-destabilizing agent, paclitaxel, a microtubule-stabilizing agent, colchicine, and a cell division inhibitor, VX-680. Our decision tree algorithm provided five cancer cell outcome categories and three subjective Groups to enable determination of dual  $EC_{50}$  curves for each drug and tumor cell type examined. For a particular drug and tumor type, the extent of  $EC_{50}$  value separation,  $EC_{50}$ , predicts the likelihood of undesirable multipolar divisions or endocycling that can result in aneuploidy and increased therapy resistance, providing useful concentration ranges to avoid in which aberrant mitotic exits prevail.

We note that there is a difference between the dual  $EC_{50}$ s determined via QPI and the single  $EC_{50}$  determined via live cell counting, especially for paclitaxel and colchicine. Paclitaxel and colchicine induced large numbers of mitotic arrests in the treated cell populations. While live cell counting classifies arrested cells as live cells, the QPI data analysis method described here does not account for arrested cells in  $EC_{50 \text{ growth}}$ . Criteria used for a cell to be included in  $EC_{50 \text{ growth}}$  requires that cell to successfully divide during the imaging period. Therefore, live cell counting identified a higher proportion of cells as live cells than QPI analysis did as growing cells, producing a larger  $EC_{50}$  value than  $EC_{50 \text{ growth}}$  for paclitaxel and colchicine treatments.

QPI analyses revealed that HeLa and M202 bipolar divisions from drug failure show the shortest mitosis periods, consistent with previous studies in which prolonged mitotic arrest resulted in hypersensitivity to additional death cues.<sup>33</sup> Additional studies showed several existing chemotherapeutics that induce apoptosis at normal dosages can trigger mitotic catastrophe that directly lead to apoptosis at very low dosages in aneuploid and polyploid cells.<sup>48</sup> This can dramatically increase chemotherapeutic tolerance in patients and hint at effective combinatorial therapies in cancer treatments using mitotic inhibitors. Therefore, future QPI studies include screens for cell fate outcomes under combined exposure of energy stressors or low dose chemotherapeutics and mitotic inhibitors that induce aneuploidy.

## CONCLUSIONS

In this study, we demonstrate a novel application of QPI in screening and identifying aberrant cell fate outcomes as a result of suboptimal mitotic inhibitor doses. Conventional growth inhibition assays rely on live cell counting to generate  $EC_{50}$  values that infer pharmacokinetics of mitotic inhibitors, and fail to reveal vital information on negative effects of the inhibitors. QPI analyzes cell fate outcome profiles,  $EC_{50, \text{Growth}}$ , and  $EC_{50, \text{Death}}$  values that provide in-depth insights into mechanism of action and risky dosing windows of mitotic inhibitors. Importantly, this QPI technique is compatible with patient derived organoids that resemble heterogeneous patient tumors better than *in vitro* single cell type cultures. Screening mitotic inhibitor cocktails on tumor organoids with QPI can facilitate the development of next generation cancer therapies.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## ACKNOWLEDGEMENTS

D.H. conducted the experiments, analyzed the data, prepared the figures and wrote the manuscript. I.J.R. helped perform experiments and analyzed data under supervision of D.H. G.M. and J.R. analyzed results and edited the manuscript. T.A.Z. supervised the study along with M.A.T., who initiated, supervised and obtained funding for the study. We thank Fasih Ahsan for advice on statistical analysis. Funded by National Institutes of Health grant R21CA227480 to M.A.T. and R01CA185189 to M.A.T. and J.R., and in part, by NCI Cancer Center Support Grant P30CA016042 to the UCLA Jonsson Comprehensive Cancer Center and P30CA016059 to the VCU Massey Cancer Center.

## REFERENCES

1. Shtivelman E, Davies MQ, Hwu P, Yang J, Lotem M, Oren M, Flaherty KT and Fisher DE, *Oncotarget*, 2014, 5, 1701–1752. [PubMed: 24743024]
2. Shi H, Hugo W, Kong X, Hong A, Koya RC, Moriceau G, Chodon T, Guo R, Johnson DB, Dahlman KB, Kelley MC, Kefford RF, Chmielowski B, Glaspy JA, Sosman JA, van Baren N, Long GV, Ribas A and Lo RS, *Cancer Discov*, 2014, 4, 80–93. [PubMed: 24265155]
3. Weaver BA and Cleveland DW, *Cancer Cell*, 2005, 8, 7–12. [PubMed: 16023594]
4. Manchado E, Guillaumot M and Malumbres M, *Cell Death Differ*, 2012, 19, 369–377. [PubMed: 22223105]
5. Brito DA and Rieder CL, *Curr Biol*, 2006, 16, 1194–1200. [PubMed: 16782009]
6. Chen JG and Horwitz SB, *Cancer Res*, 2002, 62, 1935–1938. [PubMed: 11929805]
7. Fujiwara T, Bandi M, Nitta M, Ivanova EV, Bronson RT and Pellman D, *Nature*, 2005, 437, 1043–1047. [PubMed: 16222300]
8. Ganem NJ, Storchova Z and Pellman D, *Curr Opin Genet Dev*, 2007, 17, 157–162. [PubMed: 17324569]
9. Schvartzman JM, Sotillo R and Benezra R, *Nat Rev Cancer*, 2010, 10, 102–115. [PubMed: 20094045]
10. Gascoigne KE and Taylor SS, *Cancer Cell*, 2008, 14, 111–122. [PubMed: 18656424]
11. Park Y, Depeursinge C and Popescu G, *Nature Photonics*, 2018, 12, 578–589.
12. Marquet P, Depeursinge C and Magistretti P, *Neurophotonics*, 2014, 1.
13. Greve B, Sheikh-Mounessi F, Kemper B, Ernst I, Gotte M and Eich H, *Strahlentherapie Und Onkologie*, 2012, 188, 1038–1047. [PubMed: 23053158]
14. Kühn J, Shaffer E, Mena J, Breton B, Parent J, Rappaz B, Chambon M, Emery Y, Magistretti P, Depeursinge C, Marquet P and Turcatti G, *Assay Drug Dev Technol*, 2013, 11, 101–107. [PubMed: 23062077]
15. Bon P, Maucort G, Wattellier B and Monneret S, *Opt Express*, 2009, 17, 13080–13094. [PubMed: 19654713]
16. Aknoun S, Savatier J, Bon P, Galland F, Abdeladim L, Wattellier B and Monneret S, *Journal of Biomedical Optics*, 2015, 20.
17. Reed J, Chun J, Zangle TA, Kalim S, Hong JS, Pefley SE, Zheng X, Gimzewski JK and Teitell MA, *Biophys J*, 2011, 101, 1025–1031. [PubMed: 21889438]
18. Zangle TA, Chun J, Zhang J, Reed J and Teitell MA, *Biophys J*, 2013, 105, 593–601. [PubMed: 23931307]
19. Chun J, Zangle TA, Kolarova T, Finn RS, Teitell MA and Reed J, *Analyst*, 2012, 137, 5495–5498. [PubMed: 23057068]
20. O'Brien NA, Browne BC, Chow L, Wang Y, Ginther C, Arboleda J, Duffy MJ, Crown J, O'Donovan N and Slamon DJ, *Mol Cancer Ther*, 2010, 9, 1489–1502. [PubMed: 20501798]
21. Huang D, Leslie KA, Guest D, Yeshcheulova O, Roy IJ, Piva M, Moriceau G, Zangle TA, Lo RS, Teitell MA and Reed J, *Anal Chem*, 2018, 90, 3299–3306. [PubMed: 29381859]
22. Murray GF, Turner TH, Leslie KA, Alzubi MA, Guest D, Sohal SS, Teitell MA, Harrell JC and Reed J, *ACS Omega*, 2018, 3, 17687–17692. [PubMed: 30613814]
23. Barer R, *Nature*, 1952, 169, 366–367. [PubMed: 14919571]
24. Ross KFA, *Phase contrast and interference microscopy for cell biologists*, St. Martin's Press, New York., 1967.
25. Zangle TA, Teitell MA and Reed J, *PLoS One*, 2014, 9, e115726. [PubMed: 25531652]
26. Chowdhury I, Tharakan B and Bhat GK, *Comp Biochem Physiol B Biochem Mol Biol*, 2008, 151, 10–27. [PubMed: 18602321]
27. Cotran RS, Kumar V, Collins T and Robbins SL, *Robbins pathologic basis of disease*, Saunders, Philadelphia, 6th edn., 1999.
28. Ganem NJ, Godinho SA and Pellman D, *Nature*, 2009, 460, 278–282. [PubMed: 19506557]

29. Edgar BA, Zielke N and Gutierrez C, *Nat Rev Mol Cell Biol*, 2014, 15, 197–210. [PubMed: 24556841]
30. Arlot-Bonnemains Y, Baldini E, Martin B, Delcros JG, Toller M, Curcio F, Ambesi-Impiombato FS, D'Armiento M and Ulisse S, *Endocr Relat Cancer*, 2008, 15, 559–568. [PubMed: 18430894]
31. Huang XF, Luo SK, Xu J, Li J, Xu DR, Wang LH, Yan M, Wang XR, Wan XB, Zheng FM, Zeng YX and Liu Q, *Blood*, 2008, 111, 2854–2865. [PubMed: 18160664]
32. Traynor AM, Hewitt M, Liu G, Flaherty KT, Clark J, Freedman SJ, Scott BB, Leighton AM, Watson PA, Zhao B, O'Dwyer PJ and Wilding G, *Cancer Chemother Pharmacol*, 2011, 67, 305–314. [PubMed: 20386909]
33. Bekier ME, Fischbach R, Lee J and Taylor WR, *Mol Cancer Ther*, 2009, 8, 1646–1654. [PubMed: 19509263]
34. Cetin AE, Stevens MM, Calistri NL, Fulciniti M, Olcum S, Kimmerling RJ, Munshi NC and Manalis SR, *Nat Commun*, 2017, 8, 1613. [PubMed: 29151572]
35. Li Y, Fanous MJ, Kilian KA and Popescu G, *Sci Rep*, 2019, 9, 248. [PubMed: 30670739]
36. Kang JH, Miettinen TP, Chen L, Olcum S, Katsikis G, Doyle PS and Manalis SR, *Nat Methods*, 2019, 16, 263–269. [PubMed: 30742041]
37. Zangle TA, Burnes D, Mathis C, Witte ON and Teitell MA, *PLoS One*, 2013, 8, e68916. [PubMed: 23935904]
38. Sakurikar N, Eichhorn JM and Chambers TC, *J Biol Chem*, 2012, 287, 39193–39204. [PubMed: 22965228]
39. Storchova Z and Kuffer C, *J Cell Sci*, 2008, 121, 3859–3866. [PubMed: 19020304]
40. Weaver BA and Cleveland DW, *J Cell Biol*, 2009, 185, 935–937. [PubMed: 19528293]
41. Sotillo R, Schwartzman JM, Socci ND and Benezra R, *Nature*, 2010, 464, 436–440. [PubMed: 20173739]
42. Orth JD, Kohler RH, Foiijer F, Sorger PK, Weissleder R and Mitchison TJ, *Cancer Res*, 2011, 71, 4608–4616. [PubMed: 21712408]
43. Dominguez-Brauer C, Thu KL, Mason JM, Blaser H, Bray MR and Mak TW, *Mol Cell*, 2015, 60, 524–536. [PubMed: 26590712]
44. Jordan MA, Wendell K, Gardiner S, Derry WB, Copp H and Wilson L, *Cancer Res*, 1996, 56, 816–825. [PubMed: 8631019]
45. Torres K and Horwitz SB, *Cancer Res*, 1998, 58, 3620–3626. [PubMed: 9721870]
46. Miwa S, Yano S, Kimura H, Yamamoto M, Toneri M, Matsumoto Y, Uehara F, Hiroshima Y, Murakami T, Hayashi K, Yamamoto N, Bouvet M, Fujiwara T, Tsuchiya H and Hoffman RM, *Cell Cycle*, 2015, 14, 621–629. [PubMed: 25551170]
47. Orth JD, Tang Y, Shi J, Loy CT, Amendt C, Wilm C, Zenke FT and Mitchison TJ, *Mol Cancer Ther*, 2008, 7, 3480–3489. [PubMed: 18974392]
48. Denisenko TV, Sorokina IV, Gogvadze V and Zhivotovsky B, *Drug Resist Updat*, 2016, 24, 1–12. [PubMed: 26830311]



**Fig. 1.** Experimental design and analysis schematics. (A) Timeline of the experiment, showing cell synchronization and period of QPI. (B) Example QPI images of single HeLa cells under 50nM of colchicine treatment. The same cells outlined in green and purple were tracked continuously over time to determine their post-mitotic entry fates, in this case are death and prolonged mitotic arrest. Color bar indicates phase shift in nm. Time stamps indicate time since start of drug exposure. (C) Representative schematic of biomass, mean phase-shift and shape factor data over time for two cell fates shown in (B): death and mitotic arrest. (D) Representative schematic of two  $EC_{50}$  curves generated based on normalized cell-fate outcomes distributions from randomly sampled single cells over a range of drug

concentrations. The difference between the two  $EC_{50}$  values,  $EC_{50}$ , provides a dosing range in which the drug can potentially cause aberrant mitotic exits.


Author Manuscript

Author Manuscript


Author Manuscript

Author Manuscript






**Fig. 2.** Cell fate algorithm with examples. (A) A decision tree diagram showing the MATLAB algorithm for determining cell fates based on dynamic changes in cell biomass, mean phase-shift, and shape factor measured by QPI. (B) Example of a HeLa cell undergoing a bipolar division. (C) Example of a HeLa cell dying. (D) Example of a HeLa cell in prolonged mitotic arrest. (E) Example of a HeLa cell undergoing a multipolar division. (F) Example of a HeLa cell undergoing endocycling.



**Fig. 3.** Cell fate distribution analysis. (A) Cell fate distribution of HeLa cells with paclitaxel exposure ( $n = 187$ ). (B) Cell fate distribution of HeLa cells with colchicine exposure ( $n = 380$ ). (C) Cell fate distribution of HeLa cells with VX-680 exposure ( $n = 223$ ). (D) Cell fate distribution of M202 cells with paclitaxel exposure ( $n = 128$ ). (E) Cell fate distribution of M202 cells with colchicine exposure ( $n = 42$ ). (F) Cell fate distribution of M202 cells with VX-680 exposure ( $n = 63$ ). A solid line in each panel represents a dose response curve fit to the bipolar division distribution. A dotted line in each panel represents a dose response curve fit to the prolonged mitotic arrest or cell death distribution. Numbers represent  $EC_{50}$  values of for each curve. Legend shows color code for cell fate outcomes.



**Fig. 4.** Cell fate responses and mitosis durations. (A) Time spent in mitosis for HeLa single cells with colchicine exposure. The length of each color bar indicates the amount of time spent in mitosis for that individual cell. The color of the bar corresponds to its cell fate outcome. (B) Time spent in mitosis for HeLa single cells with palitaxel exposure. (C) Time spent in mitosis for HeLa single cells with VX-680 exposure. (D) Box plot of time spent in mitosis for five cell fate outcomes. One-way ANOVA with unbalanced sample groups was performed comparing all samples. There are statistically significant differences between the mean values of mitosis durations for different cell fate outcomes ( $p = 1.8E-54$ ).

**Table 1**

Mitotic Inhibitors Drug Concentrations Administered to M202 and HeLa Cell Lines in Live Cell Interferometer Study.

Cell Lines	Paclitaxel (nM)	Colchicine (nM)	VX-680 (nM)
HeLa	10, 50, 100, 500	50, 150, 500, 1500, 2500	100, 300, 600
M202	1, 3, 9, 50, 100, 500	1, 5, 10, 20, 50	30, 90, 150, 300, 600

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript