UC Riverside

UC Riverside Previously Published Works

Title

Synthesis of the C₅₉N⁺ carbocation. A monomeric azafullerene isoelectronic to C₆₀.

Permalink

<https://escholarship.org/uc/item/9zm1w8x3>

Journal

Journal of the American Chemical Society, 125(14)

ISSN

0002-7863

Authors

Kim, Kee-Chan
Hauke, Frank
Hirsch, Andreas
et al.

Publication Date

2003-04-01

DOI

10.1021/ja034014r

Peer reviewed

Synthesis of the $C_{59}N^+$ Carbocation. A Monomeric Azafullerene Isoelectronic to C_{60}

Kee-Chan Kim,[†] Frank Hauke,[‡] Andreas Hirsch,[‡] Peter D. W. Boyd,[§] Elizabeth Carter,[⊥]
Robert S. Armstrong,[⊥] Peter A. Lay,[⊥] and Christopher A. Reed^{*,†}

Department of Chemistry, University of California, Riverside, California 92521-0403, Institut für Organische Chemie der Universität Erlangen-Nürnberg, Henkestrasse 42, 91054 Erlangen, Germany, Department of Chemistry, The University of Auckland, Private Bag, Auckland, New Zealand, 92521-0403, and School of Chemistry, University of Sydney, NSW 2006, Australia

Received January 2, 2003; E-mail: chris.reed@ucr.edu

By isoelectronic analogy to C_{60} , heterofullerenes such as $C_{59}N^+$, $C_{59}B^-$, $C_{59}Si$, and $C_{59}P^+$ should be stable entities. Differences in size and electronegativity between C and its neighboring nonmetallic elements are unlikely to be large enough to disrupt covalent bonding in the icosahedral cage. Indeed, ions having these elemental compositions have been relatively easy to detect via mass spectrometry on fullerene soots prepared in the presence of heteroatom sources.^{1–4} Nevertheless, only one analogue of C_{60} has been carried to the level of practical synthesis and compositional purity, the *N*-heterofullerene $C_{59}N$. With one more electron than C_{60} , its stable form is a C–C bonded dimer, $(C_{59}N)_2$.^{5–8} The C–H bonded monomer, $HC_{59}N$, has also been characterized.⁹ We now report that the monomeric $C_{59}N^+$ cation, which is isoelectronic with C_{60} , can be isolated as a carborane anion salt. The synthesis involves a rare example of oxidation of an sp^3 – sp^3 C–C bond to produce a carbenium ion.

Fullerenes are typically easy to reduce but hard to oxidize¹⁰ and $(C_{59}N)_2$ is no exception. A threshold oxidation potential of ca. +0.9 V (vs Fc/Fc⁺) has been reported in its irreversible anodic cyclic voltammetry.⁹ In addition, $(C_{59}N)_2$ has an sp^3 – sp^3 C–C bond and such bonds typically present high barriers to oxidation. Chemistry has few “electron–hole” oxidants that operate above ca. +0.7 V¹¹ because most strong oxidants (e.g., Cl_2 , SbF_5 , XeF_2) come partnered with nucleophiles (e.g., halides) that immediately react with oxidized species. However, when partnered with exceptionally inert carborane anions^{12,13} such as $CB_{11}H_6Cl_6^-$, stable radical cations can be used to extend the range to +1.34 V, sufficient to oxidize C_{60} to the $C_{60}^{+\bullet}$ radical cation.¹⁴

The radical cation of crude hexabromo(phenyl)carbazole (HBPC^{•+})^{14,15} oxidizes $(C_{59}N)_2$ ⁶ to $C_{59}N^+$ in dry *o*-dichlorobenzene (eq 1). The counterion is the silver(I) bis-carborane complex ion

$[Ag(CB_{11}H_6Cl_6)_2]^-$. $[C_{59}N][Ag(CB_{11}H_6Cl_6)_2]$, **1**, was isolated as a brown precipitate in good yield (>75%) by addition of hexane or crystallized as dark green crystals by diffusion of hexane vapor.¹⁵ Anal. Calcd for $C_{61}H_{12}NB_{22}Cl_{12}Ag$: C, 47.89; H, 0.79; N, 0.92. Found: C, 47.49; H, 0.92; N, 1.10. The solid is reasonably air stable.

The dark green color of $(C_{59}N)_2$ in *o*-dichlorobenzene lightens slightly upon oxidation. The visible spectrum of $C_{59}N^+$ is quite featureless (blue line in Figure 1) compared to that of $(C_{59}N)_2$.

Figure 1. Visible spectra of $C_{59}N^+$ (blue, 1.38×10^{-3} M) and $(C_{59}N)_2$ (red, 6.90×10^{-4} M) in *o*-dichlorobenzene.

Stirring solutions of the $C_{59}N^+$ salt over Hg-amalgamated zinc pellets slowly returned the spectrum to that of $(C_{59}N)_2$, indicating reversibility of the redox chemistry.

The infrared spectrum of the $C_{59}N^+$ ion can be distinguished from that of $(C_{59}N)_2$ and $HC_{59}N$ in the 1600–1150 and 600–500 cm^{-1} regions. Bands at 1586, 1580, 1575; 1468, 1455, 1434; ~1183 sh, 1178, 1165; 579, 537, 532, 527, 525, 519 cm^{-1} in **1**¹⁵ contrast with those at 1574, 1565; 1460, 1443, 1423, 1416; 1196, 1186, 1174; 579, 576, 568, 529, 523 cm^{-1} in $(C_{59}N)_2$ and 1574, 1565, 1461, 1443, 1422, 1414; 1197, 1186, 1174; 579, 574, 568, 529, 523 cm^{-1} in $HC_{59}N$.¹⁶ Other fingerprint bands for the $C_{59}N$ moiety near 800 cm^{-1} are masked by bands from the carborane anion. In the resonance Raman spectrum of **1** with 488 nm excitation, the prominent $A_g(2)$ global breathing mode appears at 1467 cm^{-1} .¹⁵ This is higher in energy than that of $(C_{59}N)_2$ (1462 cm^{-1}),¹⁶ consistent with a tightening of cage bonding from the sp^3 to sp^2 bonding change at C. The frequency in $C_{59}N^+$ is essentially the same as that in C_{60} (1466 cm^{-1}). More bands appear in $C_{59}N^+$ because of lower symmetry.

Consistent with a positive ion already being present, MALDI mass spectroscopy of **1** in positive ion mode gave a dominant peak at $m/z = 722$ for the $C_{59}N^+$ cation using low laser powers.¹⁵ Under comparable conditions, $(C_{59}N)_2$ gave no signal. The isotope pattern confirms the formulation as $C_{59}N^+$.¹⁷ In negative ion mode, a broad, strong peak centered at $m/z = 350$ identifies the carborane anion $CB_{11}H_6Cl_6^-$, indicating expected dissociation of the weakly bound $Ag(CB_{11}H_6Cl_6)_2^-$ complex anion.

As expected from C_s symmetry, the ^{13}C NMR spectrum of **1** in tetrachloroethane-*d*₂ shows 31 peaks that can be ascribed to the $C_{59}N^+$ ion (Figure 2). A near accidental degeneracy occurs at 142.7/142.8 ppm. Because solubility is higher, a spectrum with a better signal-to-noise ratio could be obtained in *o*-dichlorobenzene-*d*₂ but slight solvent shifts in the resonances exacerbate the problem of accidental degeneracies.¹⁵ Peak positions distinguish $C_{59}N^+$ from $(C_{59}N)_2$ and $HC_{59}N$. Consistent with the overall positive charge on

[†] University of California, Riverside.

[‡] Universität Erlangen-Nürnberg.

[§] The University of Auckland.

[⊥] University of Sydney.

Figure 2. Experimental ^{13}C NMR spectrum of C_{59}N^+ in **1** in $\text{TCE-}d_2$ (above) and calculated spectrum (stick diagram below).

Figure 3. Selected ^{13}C assignments (ppm) in C_{59}N^+ and resonance forms showing both carbenium and iminium ion character.

Figure 4. Perspective view of $[\text{C}_{59}\text{N}^+][\text{Ag}(\text{CB}_{11}\text{H}_6\text{Cl}_6)_2^-]$ in **1**. Three lattice *o*-dichlorobenzene molecules are omitted for clarity.

the ion, the majority of the peaks cluster around 144 ppm, somewhat downfield of 142.9 ppm in C_{60} . Individual assignments can be made on the basis of the reasonable agreement with the DFT-calculated spectrum (B3LYP/6-31G*/B3LYP/6-311G(d,p)), shown as a stick diagram in Figure 2. A downfield peak at 153.8 ppm (1C intensity) is ascribed to the formal carbenium ion center (atom 60 in Figure 3). This compares to 182 ppm in the isoelectronic HC_{60}^+ ion,¹⁴ indicating greater delocalization of the positive charge in C_{59}N^+ and iminium ion character, favored by NBO analysis¹⁵ (Figure 3). The most upfield shifted resonances (133.7, 135.0, 137.4 ppm) arise from C atoms β to the N atom (C46, C56, C57, respectively).

The X-ray crystal structure of **1** with three *o*-dichlorobenzene solvate molecules provides final proof of formulation and structure (Figure 4). Although the anion is well ordered, disorder in the cation and the solvate molecules prevents accurate bond length data from being obtained for C_{59}N^+ . The cation is located on a crystallographic mirror plane that does not bisect the [6,6] C–N bond. It was successfully modeled with 50:50 site occupancies and the assump-

tion that the closest anion/cation contact identifies the carbocationic C atom (C4F in Figure 4).

In conclusion, an azafullerene analogue of C_{60} is accessible via chemical oxidation of dimeric $(\text{C}_{59}\text{N})_2$ with a strong electron–hole oxidant. As with reactive carbocations such as the benzenium ion (C_6H_7^+)¹⁸ and the radical cation $\text{C}_{60}^{\bullet+}$,¹⁴ the stabilization of C_{59}N^+ as an isolable salt profited from use of an inert carborane as counterion. The properties of C_{59}N^+ reflect the isoelectronic analogy to C_{60} and some delocalization of charge.

Acknowledgment. We thank Drs. Robert Bolskar, Fook Tham, and Alex M. Clark for very valuable experimental assistance. This work was supported by the NSF (CHE 00952 to C.A.R.), NIH (GM 23851 to C.A.R.), Deutsche Forschungsgemeinschaft to A.H.), Australian Research Council (Large and IREX to R.S.A. and P.A.L.), and Marsden Fund of the Royal Society of New Zealand (00-UOA-030 to P.D.W.B.).

Supporting Information Available: Full experimental details, NMR, vis, IR, Raman, and MALDI spectra, and X-ray crystallographic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Guo, T.; Jin, C.; Smalley, R. E. *J. Phys. Chem.* **1991**, *95*, 4948–4950.
- (2) Pradeep, T.; Vijayakrishnan, V.; Santa, A. K.; Rao, C. N. R. *J. Phys. Chem.* **1991**, *95*, 10564–10565.
- (3) Kimura, T.; Sugai, T.; Shinohara, H. *Chem. Phys. Lett.* **1996**, *256*, 269.
- (4) Moschel, C.; Jansen, M. Z. *Anorg. Allg. Chem.* **1999**, *625*, 175.
- (5) Hummelen, J. C.; Knight, B.; Pavlovich, J.; Gonzalez, R.; Wudl, F. *Science* **1995**, *269*, 1554–1556.
- (6) Nuber, B.; Hirsch, A. *Chem. Commun.* **1996**, 1421–1422.
- (7) Hummelen, J. C.; Bellavia-Lund, C.; Wudl, F. *Top. Curr. Chem.* **1999**, *199*, 93–134.
- (8) Hirsch, A.; Nuber, B. *Acc. Chem. Res.* **1999**, *32*, 795–804.
- (9) Keshavarz-K. M.; Gonzalez, R.; Hicks, R. G.; Srdanov, G.; Srdanov, V. I.; Collins, T. G.; Hummelen, J. C.; Bellavia-Lund, C.; Pavlovich, J.; Wudl, F.; Holczer, K. *Nature* **1996**, *383*, 147–150.
- (10) Reed, C. A.; Bolskar, R. D. *Chem. Rev.* **2000**, *100*, 1075–1119.
- (11) Connelly, N. G.; Geiger, W. E. *Chem. Rev.* **1996**, *96*, 877–910.
- (12) Reed, C. A. *Acc. Chem. Res.* **1998**, *31*, 133–139.
- (13) King, B. T.; Janousek, Z.; Gruner, B.; Trammell, M.; Noll, B. C.; Michl, J. *J. Am. Chem. Soc.* **1996**, *118*, 3313–3314.
- (14) Reed, C. A.; Kim, K.-C.; Bolskar, R. D.; Mueller, L. J. *Science* **2000**, *122*, 4660–4667.
- (15) See Supporting Information for details.
- (16) Krause, M.; Baes-Fischlmair, S.; Pfeiffer, R.; Plank, W.; Pichler, T.; Kuzmany, H.; Tagmatarchis, N.; Prassides, K. *J. Phys. Chem. B* **2001**, *105*, 11964–11969.
- (17) Lamparth, I.; Nuber, B.; Schick, G.; Skiebe, A.; Grosser, T.; Hirsch, A. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 2257–2259.
- (18) Reed, C. A.; Kim, K.-C.; Stoyanov, E. S.; Stasko, D.; Tham, F. S.; Mueller, L. J.; Boyd, P. D. W. *J. Am. Chem. Soc.* **2003**, *125*, 1796–1804.

JA034014R