

Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

Spallation-Fission Competition in Heaviest Elements; Triton Production

Permalink

<https://escholarship.org/uc/item/30j1r8mm>

Authors

Wade, William H.
Gonzalez-Vidal, Jose
Glass, Richard A.
et al.

Publication Date

1957-03-01

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-3640

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W 7405 eng 48

SPALLATION-FISSION COMPETITION IN HEAVIEST ELEMENTS;

TRITON PRODUCTION

William H. Wade, Jose Gonzalez-Vidal, Richard A. Glass,* and Glenn T. Seaborg

March, 1957

Printed for the U. S. Atomic Energy Commission

SPALLATION-FISSION COMPETITION IN HEAVIEST ELEMENTS;
TRITON PRODUCTION

Contents

Abstract	3
Introduction	4
Experimental	5
Results	
The (α ,t) reaction	6
The (d,t) and (p,t) reactions	6
Discussion	7
References	11

SPALLATION-FISSION COMPETITION IN HEAVIEST ELEMENTS:
TRITON PRODUCTION

William H. Wade, Jose Gonzalez-Vidal, Richard A. Glass,* and Glenn T. Seaborg

Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

March, 1957

ABSTRACT

Stacked foils of Au¹⁹⁷, Th²³², and U²³⁸ were bombarded, in a series of experiments, with 48-Mev helium ions, 24-Mev deuterons, and 32-Mev protons. Tritium from each foil was collected and then measured by a gas-counting technique. The qualitative results indicate that high-energy tritons are emitted in relatively large abundance from all targets and with each of the bombarding particles. Cross sections for triton production from fissionable U²³⁸ and Th²³² and from non-fissionable Au¹⁹⁷ are comparable. The integrated (α, t) cross sections thus determined for U²³⁸ and Th²³² are nearly the same as the integrated cross sections for the " $(\alpha, p2n)$ " reactions as determined by measuring the yield of the product heavy isotopes in radiochemical experiments. All of the facts are consistent with a picture of emission of high-energy tritons (whether due to a stripping, pick-up, or other mechanism) in which fissionable intermediate nuclei are formed mainly at levels of excitation below their fission thresholds.

* Present address, Stanford Research Institute, Palo Alto, California.

SPALLATION-FISSION COMPETITION IN HEAVIEST ELEMENTS:
TRITON PRODUCTION

William H. Wade, Jose Gonzalez-Vidal, Richard A. Glass, and Glenn T. Seaborg
Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

March 1957

I. INTRODUCTION

Considerable information about nuclear reaction mechanisms can be obtained both from radiochemical yield studies of residual nuclei and from investigations of outgoing neutrons, protons, and other small particles. Initial studies of the present series¹⁻⁴ dealt with radiochemical yields while the present study is the first to measure the yield of an outgoing particle.

In the first paper of the series¹ radiochemically determined yields were reported for spallation products from helium-ion-induced reactions in plutonium isotopes. These yields were characterized by the similar magnitude of the (α, xn) and (α, pxn) type cross sections. The more recent radiochemical studies²⁻⁴ of heavy-element nuclear reactions induced by helium ions of energies less than 50 Mev also have shown that certain cross sections, particularly that for the production of the $(\alpha, p2n)$ product, are too large relative to those for the production of the (α, xn) product to be explained on the basis of a compound-nucleus, evaporation picture in which charged-particle emission would be discriminated against. It has been observed that for U^{238} ,² U^{235} ,³ U^{233} ,³ and Th^{232} ,⁴ the $(\alpha, p2n)$ reaction is, in general, the most prominent of all the (α, xn) and (α, pxn) reactions observed. These considerations led to the hypothesis that the large values of the $(\alpha, p2n)$ cross sections in the region of fissionable nuclei are due to direct interaction of helium ions and target nuclei in which are produced tritons of sufficient energy to often leave the residual nuclei in states below fission and particle-emission thresholds. If this is true, it is reasonable to expect comparable $(\alpha, p2n)$ type cross sections for the heavy fissionable and the somewhat lighter non-fissionable nuclides. The following work was undertaken to test this hypothesis, and includes work on the (d, t) and (p, t) reactions.

Only a few experiments on tritium emission have previously been described. Tritons from the α and d and p, t reactions in bombardments with pro-

jectile energies greater than 100 Mev have been observed.^{5,6,7} In experiments with 600- and 2000-Mev protons Currie, Libby, and Wolfgang⁸ collected tritium from a number of elements and measured its radioactivity. With lower-energy particles (< 25 Mev), tritium production from deuteron and proton bombardments has been the subject of experimental investigation in both the light-element^{9,10} and heavy but non-fissionable element¹¹ regions. The stripping theory of Butler¹² has been applied by News¹³ by considering the (d,t) process as the inverse of stripping, i.e., neutron pick-up.

II. EXPERIMENTAL

For the helium-ion and deuteron bombardments, stacked .05-mm foils of natural Au¹⁹⁷ and isotopically pure U²³⁸, and 0.1-mm foils of natural Th²³² were subjected to the external beams of the 60-inch Crocker Laboratory cyclotron. The proton bombardments were carried out on the Berkeley linear accelerator using the same technique. In a later paper it will be shown that tritium production by these three bombarding particles is quite general over the entire periodic table. For this reason it was impossible to vary the beam energy by imposing degrading foils in front of the target without introducing an extraneous source of tritons. Thus, only maximum-energy beams were available to the first foil in the stack (48-Mev helium ions, 24-Mev deuterons, and 32-Mev protons).

The tritium was extracted by heating the foils to 1200°C in a measured amount of hydrogen carrier followed by selective diffusion of hydrogen isotopes through a palladium thimble and introduction into a counter tube. This apparatus is essentially the same as used by Currie, Libby, and Wolfgang, who discussed the possible experimental errors inherent in the apparatus. Counting was done in the proportional region using a methane filling of 1 atmosphere and a partial pressure of hydrogen of ~0.04 atms. Counting rates varied from 2×10^2 to 1×10^5 c/m, representing a total yield of tritium from the target into the counter tube of 35 to 40%.

III. RESULTS

A. The (α ,t) reaction.

The apparent cross-sections (σ) in mb for the (α ,t) reaction calculated on the basis of thin-target approximations are shown in Figure 1-A as a function of the mg/cm^2 of gold, thorium, and uranium as measured from the end of the stack upon which the beam was incident. The number of tritium atoms found in a foil, the number of particles incident on the stack, and the atoms per cm^2 in the foil were employed in each calculation. Since the helium-ion beam ranges are 460, 485, and 500 mg/cm^2 of gold, thorium, and uranium respectively, the maximum cross section consistently seems to be observed at depths in the target beyond the range of the helium ions. The integrated (α ,t) cross sections determined for gold, thorium, and uranium are listed in Table I in units of millibarn-mm and tritons per incident helium ion.* The cross sections are compared with cross sections for the production of the corresponding residual nuclei determined by the radiochemical method in the cases of thorium⁴ and uranium². It will be noted that tritium-emission and residual-nuclei cross sections agree within 20% in both cases. Table I also shows that the integrated (α ,t) cross section for gold agrees within 20% with the radiochemical yield values for the corresponding reaction for uranium and thorium.

B. The (d,t) and (p,t) reactions

Figures 1B and 1C show plots for the yields of tritium in the (d,t) and (p,t) reactions. In these cases no radiochemical cross-section determinations are available for comparison. The deuteron beam is completely stopped by 930, 995, and 1000 mg/cm^2 of gold, thorium, and uranium respectively, and the proton beam by 2450, 2580, and 2660 mg/cm^2 of gold, thorium, and uranium respectively.

* In general, the number of tritons per incident particle is

$$t_i = 6.02 \times 10^{-7} \frac{\beta}{A. W.} \sum_0^{x(\text{max.})} \sigma_x,$$

where β is the target density in $\text{mg}/\text{cm}^2\text{-mm}$, A. W. is the atomic weight, and σ_x is the cross section in millibarns for foil x.

IV. DISCUSSION

The striking result that the apparent cross section for the yield of tritium in the (α, t) reaction reaches a peak in target foils that the beam does not reach, leads, on detailed examination of the data, to the conclusion that tritons with velocities comparable with those of the helium ions incident on the foil stack must be emitted in forward directions. The high energy of the tritons (20-30 Mev from an analysis of Fig. 1A) indicates that for a large share of interactions the residual heavy product must be left in the ground state or in a low-lying excited state, and hence little fission competition is possible for U^{238} and Th^{232} . This conclusion that these nuclei survive fission is given confirmation by the agreement seen in Table I between the cross sections for the production of the " $(\alpha, p2n)$ " product determined by radiochemical methods which measure only the events surviving fission, and the tritium-production cross sections. The fact that the integrated cross sections for tritium production are somewhat higher than the corresponding $(\alpha, p2n$ or $t)$ cross sections determined radiochemically may indicate the presence of a spectrum of triton energies of which approximately 20% results in residual nuclei being left in states sufficiently excited to undergo fission. The differences as they stand, however, are such that it is possible that the two types of cross sections are actually equal. It should be noted that although it is believed that the tritium activity collected and the corresponding heavy fragments observed radiochemically in the main represent simply (α, t) reactions, the tritium actually may result from (α, t) , (α, tn) , and (α, tf) reactions and the " $(\alpha, p2n)$ " products may result from (α, t) , $(\alpha, p2n)$, and (α, dn) reactions. For this reason and also because of the fact that the experimental procedures are so different, one must exercise caution in interpreting the data.

A direct interaction mechanism such as stripping of a proton from the helium ion is suggested by experimental data presented here. If the residual nucleus is to be left in an energy state between ground and the fission threshold, the tritons must carry off an energy of between $(48 + Q)$ Mev and $(48 + Q - 5)$ Mev, or approximately $(35 - 30)$ Mev in the case of full-energy helium ions. If tritium production is assumed most predominant in the first few mils of foil, that is, with helium-ion energies greater than 30 Mev as shown by the radiochemical data, then the energy of the tritons found deep in the stacked

foils can be calculated without too much uncertainty. The results show that these tritons, having an angular dependence strongly favored in the direction of the helium ion beam, have energies of 30 ± 5 Mev. Further, analysis of the data of Fig. 1-A shows that if the majority of tritons have energies between 25 - 35 Mev they are emitted within a cone of total included angle of 60° .

The data for triton production by deuteron projectiles appear reasonable in the light of a mechanism by which the deuteron "picks up" a neutron as it passes near the target nucleus. Harvey¹¹ has studied the excited states of the residual nuclei produced from this reaction in Au¹⁹⁷ and found the production of nuclei in their ground state quite prevalent. Table I shows that the cross sections for triton production in (d,t) reactions are actually considerably larger in Th²³² and U²³⁸ than in Au¹⁹⁷ despite the fissionability of the former two. The nature of the experiments is such that the high energy of the tritons is not as obvious in deuteron bombardments, since the range of the deuterons is greater than the range of the maximum-energy tritons so that no tritium is found in foils beyond the deuteron range.

Cross sections for triton production in (p,t) reactions are similar in all three of the nuclides although the integrated cross sections are slightly larger in the case of Th²³² and U²³⁸. Thus, the magnitudes of the cross sections again reflect a lack of fission competition with the (p,t) reaction for Th²³² and U²³⁸. This reaction is a possible example of a pick-up of two nucleons (double pick-up). Cohen¹⁰ has studied the (p,t) reaction in beryllium and iron and considers it to be a direct interaction process. Again the range of the tritons is less than that of the protons, and it is not possible to determine the range of the tritons by the present simple experimental techniques.

Thus, it has been established in a qualitative way that high-energy tritium emission is occurring to an important extent in heavy fissionable and non-fissionable isotopes; that most, if not all, of the production of " $(\alpha, p2n)$ " products is accounted for by the (α, t) process; and that tritium emission reactions proceed at these energies through a mechanism not involving formation of a compound nucleus. The (α, t) reaction probably proceeds by a stripping mechanism with a high-energy triton emergent in the forward direction. On the basis of their large production cross sections the tritons from the (d,t) and (p,t) reactions would also appear to be generated by a mechanism circumventing compound-nucleus formation, and probably represent examples of "pick-up" reac-

tions. It is interesting to note that cross-section values for the helium-ion, deuteron, and proton reactions are in the order $(\alpha,t) < (p,t) < (d,t)$, which is the generally accepted order of increasing ease of the processes, helium-ion stripping, double neutron pick-up, single neutron pick-up.

At the present time, independent determinations of the range and angular distribution of tritons produced in these various reactions are under investigation in this laboratory and will be reported shortly.

This project was performed under the auspices of the U.S. Atomic Energy Commission. Appreciation is expressed to the crews of the 60-inch cyclotron and the linear accelerator for their assistance.

Table I. Integrated Triton Production Cross Sections

Reaction	Isotope	Integral σ		Radiochemical value
		mb-mm	Tritons/incident particle	
(α ,t)	Au ¹⁹⁷	2.17	$1.28 \pm 0.11 \times 10^{-5}$	--
	Th ²³²	5.15	$1.56 \pm 0.13 \times 10^{-5}$	1.27×10^{-5} *
	U ²³⁸	2.60	$1.23 \pm 0.09 \times 10^{-5}$	1.08×10^{-5} **
(d,t)	Au ¹⁹⁷	8.55	$5.04 \pm 0.45 \times 10^{-5}$	--
	Th ²³²	37.4	$11.4 \pm 1.0 \times 10^{-5}$	--
	U ²³⁸	20.0	$9.45 \pm 0.75 \times 10^{-5}$	--
(p,t)	Au ¹⁹⁷	6.84	$4.04 \pm 0.52 \times 10^{-5}$	--
	Th ²³²	24.2	$6.15 \pm 0.70 \times 10^{-5}$	--
	U ²³⁸	13.7	$5.81 \pm 0.62 \times 10^{-5}$	--

* Assuming a counting efficiency of 100%.

** Assuming a counting efficiency of 70%.

REFERENCES

1. R. A. Glass, R. J. Carr, J. W. Cobble, and G. T. Seaborg, Phys. Rev. 104, 434 (1956).
2. S. Vandenbosch, unpublished work.
3. R. Vandenbosch, T. D. Thomas, R. A. Glass, and G. T. Seaborg, to be published.
4. B. M. Foreman, unpublished work.
5. L. E. Bailey, (thesis) University of California, 1956; also published as University of California Radiation Laboratory Report, UCRL-3334 (1956).
6. W. H. Barkas and H. Tyren, Phys. Rev. 89, 1 (1953).
7. R. Deutsch, Phys. Rev. 97, 1110 (1955).
8. L. A. Currie, W. F. Libby, and R. L. Wolfgang, Phys. Rev. 101, 1557 (1956).
9. J. R. Hoff and T. N. Marsham, Proc. Phys. Soc. (London) 66A, 1032 (1953).
10. B. L. Cohen and T. H. Handley, Phys. Rev. 93, 514 (1954).
11. J. A. Harvey, Can. J. Phys. 31, 278 (1953).
12. S. T. Butler, Proc. Roy. Soc. 208A, 559 (1951) (London).
13. H. C. Newns, Proc. Roy. Soc. 65A, 916 (1952) (London).

FIGURE CAPTION

Fig. 1. Cross section for triton production in Au^{197} , Th^{232} , and U^{238} induced by helium ions, deuterons, and protons. All abscissas are expressed in mg/cm^2 of target material with numerical scales as given below Sections 1A, 1B, and 1C. The ordinates are in units of millibarns with the individual numerical scales shown for each of the nine curves.

Fig. 1A, Cross sections for triton production induced by helium ions;

Fig. 1B, Cross sections for triton production induced by deuterons;

Fig. 1C, Cross sections for triton production induced by protons.

