

UC Riverside

Journal of Citrus Pathology

Title

Monitoring of Candidatus Liberibacter asiaticus in Citrus Seedlings at Greenhouse Conditions and Commercial Orchards of Sweet Orange and Tahiti Lime in the Northwest of Parana State, Brazil

Permalink

<https://escholarship.org/uc/item/5p00v93w>

Journal

Journal of Citrus Pathology, 1(1)

Authors

Sauer, A. V.
Barbieri, B. R.
Coletta-Filho, H. D.
et al.

Publication Date

2014

DOI

10.5070/C411025187

Copyright Information

Copyright 2014 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

8.15 P

Monitoring of *Candidatus Liberibacter asiaticus* in Citrus Seedlings at Greenhouse Conditions and Commercial Orchards of Sweet Orange and Tahiti Lime in the Northwest of Parana State, Brazil

Sauer, A.V.¹, Barbieri, B.R.¹, Coletta-Filho, H.D.², Machado, M.A.², Corazza, M.J.¹, and Nunes, W.M.C.¹

¹Nucleo de Pesquisa em Biotecnologia Aplicada-NBA, Universidade Estadual de Maringa - UEM, Maringa-PR, Brazil

²Centro de Citricultura Sylvio Moreira, Cordeiropolis, SP, Brazil

This study aimed to monitor the behavior of *Ca. Liberibacter asiaticus* in grafted seedlings in the greenhouse and in commercial orchards of sweet orange and Tahiti lime. Plants of sweet orange and Tahiti lime naturally infected with HLB were protected with screens aphid-proof to prevent the spread of disease. In greenhouse we used 17 Pêra variety seedlings that were inoculated in 2008 with infected budwood. The detection of HLB in the plants was carried out using conventional PCR. Ten leaves of each plant were collected for DNA extraction and for their full monitoring of the bacterial population by quantitative PCR for a period of 19 months. We observed erratic behavior of the bacterium. Even after the bacterium detection in the plant it was not possible to verify the presence of the etiologic agent in the same place after a few months. In the seedlings the bacterium was not detected in 82.3% of the plants in certain months, coinciding with the warmer months of the year in Brazil, despite its presence has being detected at earlier dates. For the plants in commercial orchards, the detected Ct values were between 18 and 33. The pathogen was even found when the expression of symptoms was small. The study of the population behavior of this agent contributes to the understanding of the epidemiology of this disease.

Support: CNPq, Capes, Araucaria Foundation

Topic Categories: Survey, Detection and Diagnosis